


Title	Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts
Authors	Payelleville, Amaury;Blackburn, Dana;Lanois, Anne;Pagès, Sylvie;Cambon, Marine C.;Ginibre, Nadege;Clarke, David J.;Givaudan, Alain;Brillard, Julien
Publication date	2019-10-09
Original Citation	Payelleville, A., Blackburn, D., Lanois, A., Pagès, S., Cambon, M. C., Ginibre, N., Clarke, D. J., Givaudan, A. and Brillard, J. [2019] 'Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts', PLOS ONE, 14(10), e0212655. (14pp.) DOI: 10.1371/journal.pone.0212655
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0212655 - 10.1371/journal.pone.0212655
Rights	©2019 Payelleville et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited - http://creativecommons.org/licenses/by/4.0/
Download date	2023-06-06 10:07:52
Item downloaded from	http://hdl.handle.net/10468/8872



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Fig. S1



Relative Expression Results

Parameter	Value
Iterations	2000

Gene	Type	Reaction Efficiency	Expression	Std. Error	95% C.I.	P(H1)	Result
GyrB	REF	0,7929	1,157				
Dam	TRG	1,1132	14,217	8,224 - 23,010	5,785 - 41,381	0,000	UP
fis	REF	0,7791	0,864				

Interpretation

Dam is UP-regulated in sample group (in comparison to control group) by a mean factor of 14,217 (S.E. range is 8,224 - 23,010).
Dam sample group is different to control group. P(H1)=0,000

Figure S1. Relative expression of the *dam* gene in the *P. luminescens* *dam*-overexpressing strain.

qRT-PCR was carried out with total RNA extracted from 3 independent samples of exponentially growing cells of the *dam*-overexpressing strain and of the control strain. Left panel, Box-plot representing the expression ratio between the two strains is represented for the *dam* gene. Right panel, detailed report of the REST-analysis using with *gyrB* and *fis* used as control genes (see Materials & Methods section for details). The level of expression between the 2 strains was different at $p < 0.001$.