

|                             |  |
|-----------------------------|--|
| Title                       | Substance use in young persons in Ireland, a systematic review   |
| Authors                     | Murphy, Kevin D.;Sahm, Laura J.;Lambert, Sharon;Byrne, Stephen   |
| Publication date            | 2012-10-29   |
| Original Citation           | Murphy K., Sahm L., Lambert S., Byrne S. (2012) 'Substance use in young persons in Ireland, a systematic review', European Society of Clinical Pharmacy (ESCP) 41st Symposium on Clinical Pharmacy, Barcelona, Spain, 29 - 31 October, in International Journal of Clinical Pharmacy (2013) 35(5), pp. 997-998. doi: 10.1007/s11096-013-9801-0 |
| Type of publication         | Conference item  |
| Link to publisher's version | <a href="http://www.escpweb.org/">http://www.escpweb.org/</a> - 10.1007/s11096-013-9801-0  |
| Rights                      | © Springer Science+Business Media Dordrecht 2013   |
| Download date               | 2023-09-27 22:37:06  |
| Item downloaded from        | <a href="https://hdl.handle.net/10468/6204">https://hdl.handle.net/10468/6204</a>  |



# UCC

**University College Cork, Ireland**  
 Coláiste na hOllscoile Corcaigh



# Substance Use in Young Persons in Ireland, a Review

Kevin D Murphy<sup>1</sup>, Laura Sahn<sup>1</sup>, Sharon Lambert<sup>2</sup>, and Stephen Byrne<sup>1</sup>.

<sup>1</sup>Pharmaceutical Care Research Group, University College Cork, Ireland <sup>2</sup>Matt Talbot Services, Cork, Ireland



## Background:

- 22.5% of all deaths in the European region were directly caused by addictive substances<sup>1</sup>.
- Adolescence is a time when people try many addictive substances for the first time<sup>2</sup>.
- Experimenting with addictive substances at an early age increases the risk of polysubstance use<sup>3</sup>, problem drug use<sup>4</sup>, health<sup>5</sup> and psychological<sup>6</sup> problems.

## Objective:

- The objective of this study was to conduct a literature review with special emphasis on the prevalence of adolescent use of four commonly used substances in Ireland: tobacco, alcohol, cannabis, and benzodiazepines<sup>7</sup>.

## Design:

- The review followed the Preferred Reporting Items for Systematic reviews and Meta-analyses (PRISMA) guidelines.
- Boolean searches were conducted on the following databases: PubMed, Scopus, Web of Knowledge, Google Scholar, ERIC, EMBASE, and CINAHL.
- Search terms used were *adolescent, cannabis, tobacco, benzodiazepines, and Ireland*.
- A manual search of the Health Research Board drug-related website, National Documentation Centre on Drug Use was conducted.
- Inclusion criteria were English language articles, and articles published from January 2000.

## Results:


- The study selection process is detailed in Figure 1.
- The search returned 18 studies; 1 randomised control trial and 17 cross-sectional studies.
- Eight studies looked at a single substance, while there was a single substance that looked at 2 substances. Five studies looked at 3 substances and four of the studies looked at all 4 substances.
- There were 11 studies that surveyed a national population, while the other 7 studies sampled at provincial or county level.

## Tobacco

- There were sixteen studies that examined tobacco use.
- Over the timeframe of this systematic review, there was a fall of lifetime usage levels from approximately 60% to 48%.
- A fall was also found in the percentage that smoked in the previous month from approximately 30% to 10-20%.

## Alcohol

- There were eleven studies that examined alcohol use.
- Lifetime use of alcohol fell from approximately 90% to 60%, though some of the studies included in this section did not have rigorous design.



**Figure 1. Study selection process using PRISMA diagram**

- Early initiation of alcohol (prior to 13 years of age) levels fell by approximately 10% to 25%.

## Cannabis

- There were nine studies that examined cannabis use.
- Previous 12-month usage fell from approximately 30% around 2002, to about 15% after 2009.
- Recent use of cannabis (use in the previous month) also fell from approximately 15% to 10%.

## Benzodiazepines

- There were six studies that examined benzodiazepine use.
- Lifetime use of benzodiazepines remained constant at approximately 10%.
- Some studies differentiated between prescription use and non-prescription use. The latter was low but steady at 2.0-5.6%.

## Conclusions:

- This review has shown that despite some decline, substance use is still quite prevalent within the adolescent population of Ireland.
- Continued tracking of substance use levels will elucidate whether this trend is short-term or long-term.
- Due to the serious consequences for both the substance user and for society as a whole, future work will examine the reasons underlying substance use.

## Limitations:

- 13 of the 18 studies conducted their surveys on school children. This omits early school-leavers who are more likely to use both legal and illegal substances<sup>8</sup>.
- The questions asked in the studies varied, so only an approximate comparison was possible.