

Title	Dissecting the regulation of bile-induced biofilm formation in <i>Staphylococcus aureus</i>
Authors	Ulluwishewa, Dulantha;Wang, Liang;Pereira, Callen;Flynn, Stephanie;Cain, Elizabeth;Stick, Stephen;Reen, F. Jerry;Ramsay, Joshua P.;O'Gara, Fergal
Publication date	2016-08-01
Original Citation	Ulluwishewa, D., Wang, L., Pereira, C., Flynn, S., Cain, E., Stick, S., Reen, F. J., Ramsay, J. P. and O'Gara, F. (2016) 'Dissecting the regulation of bile-induced biofilm formation in <i>Staphylococcus aureus</i> ', <i>Microbiology</i> , 162(8), pp. 1398-1406. doi: 10.1099/mic.0.000317
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="https://www.microbiologyresearch.org/content/journal/micro/10.1099/mic.0.000317">https://www.microbiologyresearch.org/content/journal/micro/10.1099/mic.0.000317</a> - <a href="https://doi.org/10.1099/mic.0.000317">10.1099/mic.0.000317</a>
Rights	© The Authors 2016. The definitive peer reviewed, edited version of this article is published in <i>Microbiology</i> , 162, 8, 2016, <a href="https://doi.org/10.1099/mic.0.000317">https://doi.org/10.1099/mic.0.000317</a>
Download date	2024-06-19 01:51:45
Item downloaded from	<a href="https://hdl.handle.net/10468/8865">https://hdl.handle.net/10468/8865</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

# Microbiology

## Dissecting the regulation of bile-induced biofilm formation in *Staphylococcus aureus*

--Manuscript Draft--

<b>Manuscript Number:</b>	MIC-D-16-00108
<b>Full Title:</b>	Dissecting the regulation of bile-induced biofilm formation in <i>Staphylococcus aureus</i>
<b>Short Title:</b>	Bile induced Biofilm formation in <i>Staphylococcus</i>
<b>Article Type:</b>	Standard
<b>Section/Category:</b>	Host-microbe interaction
<b>Keywords:</b>	bile; biofilm; <i>Staphylococcus</i> ; gastro-esophageal reflux
<b>Corresponding Author:</b>	Fergal O'Gara Biomerit Research Centre, University College Cork Cork, IRELAND
<b>First Author:</b>	Dulantha Ulluwishewa
<b>Order of Authors:</b>	Dulantha Ulluwishewa Liang Wang Callen Pereira Stephanie Flynn Stephen Stick F.Jerry Reen Joshua P Ramsay Fergal O'Gara
<b>Manuscript Region of Origin:</b>	AUSTRALIA
<b>Abstract:</b>	<p>Aspiration of bile into the cystic fibrosis (CF) lung has emerged as a prognostic factor for reduced microbial lung biodiversity and the establishment of often fatal, chronic pathogen infections. <i>Staphylococcus aureus</i> is one of the earliest pathogens detected in the lungs of children with CF, and once established as a chronic infection, strategies for its eradication become limited. Several lung pathogens are stimulated to produce biofilms in vitro in the presence of bile. In this study, we further investigated the effects of bile on <i>S. aureus</i> biofilm formation. Most clinical <i>S. aureus</i> strains and the laboratory strain RN4220 were stimulated to form biofilms with sub-inhibitory concentrations of bile. Additionally, we observed bile-induced sensitivity to aminoglycosides, which we exploited in a bursa aurealis transposon screen to isolate mutants reduced in aminoglycoside sensitivity and augmented in bile-induced biofilm formation. We identified five mutants that exhibited hypersensitivity to bile with respect to bile-induced biofilm formation, three of which carried transposon insertions within gene clusters involved in wall teichoic acid (WTA) biosynthesis or transport. Strain TM4 carried an insertion between the divergently oriented tagH-tagG genes, encoding the putative WTA membrane translocation apparatus. Ectopic expression of tagG in TM4 restored a wild-type bile-induced biofilm response, suggesting that reduced translocation of WTA in TM4 induced sensitivity to bile and enhanced bile-induced biofilm formation. We propose that WTA may be important for protecting <i>S. aureus</i> against exposure to bile and that bile-induced biofilm formation may be an evolved response to protect cells from bile-induced cell lysis.</p>



1 TITLE: Dissecting the regulation of bile-induced biofilm formation in *Staphylococcus aureus*

2

3 RUNNING TITLE: Bile induced Biofilm formation in *Staphylococcus*

4

5 AUTHORS AND AFFILIATIONS:

6 Dulantha Ulluwishewa<sup>1δ</sup>, Liang Wang<sup>1δ\*</sup>, Callen Pereira<sup>1</sup>, Stephanie Flynn<sup>2</sup>, Stephen Stick<sup>3</sup>, F.

7 Jerry Reen<sup>2</sup>, Joshua P. Ramsay<sup>1</sup>, and Fergal O’Gara<sup>1,2</sup>

8 <sup>1</sup> School of Biomedical Sciences, CHIRI Research Institute, Curtin University, Perth, Australia

9 <sup>2</sup>BIOMERIT Research Centre, School of Microbiology, University College Cork, Cork,

10 Ireland

11 <sup>3</sup>Telethon Kids Institute, University of Western Australia, Australia; Princess Margaret

12 Hospital for Children, Western Australia, Australia; School of Paediatric and Child Health,

13 University of Western Australia, Australia.

14 \*Present address: School of Medical Informatics, Xuzhou Medical University, Xuzhou,

15 Jiangsu Province, China

16 <sup>δ</sup>These authors contributed equally to this paper

17

18 CORRESPONDING AUTHOR DETAILS:

19 Professor Fergal O’Gara

20 +618 9266 9447

21 Fergal.Ogara@curtin.edu.au

22

23 KEYWORDS: bile, biofilm, *Staphylococcus*, gastro-esophageal reflux

24 SUBJECT CATEGORY: Host-microbe interaction, Regulation

25 WORD COUNT: 4522

## 26 **ABSTRACT**

27 Aspiration of bile into the cystic fibrosis (CF) lung has emerged as a prognostic factor for  
28 reduced microbial lung biodiversity and the establishment of often fatal, chronic pathogen  
29 infections. *Staphylococcus aureus* is one of the earliest pathogens detected in the lungs of  
30 children with CF, and once established as a chronic infection, strategies for its eradication  
31 become limited. Several lung pathogens are stimulated to produce biofilms *in vitro* in the  
32 presence of bile. In this study, we further investigated the effects of bile on *S. aureus* biofilm  
33 formation. Most clinical *S. aureus* strains and the laboratory strain RN4220 were stimulated to  
34 form biofilms with sub-inhibitory concentrations of bile. Additionally, we observed bile-  
35 induced sensitivity to aminoglycosides, which we exploited in a *bursa aurealis* transposon  
36 screen to isolate mutants reduced in aminoglycoside sensitivity and augmented in bile-induced  
37 biofilm formation. We identified five mutants that exhibited hypersensitivity to bile with  
38 respect to bile-induced biofilm formation, three of which carried transposon insertions within  
39 gene clusters involved in wall teichoic acid (WTA) biosynthesis or transport. Strain TM4  
40 carried an insertion between the divergently oriented *tagH-tagG* genes, encoding the putative  
41 WTA membrane translocation apparatus. Ectopic expression of *tagG* in TM4 restored a wild-  
42 type bile-induced biofilm response, suggesting that reduced translocation of WTA in TM4  
43 induced sensitivity to bile and enhanced bile-induced biofilm formation. We propose that WTA  
44 may be important for protecting *S. aureus* against exposure to bile and that bile-induced biofilm  
45 formation may be an evolved response to protect cells from bile-induced cell lysis.

## 46 **INTRODUCTION**

47 Chronic infections by pathogens such as *Pseudomonas aeruginosa* and *Staphylococcus aureus*  
48 are a leading cause of morbidity and mortality in cystic fibrosis (CF) patients. Chronic bacterial  
49 infections can rarely be eradicated by antimicrobial treatment, and thus chronic infection of the  
50 lung can eventually lead to a fatal decline in lung function (Furukawa *et al.*, 2006). This is

51 becoming an ever increasing clinical issue where many have predicted the onset of a post-  
52 antibiotic era (Cooper & Shlaes, 2011). Therefore, innovative and alternative strategies are  
53 urgently needed, away from the classical antibiotic approach. The refractory nature of chronic  
54 infections to conventional therapies is largely attributed to bacteria adopting a biofilm-like  
55 mode of growth. Microorganisms in biofilms are embedded within a matrix of extracellular  
56 polymeric substances which provides a barrier against host immune defenses and antimicrobial  
57 therapy (Flemming & Wingender, 2010). The biofilm matrix is composed of polysaccharides,  
58 proteins, lipids, and genomic DNA that is released by lysed resident bacterial cells. While the  
59 molecular mechanisms underlying the formation of biofilms in a broad spectrum of pathogens  
60 has been well studied, until recently the molecular triggers which cause lung-colonizers to  
61 adopt a chronic lifestyle and the associated biofilm mode of growth remained largely  
62 uncharacterized. Understanding how bacteria adopt this chronic lifestyle in the lung would  
63 provide new therapeutic options for prevention and treatment.

64         The aspiration of bile acids into the lungs arising from gastro-esophageal reflux disease  
65 (GERD) has since emerged as a host-trigger of chronic bacterial infection (Reen *et al.*, 2012;  
66 Reen *et al.*, 2014) and chronic inflammation (Legendre *et al.*, 2014), particularly in CF, where  
67 up to 40% of children and 80% of adult patients can suffer from this complication (Legendre  
68 *et al.*, 2014; Pauwels *et al.*, 2012; Reen *et al.*, 2012; Reen *et al.*, 2014; Stringer *et al.*, 1988).  
69 Indeed, the incidence may be underestimated as clinical diagnosis of GER is often not sufficient  
70 to determine bile aspiration. Bile acids have been detected in the sputum and bronchoalveolar  
71 lavage fluid of patients that do not present with classical GER symptoms. This ‘silent  
72 aspiration’ phenomenon is particularly severe in CF patients and highlights the urgent unmet  
73 need for rapid diagnosis of bile acid profiles in biological samples from respiratory and lung  
74 transplant patients (Button *et al.*, 2005). Over recent years a number of research publications  
75 have suggested that acid-and non-acid reflux may negatively influence the progression of

76 respiratory disease (D'Ovidio *et al.*, 2005a; D'Ovidio *et al.*, 2005b; el-Serag & Sonnenberg,  
77 1997; Pauwels *et al.*, 2012; Perng *et al.*, 2007; Wu, 2008; Wu *et al.*, 2009). el-Serag and  
78 Sonnenberg showed that patients with erosive esophagitis, a sign of significant GER, had  
79 increased incidence of pulmonary fibrosis, chronic bronchitis or chronic obstructive pulmonary  
80 disease in a case control study of more than 200,000 patients (el-Serag & Sonnenberg, 1997).  
81 A strong correlation between GER-derived reflux, pulmonary aspiration, and increased lung  
82 damage also extends to several other respiratory diseases (Navarro *et al.*, 2001), including  
83 idiopathic pulmonary disease and advanced lung damage arising from lung transplantation  
84 (Sweet *et al.*, 2006; Sweet *et al.*, 2007a; Sweet *et al.*, 2007b), ventilator associated pneumonia  
85 (Wu *et al.*, 2009), Barrett's esophagus and esophageal adenocarcinoma (Nassr *et al.*, 2011) and  
86 Bile Acid Pneumonia in neonates (Zecca *et al.*, 2004; Zecca *et al.*, 2008). Therefore, the  
87 implications of elucidating the link between bile aspiration and chronic pathogen behavior has  
88 consequences for a range of clinical conditions.

89         Apart from an association between GERD and increased colonization by *P. aeruginosa*  
90 and *S. aureus* (Palm *et al.*, 2012; van der Doef *et al.*, 2009), bile has been shown to influence  
91 the behavior of *P. aeruginosa* and other respiratory pathogens *in vitro*, suppressing phenotypes  
92 associated with acute infection, while up-regulating phenotypes associated with chronic  
93 infection, including biofilm formation (Reen *et al.*, 2012). Microbial diversity has also shown  
94 to be significantly reduced in bile-aspirating patients compared with non-aspirating patients,  
95 suggesting that aspirated bile may be a major factor in shaping the pervasive lung microbiota  
96 signature in CF patients (Blainey *et al.*, 2012; Reen *et al.*, 2014). Furthermore, bile acids have  
97 been shown to trigger the IL-6 pro-inflammatory cytokine *in vitro*, suppressing HIF-1 signaling  
98 in *P. aeruginosa* infected cells (Legendre *et al.*, 2014), while reflux and aspiration have been  
99 shown to correlate with increased levels of cytokines and neutrophils *in vivo* (D'Ovidio *et al.*,  
100 2005a). Therefore, the possibility that the aspiration of bile into the lungs of pediatric patients

101 must be considered as a potential underlying factor in the emergence of chronic microbial  
102 infections.

103         In this study we examined the effect of bile on virulence related behavior in *S. aureus*,  
104 the primary pathogen associated with early stage CF infection. While *S. aureus* is considered  
105 a commensal bacterium, as it is a common colonizer of the human skin and respiratory tract, it  
106 is also a frequent cause of clinically important infections (Wertheim *et al.*, 2005). In many  
107 cases *S. aureus* is the earliest colonizer in CF patients, and is the most prevalent CF pathogen  
108 in children and adolescents (Kahl, 2010; Souza *et al.*, 2006). First elucidating the impact of  
109 physiologically relevant concentrations of bile exposure on this important pediatric pathogen,  
110 we focused on antibiotic tolerance and biofilm formation. In order to probe more deeply the  
111 regulatory mechanisms governing the bile-mediated biofilm response, we utilized a random  
112 transposon mutagenesis screen to isolate *S. aureus* mutants with an altered bile response. This  
113 uncovered a previously unforeseen switch to a biofilm lifestyle by *S. aureus* in the presence of  
114 bile, with some intriguing insights into the molecular mechanism underpinning this key  
115 pathogenic determinant.

## 116 **MATERIALS AND METHODS**

### 117 **Bacterial strains, plasmids, and growth conditions**

118 The bacterial strains and plasmids used in this study are outlined in Table 1. *S. aureus* strains  
119 were cultured at 37°C in Tryptic Soy Broth (TSB; Becton Dickinson) or Tryptic Soy Agar  
120 (TSA; TSB containing 1.5% (w/v) agar). *Escherichia coli* EPI300 was used as a cloning host,  
121 and was cultured at 37°C in Lysogeny Broth (LB), or LB agar. Where required, media were  
122 supplemented with the following concentrations of antibiotics unless otherwise stated: 100  
123 µg/mL ampicillin, 10 µg/mL chloramphenicol, 2.5 µg/mL tetracycline, 10 µg/mL  
124 erythromycin. Media were supplemented with bovine bile (Sigma-Aldrich), sodium cholate

125 (SC; Sigma-Aldrich), sodium deoxycholate (SDC; Sigma-Aldrich), or sodium dodecyl sulfate  
126 (SDS; Amresco) when required. Stock solutions of bile, SC and SDC were prepared in  
127 deionized distilled water and filter sterilized prior to addition to media.

#### 128 **DNA manipulations**

129 *S. aureus* DNA was extracted from broth using the FavorPrep Blood / Cultured Cell Genomic  
130 DNA Extraction Mini Kit (Favorgen Biotech Corp) following cell lysis with lysostaphin  
131 (Sigma-Aldrich) unless otherwise stated. Electroporation of *S. aureus* strains was carried out  
132 using the method described by Schenk & Laddaga (1992). PCR products were amplified using  
133 Phusion High-Fidelity DNA Polymerase (New England BioLabs), and purified using the  
134 FavorPrep GEL/PCR Purification Kit (Favorgen Biotech Corp).

#### 135 **Biofilm attachment assay**

136 Stationary-phase cultures of *S. aureus* cultures were diluted 1:200 in TSB or TSB  
137 supplemented with appropriate treatment, and aliquots transferred to 96-well or 24-well plates.  
138 Following incubation at 37°C, the wells were washed twice in water to remove planktonic cells.  
139 Attached cells (biofilm) were stained with 0.1% (w/v) crystal violet solution, washed twice to  
140 remove unincorporated stain, and solubilized with acetone:ethanol (3:7) before quantification  
141 by measuring the absorbance at 595 nm. At least three independent biological replicates were  
142 performed for each experiment. For each strain of *S. aureus*, treatment samples were compared  
143 to untreated samples using a two-tailed paired or unpaired Student's t-test.

#### 144 **Growth analysis**

145 To determine the effects of bile on the growth of *S. aureus*, stationary-phase cultures of *S.*  
146 *aureus* were diluted 1:500 in 25 mL TSB or TSB supplemented with bovine bile. Cultures were  
147 grown at 37°C with agitation at 180 rpm and samples were taken over a 24 hour period and the  
148 optical density (OD) measured at a wavelength of 600 nm.



149 **Antibiotic sensitivity determinations**

150 For disc diffusion testing, a single colony of *S. aureus* RN4220 was resuspended in 1 mL of  
151 phosphate buffered saline, and evenly spread on TSA plates or TSA plates containing 0.3%  
152 bile to prepare a lawn culture. Antibiotic discs, containing gentamicin (10 µg),  
153 chloramphenicol (30 µg), ampicillin (10 µg), erythromycin (15 µg), or tetracycline (30 µg),  
154 were placed on the agar, and the plates were incubated at 37°C for 18 hours, following which  
155 the diameter of the inhibition zone was observed.

156 The minimum inhibitory concentrations (MICs) of the aminoglycosides gentamycin,  
157 streptomycin, neomycin, and kanamycin were determined in duplicate by broth macrodilution  
158 method. Antibiotic solutions (0.12, 0.25, 0.5, 1, 2, 4, 8, 16, 32, 64 µg/mL) were prepared in  
159 TSB or TSB containing 1 mM SC. An inoculum of *S. aureus* RN4220 was prepared by saline  
160 (0.85% NaCl) suspension of isolated colonies, adjusted to achieve a turbidity equivalent to a  
161 0.5 McFarland standard, and diluted 1:300 in TSB containing 1 mM SC and/or antibiotics such  
162 that the final inoculum was  $\sim 5 \times 10^5$  colony-forming units (CFU)/mL. Cultures were incubated  
163 overnight at 37°C with shaking. The lowest concentration of antibiotic which inhibited all  
164 visually apparent growth was considered the MIC.

165 **Transposon mutagenesis**

166 Plasmids pBursa and pFA545 were transformed together into *S. aureus* RN4220 by  
167 electroporation, and transformants were selected on TSA containing chloramphenicol and  
168 tetracycline following incubation at the permissive temperature of 30°C for 48 hours. Resulting  
169 colonies were resuspended in sterile water and incubated at the non-permissive temperature of  
170 43°C for 1 hour before spreading on selective media (TSA containing 1 erythromycin and SDC,  
171 or TSA containing erythromycin, SC, and either gentamycin, neomycin, or kanamycin) to  
172 screen for mutants of interest.

173 **Determination of site of transposon insertion**

174 Transposon insertion sites were determined by random-primed PCR as described below.  
175 Oligonucleotide primers are listed in Table 2. DNA was purified from transposon mutants  
176 using PrepMan Ultra Sample Preparation Reagent (Life Technologies) following cell lysis with  
177 lysostaphin. PCR was performed using the purified DNA as template, with a random primer  
178 mix (PF106, PF107 and PF108) and a transposon-specific primer (GFP1). A second PCR was  
179 performed using an aliquot of amplified product from the first PCR, with primers GFP2 and  
180 PF109. Following confirmation of the presence of amplified product by gel electrophoresis,  
181 the PCR reactions were purified and subsequently sequenced using primer GFP3. Sequencing  
182 reactions were conducted by the Australian Genome Research Facility. Each sequence was  
183 compared using the National Center for Biotechnology Information (NCBI) nucleotide Basic  
184 Local Alignment Search (BLASTN) tool to the *S. aureus* NCTC8325 complete genome  
185 (CP000253.1). The location at which the query sequence first matched the subject sequence  
186 was determined as the transposon insertion site.

187 **RESULTS**

188 **Bile induces biofilm formation in most *S. aureus***

189 Physiological concentrations of bile or bile acids stimulate *in vitro* biofilm formation in several  
190 lung-colonizing pathogens (Reen *et al.*, 2012). In contrast, bile suppresses biofilm formation  
191 by *S. aureus* strain NCDO949, a common laboratory strain originally isolated from pleural  
192 fluid. To establish whether NCDO949 bile-response phenotype was representative of *S. aureus*  
193 strains, biofilm formation was analyzed for pediatric isolates obtained from the CF unit at Cork  
194 University Hospital, Ireland, using the crystal violet attachment assay. In contrast to  
195 NCDO949, biofilm formation was stimulated by bile in these isolates (Fig. 1a). Furthermore,  
196 bile-stimulated biofilm formation was observed for community-acquired MRSA strains

197 JKD6159 and MW2 and the common laboratory strain RN4220 (Fig. 1b). In contrast, USA300  
198 strain JE2 did not exhibit bile-induced biofilm formation. Therefore these data suggest like  
199 other lung pathogens, most *S. aureus* strains are stimulated to form biofilms in the presence of  
200 bile, but, that there is variation in this response amongst strains.

201         Since the well-characterized and genetically tractable *S. aureus* strain RN4220  
202 exhibited a similar response to bile as most clinical *S. aureus* isolates, we further investigated  
203 the effects of the addition of bile on this strain. Analysis of RN4220 growth in TSB broth  
204 culture revealed that the log-phase growth rate was uninhibited by the addition of up to 0.3%  
205 bovine bile (Fig. 1c). Bile acids make up over 50 % of the total solute concentration of bile  
206 (Kristiansen *et al.*, 2007) and their salt equivalents have been implicated in biofilm formation  
207 in *P. aeruginosa* (Reen *et al.*, 2012). The addition of sodium cholate (SC) (Fig. 2a) or sodium  
208 deoxycholate (SDC) (Fig. 2b) at sub-inhibitory concentrations resulted in a dose dependent  
209 increase in *S. aureus* biofilm formation, indicating that these individual bile components were  
210 able to induce a similar response to whole bile in *S. aureus*. Interestingly, addition of 0.1 mM  
211 of the anionic detergent SDS also causes a similar but statistically insignificant increase in  
212 biofilm formation, possibly indicating that the biofilm formation by *S. aureus* may be a  
213 response to the common detergent activities of these molecules (Fig. 2c).

#### 214 **Development of a *bursa aurealis* transposon mutagenesis screen to isolate bile-response** 215 **mutants**

216 Bile acids have been demonstrated to enhance the activity of penicillin and neomycin against  
217 staphylococcal strains. Bile has no effect on the efficacy of other antibiotics such as  
218 chloramphenicol or erythromycin, but can weaken the activity of some antibiotics including  
219 vancomycin (Schneierson & Amsterdam, 1958; Stanley Schneierson *et al.*, 1962). In a study  
220 which compared the effects of various bile acids in their salt form, as well as other components  
221 of bile such as cholesterol, SC and SDC were found to be the most effective at increasing the

222 anti-staphylococcal activity of neomycin (Stanley Schneierson *et al.*, 1962). We investigated  
223 the effect of SC on antibiotic tolerance in *S. aureus* RN4220. Disc diffusion antibiotic  
224 sensitivity testing revealed that 0.3% bile increased the sensitivity of *S. aureus* RN4220  
225 towards gentamicin, but had no effect on chloramphenicol, ampicillin, erythromycin or  
226 tetracycline (Fig. S1). We suspected that bile acids may specifically induce sensitivity to  
227 aminoglycosides in *S. aureus* RN4220. After confirming that SC at a concentration of 1 mM  
228 did not have bactericidal or bacteriostatic effects on this strain (Fig. S2), we proceeded to test  
229 the MIC of gentamycin, streptomycin, neomycin and kanamycin both in the presence of 1mM  
230 SC. The MIC of all aminoglycosides was reduced in the presence of 1 mM SC (Table 3),  
231 consistent with a previous study in *Lactobacillus* species (Elkins & Mullis, 2004), where the  
232 authors implicated increased antibiotic uptake following bile exposure.

233         We hypothesized that the molecular control of bile-induced aminoglycoside sensitivity  
234 and bile-induced biofilm formation in *S. aureus* might be linked at a regulatory level. Following  
235 this hypothesis, we predicted that mutant RN4220 strains that overcame bile-induced  
236 aminoglycoside sensitivity might also exhibit an altered bile-induced biofilm formation  
237 phenotype. In order to isolate genetically marked mutant strains that had overcome bile-  
238 induced aminoglycoside sensitivity, we utilized random transposon mutagenesis using the  
239 mariner-based transposon *bursa aurealis* to generate pools of mutant RN4220 and then  
240 selected for mutants that had overcome bile-induced aminoglycoside sensitivity in the presence  
241 of 1 mM sodium cholate. A range of aminoglycosides was utilized to avoid bias towards  
242 mutations which conferred resistance via an antibiotic-specific mechanism. Aminoglycoside  
243 concentrations were adjusted in selection plates to a level at which we could observe over ten  
244 colonies per plate.

245         In multiple rounds of mutagenesis with various aminoglycosides, several hundred *S.*  
246 *aureus* mutants were isolated. For 44 mutants we identified the site of the transposon using

247 random-primed PCR. Table S1 shows the list of transposon mutants, the antibiotic and  
248 concentration used to screen for the mutant, and the site of transposon insertion. Thirty four of  
249 the mutants carried the transposon insertion within a defined open reading frame (ORF), while  
250 the remaining 10 mutants had the transposon insertion occurring at between two coding  
251 sequences. Where possible the putative gene and/or gene product associated with the  
252 transposon disruption were identified (Table S1).

### 253 **Regulatory mutations in wall-teichoic acid synthesis stimulate a hypersensitive bile-** 254 **induced biofilm response**

255 We screened all 44 mapped mutants for differential biofilm formation in response to bile using  
256 the crystal violet assay (Fig. S3). Of these, five exhibited an increased sensitivity to bile, in that  
257 they exhibited increased attachment in the presence of 0.03% bile, compared to wild-type  
258 RN4220. As previously discussed, although the parental RN4420 strain displayed a significant  
259 increase biofilm formation in the presence of 0.3% bile, it failed to respond to bile at the level  
260 of 0.03%. In contrast, five mutants, namely TM4, TM19, TM26, TM28 and TM39, showed an  
261 enhanced bile-response, displaying substantially increased biofilm formation in the presence  
262 of 0.03% bile (Fig. 3), a phenotype akin to several of the clinical isolates investigated.

263         Of the five mutants that exhibited biofilm stimulation in the presence of 0.03% bile,  
264 three carried the transposon insertion within a defined ORF (TM19, TM26, TM28), while two  
265 had the transposon insertion occurring at between two coding sequences (TM4, TM39).  
266 Interestingly, in both mutants from the latter category, the transposon insertion was between  
267 divergently oriented genes where at least one gene was associated with wall teichoic acid  
268 (WTA) biosynthesis. TM4 carried an insertion between divergently oriented WTA  
269 biosynthesis genes, *tagG* and *tagH* (Lazarevic & Karamata, 1995; Schirner *et al.*, 2011). In  
270 TM39 the insertion was between *tagO*, which is associated with WTA biosynthesis (Soldo *et*  
271 *al.*, 2002; Xia *et al.*, 2010), and *gdpS*, the only conserved GGDEF domain protein identified

272 thus far in *Staphylococcus* (Shang *et al.*, 2009). Furthermore, the transposon insertion in TM26  
273 was within *tarJ*, another gene involved in the WTA biosynthetic pathway (Brown *et al.*, 2013).  
274 It should be noted however, that although the insertion disrupted *tarJ*, this gene is duplicated  
275 in many *S. aureus* strains (Qian *et al.*, 2006), and the locus SAOUHSC\_00226 in the strain  
276 NCTC8325 most likely represents a second copy of *tarJ*. These data indicated that WTA  
277 synthesis or its regulation might be involved in the response to bile and control of biofilm  
278 production. The remaining two hypersensitive mutants carried insertions within *ctaB* (TM19),  
279 which encodes for heme O synthase, and a possible D-galactonate transporter (TM28).

280         The isolation of mutants in WTA-associated genes suggested that WTA synthesis  
281 and/or translocation were involved in the bile-induced biofilm response, but because two  
282 insertions were within intergenic regions, it was unclear if this was due to increased or  
283 decreased WTA. We PCR-amplified and cloned regions of *tagG-tagH* from RN4220 into the  
284 *S. aureus* shuttle vector pLI50, and introduced each clone into both TM4 and wild type  
285 RN4220. While the introduction of pLI50 into RN4220 did not alter its bile-induced biofilm  
286 response, introduction of pLI50 containing carrying *tagG* (along with the *sod* ribosomal  
287 binding site (Malone *et al.*, 2009); pLI50::*tagG*) into TM4 led to the restoration of the wild  
288 type phenotype (Fig. 4). Introduction of a similar construct carrying *tagH* had no effect (not  
289 shown). Interestingly, introduction of pLI50 containing a copy of the non-coding intergenic  
290 region located between *tagH* and *tagG* (pLI50::int\_*tagHG*) into in RN4220, produced a strain  
291 with an almost identical phenotype to TM4 (Fig. 4). Together these data suggest that the  
292 enhanced bile-induced biofilm phenotype in TM4 may be due to a reduced expression of *tagG*,  
293 caused by the transposon insertion disrupting an operator site for an activator of *tagG*. The  
294 introduction of pLI50::*tagG* into TM4 likely restores the expression of *tagG* independently of  
295 the activator, producing the observed wild type phenotype, while the introduction of the  
296 intergenic region in RN4220 may sequester a DNA-binding activator of *tagG* expression.

297 **DISCUSSION**

298 The pathogenesis of *S. aureus* in the lungs of respiratory patients is characterized by successful  
299 colonization and persistence, particularly in pediatric CF patients where it dominates the early  
300 developing lungs. The environmental factors that cause *S. aureus* to adopt this pervasive  
301 lifestyle are as yet unknown, but our data presents a strong case for a role for bile aspiration  
302 derived from GERD in the pathogenesis of this organism. In this study we demonstrate that  
303 physiologically relevant concentrations of bile, as well as sub inhibitory concentrations of bile  
304 acids, can alter the behavior of *S. aureus*, leading to enhanced biofilm formation by this  
305 important clinical pathogen. The biofilm mode of growth is frequently associated with chronic  
306 infections as it allows the bacterium to evade host defenses, and persist for extended periods  
307 of time (Furukawa *et al.*, 2006). Our observation is consistent with the increased incidence of  
308 *S. aureus* in pediatric patients suffering from GERD (Palm *et al.*, 2012; van der Doef *et al.*,  
309 2009), the likely source of bile acids in the lungs of these patients (Aseeri *et al.*, 2012; Pauwels  
310 *et al.*, 2012; Reen *et al.*, 2014).

311 This bile-mediated response presented an opportunity to probe more deeply the  
312 regulatory mechanisms governing the switch between the phenotypes associated with acute  
313 and chronic infection in *S. aureus*. Using a random transposon mutagenesis approach we  
314 identified five *S. aureus* mutants with an enhanced bile-induced biofilm response. Of these,  
315 three mutants had a transposon insertion within or directly upstream of genes involved in WTA  
316 biosynthesis. WTA are surface-exposed anionic glycopolymers present in many gram positive  
317 species of bacteria, covalently bound to the peptidoglycan layer (Brown *et al.*, 2013). Indeed  
318 WTA are the most abundant peptidoglycan bound glycopolymer in gram positive species,  
319 making up over half of the dry weight of the cell wall. WTA have been shown to play key roles  
320 in maintenance of cell shape, several aspects of cell division, modulation of antibiotic  
321 susceptibility, and host tissue colonization (Swoboda *et al.*, 2010; Weidenmaier *et al.*, 2004).

322 Moreover, WTA have been implicated in biofilm formation in staphylococcal species (Holland  
323 *et al.*, 2011; Vergara-Irigaray *et al.*, 2008). For example, WTA contain D-alanine modifications  
324 which allow *S. aureus* to modulate its surface charge, aiding its primary attachment to artificial  
325 surfaces before the formation of multiple cell layers (Gross *et al.*, 2001).

326 *S. aureus* WTA polymers are composed of 30-50 ribitol phosphate (Rbo-P) subunits,  
327 connected to peptidoglycan via the murein linkage unit GlcNAc-ManNAc-(glycerol phosphate  
328 [Gro-P])<sub>2-3</sub> (Yokoyama *et al.*, 1986). WTA biosynthesis is a complex multi-step process which  
329 is yet to be fully characterized (Brown *et al.*, 2013). However, it has been shown that *tagO*, the  
330 regulatory region of which is likely disrupted in TM39, is involved in initiating the synthesis  
331 of the aforementioned murein linkage unit (Soldo *et al.*, 2002; Xia *et al.*, 2010). TarJ (an  
332 alcohol dehydrogenase), the function of which is most likely abrogated in TM26, together with  
333 TarI and TarL, catalyzes the attachment of Rbo-P to the murein linkage unit. Once WTA  
334 polymer formation is complete, it is translocated across the plasma membrane by the two-  
335 component ABC (ATP-binding cassette) transporter TagGH (Lazarevic & Karamata, 1995;  
336 Schirner *et al.*, 2011), the regulatory region of which was mutated in TM4. TagG presumably  
337 facilitates translocation of WTA polymer across the plasma membrane following a  
338 conformational change in the transmembrane domain induced by TagH.

339 The occurrence of three independent mutations within or flanking genes involved in the  
340 WTA biosynthesis pathway leading to enhanced biofilm formation in the presence of bile,  
341 strongly suggests an involvement of WTA in the *S. aureus* bile-induced biofilm formation  
342 response. WTA are essential for biofilm formation and host-colonization by *S. aureus* (Gross  
343 *et al.*, 2001; Weidenmaier *et al.*, 2004), and deletion of *tagO* impairs biofilm production in  
344 *Staphylococcus epidermidis* (Holland *et al.*, 2011). Interestingly, none of our isolated mutants  
345 carried complete knockouts in genes essential for WTA biosynthesis. Mutants TM4 and TM39  
346 carried intergenic insertions, while in TM26 the insertion was in one of two *tarJ* genes in *S.*



347 *aureus* (Qian *et al.*, 2006). This pattern of insertions is consistent with bias against WTA-  
348 associated insertions in the Nebraska Transposon Mutant Library, which consists of 1,952  
349 strains, each containing a single mutation within a nonessential gene of *S. aureus* isolate  
350 USA300. This collection also lacks mutations within *tagO*, *tagG*, and *tagH*, suggesting these  
351 may be essential genes (Fey *et al.*, 2013). Thus the mutations within the hyper-biofilm mutants  
352 likely do not knock out WTA synthesis, but may merely reduce it and concomitantly induce  
353 sensitivity to bile, leading to an increased level of biofilm production on exposure to lower  
354 concentrations of bile.

355         As bile is known to induce cell membrane damage (Begley *et al.*, 2005), and biofilms  
356 protect microorganisms from external damaging-agents (Flemming & Wingender, 2010), our  
357 observations lead us to propose that the bile-induced biofilm phenotype is an adaptive response  
358 to increased cell wall stress. In line with this, the inhibition of WTA synthesis by deletion of  
359 LytR-CpsA-Psr proteins has been shown to increase the basal expression of the ‘cell wall stress  
360 stimulon’, a collection of genes responsible for mounting a general cell wall stress response in  
361 the presence of cell wall damaging agents (Dengler *et al.*, 2012). The recently discovered  
362 antibiotic targocil, which blocks expression of *tagG*, has also been shown to induce cell wall  
363 stress in *S. aureus* (Campbell *et al.*, 2012). It is plausible that in the hyper-biofilm mutants  
364 identified in the present study, the transposon insertions reduce WTA biosynthesis, thereby  
365 stimulating a more profound cell wall stress-response in the presence of bile.

366         The transposon screen employed by this study utilized an indirect screening approach  
367 to identify genes within regulatory pathways involved in the *S. aureus* bile-response. This  
368 approach was limited in that all mutations identified as being important in biofilm formation  
369 were a subset of the mutations which conferred aminoglycoside resistance. As such, mutations  
370 which alter the bile-modulated biofilm response, but do not affect bile-induced aminoglycoside  
371 sensitivity, were not able to be detected in this study. As the bile induced aminoglycoside

372 sensitivity is likely caused by cell membrane perturbation by bile, leading to increased uptake  
373 of aminoglycosides, it was not surprising to observe a large number of mutations within genes  
374 involved in cell envelope integrity. Despite these limitations, the data presented in this study  
375 provides several insights into the possible regulatory mechanisms governing the bile-response  
376 in *S. aureus*. Further investigation of these mutants will provide insights into bile-responsive  
377 pathways linking biofilm formation to host-triggers of chronic infection, thereby facilitating  
378 the development of innovative strategies for the prevention and treatment of respiratory  
379 disease.

## 380 **ACKNOWLEDGMENTS**

381 The authors thank Prof Dominique Missiakas for providing pBursa and pFA545.

382

383 This work was supported in part by research grants awarded to FOG by the Health Science  
384 Faculty Curtin University, the European Commission (FP7-PEOPLE-2013-ITN, 607786; FP7-  
385 KBBE-2012-6, CP-TP-312184; FP7-KBBE-2012-6, 311975; OCEAN 2011-2, 287589; Marie  
386 Curie 256596; EU-634486), Science Foundation Ireland (SSPC-2, 12/RC/2275;  
387 13/TIDA/B2625; 12/TIDA/B2411; 12/TIDA/B2405; 14/TIDA/2438), the Department of  
388 Agriculture and Food (FIRM/RSF/CoFoRD; FIRM 08/RDC/629; FIRM 1/F009/MabS; FIRM  
389 13/F/516), the Irish Research Council for Science, Engineering and Technology  
390 (PD/2011/2414; GOIPG/2014/647), the Health Research Board/Irish Thoracic Society  
391 (MRCG-2014-6), the Marine Institute (Beaufort award C2CRA 2007/082) and Teagasc (Walsh  
392 Fellowship 2013).

393

394

395 **ABBREVIATIONS**

396 CF, cystic fibrosis; GERD, gastro-esophageal reflux; disease; LB, lysogeny broth; MIC,  
397 minimum inhibitory concentration; ORF, open reading frame; PIA, polysaccharide  
398 intercellular adhesion; SC, sodium cholate; SDC, sodium deoxycholate; TSA, tryptic soy agar;  
399 TSB, tryptic soy broth.

400 **REFERENCES**

401 **Aseeri, A., Brodlie, M., Lordan, J., Corris, P., Pearson, J., Ward, C. & Manning, N.**  
402 **(2012).** Bile acids are present in the lower airways of people with cystic fibrosis. *Am J Respir*  
403 *Crit Care Med* **185**, 463.

404  
405 **Baba, T., Takeuchi, F., Kuroda, M., Yuzawa, H., Aoki, K., Oguchi, A., Nagai, Y., Iwama,**  
406 **N., Asano, K. & other authors (2002).** Genome and virulence determinants of high virulence  
407 community-acquired MRSA. *Lancet* **359**, 1819-1827.

408  
409 **Bae, T., Banger, A. K., Wallace, A., Glass, E. M., Aslund, F., Schneewind, O. & Missiakas,**  
410 **D. M. (2004).** *Staphylococcus aureus* virulence genes identified by *bursa aurealis* mutagenesis  
411 and nematode killing. *Proc Natl Acad Sci U S A* **101**, 12312-12317.

412  
413 **Begley, M., Gahan, C. G. & Hill, C. (2005).** The interaction between bacteria and bile. *FEMS*  
414 *Microbiol Rev* **29**, 625-651.

415  
416 **Blainey, P. C., Milla, C. E., Cornfield, D. N. & Quake, S. R. (2012).** Quantitative analysis  
417 of the human airway microbial ecology reveals a pervasive signature for cystic fibrosis. *Sci*  
418 *Transl Med* **4**, 153ra130.

419  
420 **Brown, S., Santa Maria Jr, J. P. & Walker, S. (2013).** Wall teichoic acids of gram-positive  
421 bacteria. *Annu Rev Microbiol* **67**.

422  
423 **Button, B. M., Roberts, S., Kotsimbos, T. C., Levvey, B. J., Williams, T. J., Bailey, M.,**  
424 **Snell, G. I. & Wilson, J. W. (2005).** Gastroesophageal reflux (symptomatic and silent): a  
425 potentially significant problem in patients with cystic fibrosis before and after lung  
426 transplantation. *J Heart Lung Transplant* **24**, 1522-1529.

427  
428 **Campbell, J., Singh, A. K., Swoboda, J. G., Gilmore, M. S., Wilkinson, B. J. & Walker,**  
429 **S. (2012).** An antibiotic that inhibits a late step in wall teichoic acid biosynthesis induces the  
430 cell wall stress stimulon in *Staphylococcus aureus*. *Antimicrob Agents Chemother* **56**, 1810-  
431 1820.

432  
433 **Chua, K., Seemann, T., Harrison, P. F., Davies, J. K., Coutts, S. J., Chen, H., Haring, V.,**  
434 **Moore, R., Howden, B. P. & other authors (2010).** Complete genome sequence of  
435 *Staphylococcus aureus* strain JKD6159, a unique Australian clone of ST93-IV community  
436 methicillin-resistant *Staphylococcus aureus*. *J Bacteriol* **192**, 5556-5557.

437

438 **Cooper, M. A. & Shlaes, D. (2011).** Fix the antibiotics pipeline. *Nature* **472**, 32.  
439  
440 **D'Ovidio, F., Mura, M., Tsang, M., Waddell, T. K., Hutcheon, M. A., Singer, L. G.,**  
441 **Hadjiliadis, D., Chaparro, C., Gutierrez, C. & other authors (2005a).** Bile acid aspiration  
442 and the development of bronchiolitis obliterans after lung transplantation. *J Thorac Cardiovasc*  
443 *Surg* **129**, 1144-1152.  
444  
445 **D'Ovidio, F., Singer, L. G., Hadjiliadis, D., Pierre, A., Waddell, T. K., de Perrot, M.,**  
446 **Hutcheon, M., Miller, L., Darling, G. & other authors (2005b).** Prevalence of  
447 gastroesophageal reflux in end-stage lung disease candidates for lung transplant. *Ann Thorac*  
448 *Surg* **80**, 1254-1260.  
449  
450 **Dengler, V., Meier, P. S., Heusser, R., Kupferschmid, P., Fazekas, J., Friebe, S., Stauffer,**  
451 **S. B., Majcherczyk, P. A., Moreillon, P. & other authors (2012).** Deletion of hypothetical  
452 wall teichoic acid ligases in *Staphylococcus aureus* activates the cell wall stress response.  
453 *FEMS Microbiol Lett* **333**, 109-120.  
454  
455 **el-Serag, H. B. & Sonnenberg, A. (1997).** Associations between different forms of gastro-  
456 oesophageal reflux disease. *Gut* **41**, 594-599.  
457  
458 **Elkins, C. A. & Mullis, L. B. (2004).** Bile-mediated aminoglycoside sensitivity in  
459 *Lactobacillus* species likely results from increased membrane permeability attributable to  
460 cholic acid. *Appl Environ Microbiol* **70**, 7200-7209.  
461  
462 **Fey, P. D., Endres, J. L., Yajjala, V. K., Widhelm, T. J., Boissy, R. J., Bose, J. L. & Bayles,**  
463 **K. W. (2013).** A genetic resource for rapid and comprehensive phenotype screening of  
464 nonessential *Staphylococcus aureus* genes. *MBio* **4**, e00537-00512.  
465  
466 **Flemming, H. C. & Wingender, J. (2010).** The biofilm matrix. *Nat Rev Microbiol* **8**, 623-  
467 633.  
468  
469 **Furukawa, S., Kuchma, S. L. & O'Toole, G. A. (2006).** Keeping their options open: acute  
470 versus persistent infections. *J Bacteriol* **188**, 1211-1217.  
471  
472 **Gross, M., Cramton, S. E., Gotz, F. & Peschel, A. (2001).** Key role of teichoic acid net  
473 charge in *Staphylococcus aureus* colonization of artificial surfaces. *Infect Immun* **69**, 3423-  
474 3426.  
475  
476 **Holland, L. M., Conlon, B. & O'Gara, J. P. (2011).** Mutation of *tagO* reveals an essential  
477 role for wall teichoic acids in *Staphylococcus epidermidis* biofilm development. *Microbiology*  
478 **157**, 408-418.  
479  
480 **Kahl, B. C. (2010).** Impact of *Staphylococcus aureus* on the pathogenesis of chronic cystic  
481 fibrosis lung disease. *Int J Med Microbiol* **300**, 514-519.  
482  
483 **Kreiswirth, B. N., Lofdahl, S., Betley, M. J., O'Reilly, M., Schlievert, P. M., Bergdoll, M.**  
484 **S. & Novick, R. P. (1983).** The toxic shock syndrome exotoxin structural gene is not detectably  
485 transmitted by a prophage. *Nature* **305**, 709-712.  
486

487 **Kristiansen, T. Z., Maitra, A. & Pandey, A. (2007).** Proteomics of human bile. In Proteomics of Human Body Fluids, pp. 399-414. Edited by V. Thongboonkerd. Totowa, New Jersey: Humana Press.

488  
489  
490

491 **Lazarevic, V. & Karamata, D. (1995).** The *tagGH* operon of *Bacillus subtilis* 168 encodes a two-component ABC transporter involved in the metabolism of two wall teichoic acids. *Mol Microbiol* **16**, 345-355.

492  
493  
494

495 **Lee, C. Y., Buranen, S. L. & Ye, Z. H. (1991).** Construction of single-copy integration vectors for *Staphylococcus aureus*. *Gene* **103**, 101-105.

496  
497

498 **Legendre, C., Reen, F. J., Woods, D. F., Mooij, M. J., Adams, C. & O'Gara, F. (2014).** Bile acids repress hypoxia-inducible factor 1 signaling and modulate the airway immune response. *Infect Immun* **82**, 3531-3541.

499  
500  
501

502 **Malone, C. L., Boles, B. R., Lauderdale, K. J., Thoendel, M., Kavanaugh, J. S. & Horswill, A. R. (2009).** Fluorescent reporters for *Staphylococcus aureus*. *J Microbiol Methods* **77**, 251-260.

503  
504  
505

506 **Nassr, A. O., Gilani, S. N., Atie, M., Abdelhafiz, T., Connolly, V., Hickey, N. & Walsh, T. N. (2011).** Does impaired gallbladder function contribute to the development of Barrett's esophagus and esophageal adenocarcinoma? *J Gastrointest Surg* **15**, 908-914.

507  
508  
509

510 **Navarro, J., Rainisio, M., Harms, H. K., Hodson, M. E., Koch, C., Mastella, G., Strandvik, B. & McKenzie, S. G. (2001).** Factors associated with poor pulmonary function: cross-sectional analysis of data from the ERF. *Eur Respir J* **18**, 298-305.

511  
512  
513

514 **Palm, K., Sawicki, G. & Rosen, R. (2012).** The impact of reflux burden on *Pseudomonas* positivity in children with cystic fibrosis. *Pediatr Pulmonol* **47**, 582-587.

515  
516

517 **Pauwels, A., Decraene, A., Blondeau, K., Mertens, V., Farre, R., Proesmans, M., Van Bleyenbergh, P., Sifrim, D. & Dupont, L. J. (2012).** Bile acids in sputum and increased airway inflammation in patients with cystic fibrosis. *Chest* **141**, 1568-1574.

518  
519  
520

521 **Perng, D. W., Chang, K. T., Su, K. C., Wu, Y. C., Wu, M. T., Hsu, W. H., Tsai, C. M. & Lee, Y. C. (2007).** Exposure of airway epithelium to bile acids associated with gastroesophageal reflux symptoms: a relation to transforming growth factor-beta1 production and fibroblast proliferation. *Chest* **132**, 1548-1556.

522  
523  
524  
525

526 **Qian, Z., Yin, Y., Zhang, Y., Lu, L., Li, Y. & Jiang, Y. (2006).** Genomic characterization of ribitol teichoic acid synthesis in *Staphylococcus aureus*: genes, genomic organization and gene duplication. *BMC Genomics* **7**, 74.

527  
528  
529

530 **Reen, F. J., Woods, D. F., Mooij, M. J., Adams, C. & O'gara, F. (2012).** Respiratory pathogens adopt a chronic lifestyle in response to bile. *PLoS One* **7**, e45978.

531  
532

533 **Reen, F. J., Woods, D. F., Mooij, M. J., Chróinín, M. N., Mullane, D., Zhou, L., Quille, J., Fitzpatrick, D., Glennon, J. D. & other authors (2014).** Aspirated bile: a major host trigger modulating respiratory pathogen colonisation in cystic fibrosis patients. *Eur J Clin Microbiol Infect Dis* **33**, 1763-1771.

534  
535  
536

537  
538 **Schenk, S. & Laddaga, R. A. (1992).** Improved method for electroporation of *Staphylococcus*  
539 *aureus*. *FEMS Microbiol Lett* **73**, 133-138.  
540  
541 **Schirner, K., Stone, L. K. & Walker, S. (2011).** ABC transporters required for export of wall  
542 teichoic acids do not discriminate between different main chain polymers. *ACS Chem Biol* **6**,  
543 407-412.  
544  
545 **Schneierson, S. S. & Amsterdam, D. (1958).** Effect of bile salts upon the activity of  
546 antibiotics. *Nature* **182**, 56-57.  
547  
548 **Shang, F., Xue, T., Sun, H., Xing, L., Zhang, S., Yang, Z., Zhang, L. & Sun, B. (2009).** The  
549 *Staphylococcus aureus* GGDEF domain-containing protein, GdpS, influences protein A gene  
550 expression in a cyclic diguanylic acid-independent manner. *Infect Immun* **77**, 2849-2856.  
551  
552 **Soldo, B., Lazarevic, V. & Karamata, D. (2002).** *tagO* is involved in the synthesis of all  
553 anionic cell-wall polymers in *Bacillus subtilis* 168. *Microbiology* **148**, 2079-2087.  
554  
555 **Souza, H. A., Nogueira, K. S., Matos, A. P., Vieira, R. P., Riedi, C. A., Rosario, N. A.,**  
556 **Telles, F. Q. & Costa, L. M. (2006).** Early microbial colonization of cystic fibrosis patients  
557 identified by neonatal screening, with emphasis on *Staphylococcus aureus*. *J Pediatr (Rio J)*  
558 **82**, 377-382.  
559  
560 **Stanley Schneierson, S., Amsterdam, D. & Perlman, E. L. Y. (1962).** Effect of various bile  
561 salts on the anticandidal activity of amphotericin B and gentian violet and on the anti-  
562 staphylococcal activity of neomycin. *Nature* **196**, 909-910.  
563  
564 **Stringer, D. A., Sprigg, A., Juodis, E., Corey, M., Daneman, A., Levison, H. J. & Durie,**  
565 **P. R. (1988).** The association of cystic fibrosis, gastroesophageal reflux, and reduced  
566 pulmonary function. *Can Assoc Radiol J* **39**, 100-102.  
567  
568 **Sweet, M. P., Hoopes, C., Golden, J., Hays, S., Leard, L. & Patti, M. (2006).** Prevalence of  
569 delayed gastric emptying and gastroesophageal reflux in patients with end-stage lung disease.  
570 *Ann Thorac Surg* **82**, 1570.  
571  
572 **Sweet, M. P., Patti, M., Hoopes, C., Hays, S. & Golden, J. (2007a).** Gastro-oesophageal  
573 reflux and aspiration in patients with advanced lung disease. *Thorax* **64**, 167-173.  
574  
575 **Sweet, M. P., Patti, M. G., Leard, L. E., Golden, J. A., Hays, S. R., Hoopes, C. & Theodore,**  
576 **P. R. (2007b).** Gastroesophageal reflux in patients with idiopathic pulmonary fibrosis referred  
577 for lung transplantation. *J Thorac Cardiovasc Surg* **133**, 1078-1084.  
578  
579 **Swoboda, J. G., Campbell, J., Meredith, T. C. & Walker, S. (2010).** Wall teichoic acid  
580 function, biosynthesis, and inhibition. *Chembiochem* **11**, 35-45.  
581  
582 **van der Doef, H. P., Arets, H. G., Froeling, S. P., Westers, P. & Houwen, R. H. (2009).**  
583 Gastric acid inhibition for fat malabsorption or gastroesophageal reflux disease in cystic  
584 fibrosis: longitudinal effect on bacterial colonization and pulmonary function. *J Pediatr* **155**,  
585 629-633.  
586

587 **Vergara-Irigaray, M., Maira-Litran, T., Merino, N., Pier, G. B., Penades, J. R. & Lasa,**  
588 **I. (2008).** Wall teichoic acids are dispensable for anchoring the PNAG exopolysaccharide to  
589 the *Staphylococcus aureus* cell surface. *Microbiology* **154**, 865-877.  
590

591 **Weidenmaier, C., Kokai-Kun, J. F., Kristian, S. A., Chanturiya, T., Kalbacher, H., Gross,**  
592 **M., Nicholson, G., Neumeister, B., Mond, J. J. & other authors (2004).** Role of teichoic  
593 acids in *Staphylococcus aureus* nasal colonization, a major risk factor in nosocomial infections.  
594 *Nat Med* **10**, 243-245.  
595

596 **Wertheim, H. F., Melles, D. C., Vos, M. C., van Leeuwen, W., van Belkum, A., Verbrugh,**  
597 **H. A. & Nouwen, J. L. (2005).** The role of nasal carriage in *Staphylococcus aureus* infections.  
598 *Lancet Infect Dis* **5**, 751-762.  
599

600 **Wu, J. C. (2008).** Gastroesophageal reflux disease: an Asian perspective. *J Gastroenterol*  
601 *Hepatol* **23**, 1785-1793.  
602

603 **Wu, Y. C., Hsu, P. K., Su, K. C., Liu, L. Y., Tsai, C. C., Tsai, S. H., Hsu, W. H., Lee, Y.**  
604 **C. & Perng, D. W. (2009).** Bile acid aspiration in suspected ventilator-associated pneumonia.  
605 *Chest* **136**, 118-124.  
606

607 **Xia, G., Kohler, T. & Peschel, A. (2010).** The wall teichoic acid and lipoteichoic acid  
608 polymers of *Staphylococcus aureus*. *Int J Med Microbiol* **300**, 148-154.  
609

610 **Yokoyama, K., Miyashita, T., Araki, Y. & Ito, E. (1986).** Structure and functions of linkage  
611 unit intermediates in the biosynthesis of ribitol teichoic acids in *Staphylococcus aureus* H and  
612 *Bacillus subtilis* W23. *Eur J Biochem* **161**, 479-489.  
613

614 **Zecca, E., Costa, S., Lauriola, V., Vento, G., Papacci, P. & Romagnoli, C. (2004).** Bile acid  
615 pneumonia: a "new" form of neonatal respiratory distress syndrome? *Pediatrics* **114**, 269-272.  
616

617 **Zecca, E., De Luca, D., Baroni, S., Vento, G., Tiberi, E. & Romagnoli, C. (2008).** Bile acid-  
618 induced lung injury in newborn infants: a bronchoalveolar lavage fluid study. *Pediatrics* **121**,  
619 e146-149.  
620

621

622 **FIGURES**

623 **Fig. 1 Bile-dependent biofilm formation in *Staphylococcus aureus*.** (a) Pediatric clinical isolates  
624 compared to NCDO949, and (b) Community acquired methicillin-resistant isolates compared to  
625 RN4220. Data is presented as a ratio of the Abs595 nm from treated vs untreated samples. Graphs show  
626 mean (+/- SEM) relative absorbance of at least three independent experiments, each carried out in  
627 triplicate. Statistical analysis was performed by two-tailed paired Student's t-test (\* P < 0.05, \*\* P <  
628 0.01, compared with untreated control). (c) Growth of *S. aureus* RN4220 in presence of bovine bile.  
629 Graphs show mean (+/- SD) OD600 nm of two biological replicates.

630 **Fig. 2 Biofilm formation in *Staphylococcus aureus* RN4220 in response to bile and bile salts.** *S.*  
631 *aureus* strain RN4220 cultured for 18 h in TSB (untreated), or TSB supplemented with (a) sodium  
632 cholate (SC), (b) sodium deoxycholate (SDC), and (c) sodium dodecyl sulphate (SDS). Graphs show  
633 mean (+/- SD) relative absorbance of three biological replicates. Statistical analysis was performed a  
634 two-tailed unpaired Student's t-test (\* P < 0.05, \*\* P < 0.01 compared with untreated).

635 **Fig. 3 Altered bile-dependent biofilm formation in *Staphylococcus aureus* transposon mutants.**  
636 Graphs show mean (+/- SD) absorbance from three independent experiments each performed in  
637 triplicate. Statistical analysis was performed a two-tailed paired Student's t-test (\*\* P < 0.01).

638 **Fig. 4 Complementation of bile induced biofilm phenotype.** Transposon mutant TM4 and wild type  
639 RN4220 strains were transformed with plasmid pLI50 containing parts of the *tagG*-*tagH* genomic  
640 region from RN4220 (pLI50::int\_tagHG, pLI50 containing intergenic region between *tagH* and *tagG*;  
641 pLI50::tagG, pLI50 containing *tagG*). Strains were cultured for 18h in TSB (untreated), or TSB  
642 supplemented 0.03% bile, following which biofilm attachment was measured. Graphs show mean (+/-  
643 SD) absorbance from 3 replicates.

644



## 645 TABLES

646 Table 1 Bacterial strains and plasmids

Strain/ Plasmid	Description	Source/ Reference
<i>S. aureus</i>		
NCDO949	Type strain, pleural fluid isolate	Shinfield, UK
CUHT	Pediatric Clinical Isolate, CUH clinic	BRC
CUHE	Pediatric Clinical Isolate, CUH clinic	BRC
JKD6159	Dominant CA-MRSA strain in Australia, ST93-IV	(Chua <i>et al.</i> , 2010)
MW2	First community acquired CA-MRSA, ST1-MRSA-IVa (2B) (pMW2)	(Baba <i>et al.</i> , 2002)
RN4220	Highly transformable restriction-minus derivative of NCTC8325-4	(Kreiwirth <i>et al.</i> , 1983)
<i>E. coli</i>		
EPI300	Restriction-minus [ <i>mcrA</i> , $\Delta$ ( <i>mcrCB-hsdSMR-mrr</i> )], recombination-minus ( <i>recA1</i> ), endonuclease-minus ( <i>endA1</i> )	Epicentre
Plasmids		
pBursa	Encodes <i>bursa aurealis</i> transposable element, a temperature sensitive origin of replication, and chloramphenicol resistance selection marker	(Bae <i>et al.</i> , 2004)
pFA545	Encodes a transposase, origin of replication, and a tetracycline resistance selection marker	(Bae <i>et al.</i> , 2004)
pLI50	<i>S. aureus</i> (chloramphenicol resistance selection marker) / <i>E. coli</i> (ampicillin resistance selection marker) shuttle vector	(Lee <i>et al.</i> , 1991)
pLI50::int_tagHG	pLI50 containing the <i>tagG-tagH</i> intergenic amplified from RN4220 genomic DNA using primers TagHGIntFor and TagHGIntRev and cloned as EcoRI-BamHI fragment	This study
pLI50::tagG	pLI50 containing <i>tagG</i> amplified from RN4220 genomic DNA using primers TagGForRBS and TagHGRev and cloned as EcoRI-BamHI fragment	This study

647 CA-MRSA, Community-acquired Methicillin-resistant *Staphylococcus aureus*; ST, sequence type.

648

649 **Table 2** **Oligonucleotide sequences**

Primer name	Sequence (5' - 3')
GFP1	TCCACTGACAGAAAATTTGTGCCCATTAAC
GFP2	CATTAACATCACCATCTAATTCAACAAGAA
GFP3	ACAAGAATTGGGACAACCTCCAGTGA
PF106	GACCACACGTCGACTAGTGCNNNNNNNNNNNAGAG
PF107	GACCACACGTCGACTAGTGCNNNNNNNNNNNACGCC
PF108	GACCACACGTCGACTAGTGCNNNNNNNNNNNGATAC
PF109	GACCACACGTCGACTAGTGC
TagHGIntFor	TAGTAAAGCTTGAATTCTTGTAGACCTTCCTTATTCACATT
TagHGIntRev	TAGTAGGATCCTCCATTAACCACACTTTCAAATGT
TagGForRBS	TAGTAGAATTCTTAGGAGGATGATTATTTATGTCAGCAATAGGAACAG TTTTT
TagHGRev	TAGTAGGATCCTTACAAGAAGTCTGCAAATTGATCTC

650

651 **Table 3** **Effect of SC on the minimum inhibitory concentration of aminoglycosides in *S.***  
 652 ***aureus*.**

Antibiotic	MIC	
	0 mM SC	1 mM SC
Gentamycin	4 µg/mL	0.5 µg/mL
Streptomycin	16 µg/mL	8 µg/mL
Neomycin	4 µg/mL	1 µg/mL
Kanamycin	8 µg/mL	2 µg/mL

653 The *S. aureus* RN4220 minimum inhibitory concentration of various aminoglycosides in the presence  
 654 and absence of 1 mM SC is shown.

655

656 **SUPPLEMENTARY FIGURES**

657 **Fig. S1**            **Effect of 0.3% bile on antibiotic sensitivity in *Staphylococcus aureus* measured by**  
658 **disc diffusion testing.**

659 **Fig. S2**            **Effect of sodium cholate on the growth of *Staphylococcus aureus* RN4220.** Growth  
660 of *S. aureus* RN4220 in the presence of increasing concentrations of sodium cholate revealed that a  
661 concentration of 1 mM did not have any bacteriostatic or bactericidal effects on this strain. To determine  
662 the bacteriostatic effects of sodium cholate, a single colony of *S. aureus* was resuspended in 1 mL of  
663 TSB or TSB supplemented sodium cholate, and 200  $\mu$ L aliquots transferred to a 96-well plate. The  
664 density of the *S. aureus* cells were quantified by measuring the absorbance at 600 nm following  
665 incubation at 37°C for 24 hours with agitation at 250 rpm. Two  $\mu$ L of the culture was subsequently  
666 dotted on to TSA plates and incubated at 37°C for 24 hours to determine the bactericidal effects of  
667 sodium cholate. Graph shows mean (+/- SD) absorbance of three replicates. Photographs show the  
668 bacterial growth resulting from the 2  $\mu$ L culture on TSA.

669 **Fig. S3**            **Mutational analysis of the bile-induced biofilm response in *S. aureus*.** *S. aureus*  
670 transposon mutants were cultured in TSB (untreated) or TSB supplemented with 0.3% bile. For each  
671 mutant biofilm formation is presented as Abs<sub>595nm</sub> in both treated and untreated samples. Graph shows  
672 the mean (+/- SD) absorbance of four biological replicates.

673

## 674 SUPPLEMENTARY TABLES

675 Table S1 Genes disrupted in *Staphylococcus aureus* by transposon mutagenesis.

Mutant	Insertion Site <sup>a</sup>	ORF <sup>b</sup>	Putative Gene/Gene Product
Gentamicin (0.5 µg/mL) and SC (1 mM)			
TM1	1775393	SAOUHSC_01868	Dipeptidase PepV
TM2	1676550	SAOUHSC_01776	<i>hemA</i> ; Glutamyl-tRNA reductase
TM3	1444671	SAOUHSC_01488	Heptaprenyl pyrophosphate synthase subunit A
Gentamicin (0.35 µg/mL) and SC (1 mM)			
TM4	630715	SAOUHSC_00641- SAOUHSC_00642	<i>tagH</i> <i>tagG</i>
Gentamicin (0.3 µg/mL) and SC (1 mM)			
TM5	252438	SAOUHSC_00230	<i>lytS</i> ; two-component sensor His-kinase
TM6	205584	SAOUHSC_00186- SAOUHSC_00187	ABC transporter substrate-binding Pyruvate formate lyase
TM7	814365	SAOUHSC_00843	ABC-transporter interface protein
TM8	302717	SAOUHSC_00290	PTS system transporter
TM9	1602302	SAOUHSC_01691	Competence protein ComEC/Rec2
TM10	2605235	SAOUHSC_02826	MarR HTH regulator
TM11	2385772	SAOUHSC_02595	Sodium bile acid transporter
TM12	2083628	SAOUHSC_02247- SAOUHSC_02248	<i>ktrB</i> ; Potassium transporter GNAT family acetyltransferase
TM13	135192	SAOUHSC_00129	UDP-N-acetylglucosamine 2-epimerase
TM14	1691106	SAOUHSC_01793- SAOUHSC_01794	<i>nrdR</i> Glyceraldehyde 3-phosphate dehydrogenase
TM15	43138	SAOUHSC_00039	Dihydrouridine synthase
TM16	1233041	SAOUHSC_01275	<i>glpF</i> ; Glycerol uptake facilitator
TM17	433362	SAOUHSC_00435	Glutamate synthase subunit
TM18	2805147	SAOUHSC_03033	<i>nixA</i> ; Nickel transporter
TM19	1033768	SAOUHSC_01066	<i>ctaB</i> ; heme O synthase
TM20	2084622	SAOUHSC_02249	Membrane protein
TM21	974315	SAOUHSC_01001	<i>qoxB</i> ; Quinol oxidase subunit I

---

TM22	2807414	SAOUHSC_03035	Membrane protein
Streptomycin (4 µg/mL) and SC (1 mM)			
TM23	2536967	SAOUHSC_02760	Glutamate synthase
TM24	2007589	SAOUHSC_02133-	<i>nadC</i> ; Nicotinate phosphoribosyltransferase
		SAOUHSC_02134	Nitric oxide synthase oxygenase
TM25	2604892	SAOUHSC_02825	Glyoxalase
SDC (0.2 mM)			
TM26	241577	SAOUHSC_00221	<i>tarJ</i> ; Ribitol-5-phosphate dehydrogenase
TM27	2296822	SAOUHSC_02474	hypothetical protein
TM28	2554107	SAOUHSC_02777	Probable D-galactonate transporter
TM29	2004986	SAOUHSC_02131	hypothetical protein
		SAOUHSC_02132	NAD synthetase
TM30	860206	SAOUHSC_00895	Glutamate dehydrogenase
TM31	495319	SAOUHSC_R0007	Cell wall-associated hydrolase
Neomycin (0.5 µg/mL) and SC (1 mM)			
TM36	136894	SAOUHSC_00131-	Membrane spanning protein
		SAOUHSC_00132	Aldehyde dehydrogenase
Kanamycin (1.5 µg/mL) and SC (1 mM)			
TM39	744341	SAOUHSC_00760-	<i>gdpS</i>
		SAOUHSC_00762	<i>tagO</i>
TM40	44147	SAOUHSC_00040	Membrane protein
TM41	1325036	SAOUHSC_01380-	<i>nikB</i> ; Nickel ABC transporter permease
		SAOUHSC_01381	Membrane protein
TM42	2387891	SAOUHSC_02597	<i>ptsG</i> ; Phosphotransferase system
TM43	2075304	SAOUHSC_02239-	Phage integrase
		SAOUHSC_02241	Leukocidin/Hemolysin toxin family
TM44	2735026	SAOUHSC_02972	<i>isaB</i> ; Immunodominant surface antigen B
TM45	2598814	SAOUHSC_02820	ABC-2 transporter family protein
TM46	71816	SAOUHSC_00067	<i>lldP</i> ; L-lactate permease
TM47	876020	SAOUHSC_00906	Fumarylacetoacetate (FAA) hydrolase family
TM48	226300	SAOUHSC_00202-	Uroporphyrinogen-III C-methyltransferase
		SAOUHSC_00203	Membrane protein
TM49	1787295	SAOUHSC_01873	Hypothetical Protein


---


---

TM50	549173	SAOUHSC_00544	<i>sdrC</i> ; Serine-aspartate repeat protein
------	--------	---------------	---


---


676 Transposon mutants are grouped according to antibiotic and/or bile acid concentrations used to screen  
677 for the mutant. <sup>a</sup> Genomic position in the NCTC8325 genome is presented. <sup>b</sup> Corresponding ORF in the  
678 NCTC8325 genome is presented.











Gentamycin  
10  $\mu$ g






Erythromycin  
15  $\mu$ g

Chloramphenicol  
30  $\mu$ g

Tetracycline  
30  $\mu$ g

Ampicillin  
10  $\mu$ g

0% Bile



0.3% Bile

