

Title	Microbial succession and flavor production in the fermented dairy beverage kefir
Authors	Walsh, Aaron M.;Crispie, Fiona;Kilcawley, Kieran N.;O'Sullivan, Orla;O'Sullivan, Maurice G.;Claesson, Marcus J.;Cotter, Paul D.
Publication date	2016-10-04
Original Citation	Walsh, A. M., Crispie, F., Kilcawley, K., O'Sullivan, O., O'Sullivan, M. G., Claesson, M. J. and Cotter, P. D. (2016) 'Microbial succession and flavor production in the fermented dairy beverage kefir', mSystems 1(5), e00052-16 (16pp). doi:10.1128/mSystems.00052-16
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="http://msystems.asm.org/content/2/1/e00003-17 - 10.1128/mSystems.00052-16">http://msystems.asm.org/content/2/1/e00003-17 - 10.1128/mSystems.00052-16</a>
Rights	© 2016, the Authors. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International license. - <a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
Download date	2025-01-24 19:26:33
Item downloaded from	<a href="https://hdl.handle.net/10468/3850">https://hdl.handle.net/10468/3850</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

**Table S1. Absolute abundances of bacteria and fungi in kefir samples after 0, 8 and 24 hours of fermentation, as determined by quantitative PCR (qPCR) measurements.**

<b>Sample</b>	<b>Total fungi (ng of fungal DNA)</b>	<b>Total bacteria (copies of 16S rRNA gene)</b>
Milk 0 h	0.0016	1.78E+05
Fr1 08 h	0.1386	7.64E+07
Fr1 24 h	0.2179	2.49E+08
Ick 08 h	0.0542	2.62E+08
Ick 24 h	0.0972	1.63E+08
UK3 08 h	0.0896	7.86E+07
Uk3 24 h	0.3933	4.22E+08