

Title	Staphylococcus lugdunensis cultured from the amniotic fluid at caesarean section
Authors	Marchocki, Zbigniew;Collins, J. Kevin;Lehane, Eimear;O'Reilly, Paddy;O'Donoghue, Keelin
Publication date	2013
Original Citation	Marchocki Z, Collins K, Lehane E, Reilly PO, O'Donoghue K (2013) Staphylococcus lugdunensis Cultured from the Amniotic Fluid at Caesarean Section. PLoS ONE 8(2): e56373. doi:10.1371/journal.pone.0056373
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1371/journal.pone.0056373
Rights	© 2013 Marchocki et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited - http://creativecommons.org/licenses/by/4.0/
Download date	2024-12-03 23:59:43
Item downloaded from	https://hdl.handle.net/10468/2393



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Staphylococcus lugdunensis Cultured from the Amniotic Fluid at Caesarean Section

Zbigniew Marchocki¹, Kevin Collins², Eimear Lehane², Paddy O' Reilly², Keelin O'Donoghue^{1*}

1 Anu Research Centre, Department of Obstetrics and Gynaecology, University College Cork, Cork University Maternity Hospital, Cork, Ireland, **2** Department of Microbiology, University College Cork, Cork, Ireland

Abstract

Staphylococcus lugdunensis is a virulent coagulase-negative staphylococcus. It behaves like and can be mistaken in culture for *Staphylococcus aureus*. While originally thought to be a skin commensal rarely responsible for opportunistic infection, it was rapidly established as a significant human pathogen. It has been mainly associated with native and prosthetic valve endocarditis, osteomyelitis, and skin and soft tissue cellulitis, but has also been reported as a cause of fasciitis as well as peritonitis. *Staphylococcus lugdunensis* has been reported as a cause of endometritis but has not been previously isolated from amniotic fluid. Here, amniotic fluid samples were collected in the course of a larger study on amniotic fluid bacteriology, with prior ethical approval and informed patient consent. Amniotic fluid was obtained at Caesarean Section by direct needle aspiration from the intact amnion. Analysis with *Staphylococcal* API test kits led to identification of *Staphylococcus lugdunensis* in two cases. The clinical significance of the finding in these reported cases is undetermined. *Staphylococcus lugdunensis* has been shown to be a cause of serious and potentially fatal morbidities, but this is the first report of its culture from amniotic fluid. As caesarean delivery is accepted as the single most important factor associated with post-partum infectious complications in both mother and neonate, the identification of this pathogen is a new concern.

Citation: Marchocki Z, Collins K, Lehane E, Reilly PO, O'Donoghue K (2013) *Staphylococcus lugdunensis* Cultured from the Amniotic Fluid at Caesarean Section. PLoS ONE 8(2): e56373. doi:10.1371/journal.pone.0056373

Editor: Nancy E. Freitag, University of Illinois at Chicago College of Medicine, United States of America

Received: October 30, 2012; **Accepted:** January 8, 2013; **Published:** February 7, 2013

Copyright: © 2013 Marchocki et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors have no support or funding to report.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: k.odonoghue@ucc.ie

Introduction

Staphylococcus lugdunensis is a gram-positive, catalase-positive, coagulase-negative *staphylococcus*. The organism was initially isolated and described in 1988 by Freney *et al.* [1]. It was first accepted as a skin commensal or occasionally opportunistic pathogen, and found to colonise mostly the groin and perineal regions [2]. However evidence published over the last decade confirm its life threatening potential both in immunosuppressed and immune-competent individuals. *S. lugdunensis* has now been shown to be an aetiological factor in cardiovascular infections, including native and prosthetic valve endocarditis, bloodstream infection (sepsis or septic shock), toxic shock syndrome, acute oral infection, urinary tract infection, bone and joint infection (infective arthritis or osteomyelitis, as well as prosthetic joint infection), central nervous system infection (brain abscess and meningitis), peritonitis and ocular infection [3,4]. It is also reported as a cause of painful and prolonged skin and soft tissue infections [4,5].

S. lugdunensis has been previously described in isolated case reports of post-partum infection but has not been reported isolated from amniotic fluid [6,7]. Caesarean section (CS) is an important risk factor for postpartum maternal infection [8,9]. Here, we present two cases of pregnant women who underwent CS and were incidentally found to have amniotic fluid colonised with this potentially dangerous pathogen.

Methods

Amniotic fluid (AF) samples were collected in the course of a larger study on amniotic fluid bacteriology. The study was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals [ref ECM 4: 02/08/2011] and written informed consent was obtained from each participant prior to CS. In each case a different operator (both senior clinicians) in a different operating theatre performed the CS, and the deliveries were weeks apart. As per hospital policy at the time, routine intravenous antibiotic prophylaxis was administered after delivery of the infant at CS in both cases.

AF samples were obtained at the time of CS by direct syringe aspiration from the intact amnion and stored at 4°C until processed. Samples were plated on Columbia Blood Agar. Subsequent Gram Stain, Catalase Test, Baird-Parker Selection Agar Plates, Rapid Agglutination Test and finally, analysis in *Staphylococcal* API test kits (bioMérieux, UK) led to identification of the *S. lugdunensis* species.

Results

The first case was a 38-year-old multiparous white woman who underwent elective repeat caesarean section (CS) at 37 weeks and 6 days. She had an emergency CS in her first pregnancy for uterine rupture in the second stage of labour. This pregnancy course was uneventful and she had no medical co-morbidities. She was a non-smoker and was overweight (body mass index (BMI),

28). Elective surgery was carried out without any complications and a liveborn infant of normal birth weight was delivered. Mother and baby left hospital four days after CS, with no infectious post-operative complications reported.

The second case was a 30-year-old nulliparous white woman, who underwent induction of labour with prostaglandin E2 (PGE2) for post-maturity. Her pregnancy course was uneventful and her medical co-morbidities included well-controlled asthma. She smoked 20 cigarettes daily prior to pregnancy and had a BMI of 25. Membranes spontaneously ruptured 7 hours after administration of PGE2 after which she was commenced on a syntocinon infusion. She subsequently underwent an emergency CS for fetal distress- an abnormal fetal blood sample (pH 7.14) was obtained following non-reassuring features evident on cardiotocography. Emergency caesarean section was performed and amniotic fluid obtained through the bulging amnion. A liveborn infant of normal birth weight was delivered without complication. Again, mother and baby left hospital four days after CS, with no infectious post-operative complications reported.

In both cases clear amniotic fluid was obtained by direct needle aspiration from the intact amnion at CS. Bacterial colonisation was noted in both AF samples; 90 colony forming units per ml (CFU/ml) in the first sample, and 550 CFU/ml in the second under aerobic conditions, with few under anaerobic conditions. *S. lugdunensis* species was subsequently identified in both AF samples.

Discussion

For many years, it was assumed that the amniotic fluid cavity prior to onset of labour was a sterile environment. Harris and Brown in 1927 investigated the presence of bacteria in amniotic fluid of women undergoing CS, to find that all women who were in labour for <6 hours had negative cultures, whereas those who were in labour for >6 hours had positive cultures [10]. Many reports have since confirmed that the amniotic cavity is not entirely sterile. Multiple organisms have been found using traditional bacterial culture methods and more recently, using advanced molecular techniques. Organisms identified include species of *Ureaplasma*, *Mycoplasma*, *Fusobacterium*, *Streptococcus*, *Staphylococcus*, *Bacteroides* and *Prevotella*.

Caesarean section (CS) is now a common surgical procedure, performed to deliver around 25% of all pregnancies. As CS has become safer and electively scheduled, functional and cosmetic aspects, as well as risks of hospital-acquired infection have gained increased importance. A CS delivery is the single most important factor associated with post-partum infection and carries a 5–20 fold increased risk of infection compared to vaginal delivery [9]. Endometritis, surgical site or urinary tract infections occur in about 8% of women who have a CS [8]. Case reports of early neonatal sepsis also suggest that the infection process is likely to be triggered in utero [11]. Risk factors associated with infection after CS include maternal obesity, prolonged operating time, poor surgical technique and emergency delivery [8,9]. While the majority of surgical site infections are not life threatening they have important implications on the length of hospital stay, hospital

costs and re-admission rates [9]. Around 80% of infections occur after discharge from hospital. With the increase in CS rates worldwide, it is anticipated that post CS infections will become an increasing health and economic burden [8,9].

Here, we report two women in whom amniotic fluid was colonised with the gram-positive bacteria, *S. lugdunensis*, at the time of CS. This organism has been already shown to be a cause of serious and potentially fatal morbidities in other medical specialities, and has been reported as a cause of post-partum endometritis and pyomyoma [6,7]. It has never been demonstrated at CS before now. CS is accepted as a major risk factor in post-partum infectious complications and the identification of this pathogen is a new concern. Further, the identification of *S. lugdunensis* can lead to confusion or misdiagnosis, as it can be easily mistaken for *Staphylococcus aureus* due to its production of the clumping factor that gives a positive result in slide coagulase or rapid latex agglutination tests [3].

Our findings could occur secondary to contamination, as *S. lugdunensis* is a skin commensal colonising the groin area [2]. However, we found the bacterium in samples taken by different senior operators on different days. Operators were trained in an aseptic technique of sampling, and neither sample was macroscopically or microscopically contaminated by blood. We could not facilitate two separate samples of AF being taken and analysed independently, given the clinical necessity of CS. In each case, the CS was performed for a different indication, one being emergency and the other elective (although the previous serious complication in the elective CS is noted and could be relevant). In both cases, standard aseptic techniques were observed in hand decontamination and antiseptic skin preparation prior to skin incision at CS. No infectious post-operative or neonatal complications were observed in either case before discharge, but neither was followed up in the hospital. Finally, procedures were standardised in a large accredited and internationally-recognised research laboratory. The clinical significance of the findings of *S. lugdunensis* in these reported cases therefore remains undetermined.

Conclusions

This is the first case report of *S. lugdunensis* cultured from amniotic fluid during caesarean section. CS delivery is the single most important factor associated with post-partum infectious complications, and the identification of this pathogen is a new concern. Further studies are needed to examine the prevalence of *S. lugdunensis* colonisation in amniotic fluid and to establish if there is an association with post-operative maternal or neonatal infectious morbidity, particularly after emergency CS.

Author Contributions

Conceived and designed the experiments: ZM KC KO. Performed the experiments: ZM EL PO KC. Analyzed the data: ZM PO KC KO. Contributed reagents/materials/analysis tools: PO KC KO. Wrote the paper: ZM KC KO.

References

1. Freney J, Brun M, Bes H, Meugnier F, Grimont PAD, et al. (1988) *Staphylococcus lugdunensis* sp. nov. and *Staphylococcus schleiferi* sp. nov., two species from human clinical specimens. *Int. J. Syst. Bacteriol*; 38: 168–172
2. Van der Mee-Marquet N, Achard A, Mereghetti L, Danton A, Minier M, et al. (2003) *Staphylococcus lugdunensis* infections: high frequency of inguinal area carriage. *J Clin Microbiol*; 41: 1404–1409
3. Frank KL, del Pozo JL, Patel R (2008) From Clinical Microbiology to Infection Pathogenesis: How Daring To Be Different Works for *Staphylococcus lugdunensis*. *Clinical Microbiology Reviews*; 111–133
4. Babu E, Oropello J (2011) *Staphylococcus lugdunensis*: the coagulase-negative staphylococcus you don't want to ignore. *Expert Rev Anti Infect Ther*; 9(10): 901–7
5. Hung T, Zaghi S, Yousefzadeh J, Leibowitz M (2012) Necrotizing Fasciitis Associated with *Staphylococcus lugdunensis*. *Case Rep Infect Dis*; Epub 2012 May 17
6. Shaaban HS, Choo HF, Sensakovic JW (2011) A Case of *Staphylococcus lugdunensis* related pyomyoma occurring after cesarean section. *J Global Infect Dis*; 3: 101–2

7. Bello C, Eskandar M, El GR, Nour H, Shafiq H (2007) *Staphylococcus lugdunensis* endometritis: a case report. *West Afr J Med*; 26(3): 243–5
8. National Institute for Clinical Excellence (2011) NICE Clinical Guideline Number 132: Caesarean Section. National Institute for Clinical Excellence.
9. Lamont RF, Sobel J, Kusanovic JP, Vaisbuch E, Mazaki-Tovi S, et al. (2011) Current Debate on the Use of Antibiotic Prophylaxis for Caesarean Section. *BJOG*; 118(2): 193–201
10. Harris JW, Brown H (1927) Bacterial content of the uterus at Cesarean section. *Am J Obstet Gynecol*; 13:133
11. Bouziri A, Bel Hadj S, Khaldi A, Borgi A, Menif K, et al. (2011) Early neonatal infection caused by *Streptococcus pneumoniae*: report of 3 cases. *Tunis Med*; 89(2): 206–9.