

Title	Composition, morphology and pasting properties of <i>Orchis anatolica</i> tuber gum
Authors	Bulut-Solak, Birsen;Alonso-Miravalles, Loreto;O'Mahony, James A.
Publication date	2016-12-26
Original Citation	Bulut-Solak, B., Alonso-Miravalles, L. and O'Mahony, J. A. (2017) 'Composition, morphology and pasting properties of <i>Orchis anatolica</i> tuber gum', <i>Food Hydrocolloids</i> , 69, pp. 483-490. doi: 10.1016/j.foodhyd.2016.12.009
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.foodhyd.2016.12.009
Rights	© 2016 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. - http://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2025-05-24 23:13:08
Item downloaded from	https://hdl.handle.net/10468/3986



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Accepted Manuscript

Composition, morphology and pasting properties of *Orchis anatolica* tuber gum

Birsen Bulut-Solak, Loreto Alonso Miravalles, James A. O'mahony

PII: S0268-005X(16)30999-7

DOI: [10.1016/j.foodhyd.2016.12.009](https://doi.org/10.1016/j.foodhyd.2016.12.009)

Reference: FOOHYD 3716

To appear in: *Food Hydrocolloids*


Received Date: 29 July 2016


Revised Date: 9 December 2016

Accepted Date: 12 December 2016


Please cite this article as: Bulut-Solak, B., Alonso Miravalles, L., O'mahony, J.A., Composition, morphology and pasting properties of *Orchis anatolica* tuber gum, *Food Hydrocolloids* (2017), doi: 10.1016/j.foodhyd.2016.12.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.







Chemical compounds of *Orchis anatolica* tuber gum



Scanning electron micrographs of native *Orchis anatolica* tuber gum's powder particles



Light microscopy micrographs of *Orchis anatolica* tuber gum All: (a) 2.0% OaG; (b) 2.5% OaG. Left-hand column at 35 °C, middle-columns from 55 to 95 °C on heating, right-hand column at 47 °C on cooling. Scale bar 200 μ m.



Pasting curves for *Orchis anatolica* gum at concentrations of 0.5% (—), 1.5% (Δ), 2% (o) and 2.5% (*)

Composition, Morphology and Pasting Properties of *Orchis anatolica*

Tuber Gum

Birsen BULUT-SOLAK^{a,b*}, Loreto ALONSO MIRAVALLÉS^a and James A.

O'MAHONY^a

^aSchool of Food and Nutritional Sciences, University College Cork, Cork, Ireland

^bDepartment of Gastronomy and Culinary Arts, Faculty of Tourism, Selcuk University, 42000, Konya, Turkey

*Author for correspondence.

E-mail: birsenbulut@selcuk.edu.tr

Tel: +90 332 2234422

Fax: +90 332 2417047

1 Abstract

2

3 *Orchis anatolica* (*O. anatolica*) tuber is commonly used in the production of Salep gum or *O.*
4 *anatolica* tuber gum (*OaG*) for use as a thickener, flavouring agent, gelling agent, film former
5 and emulsifier in the food industry. The aim of this study was to investigate the chemical
6 composition, physical, morphological and pasting properties of *OaG*. Physical and morphological
7 analyses, and pasting properties of *OaG* were analysed using static light scattering, scanning
8 electron microscopy, light microscopy and rotational rheometry, respectively. Volume-weighted
9 mean particle diameter ($D [4,3]$) value of *OaG* was $180 \pm 1.25 \mu\text{m}$. *OaG* was composed mainly of
10 starch (41.6%), dietary fiber (32.3%) and glucomannan (18.5%). The powder of *OaG* had
11 irregular shaped particles with smooth surfaces and round edges. After pasting treatment, the
12 initial and final viscosity values of the *OaG* dispersions at a concentration of 0.5% *OaG* were
13 33.7 ± 0.24 and 34.3 ± 0.45 mPa.s, whereas, the corresponding values at a concentration of 2.5%
14 *OaG* were 1193 ± 92.0 and 1437 ± 83.3 mPa.s, respectively. The glucomannan and dietary fiber
15 components and their possible interactions with starch, in *OaG* appear to have influenced the
16 peak temperature and viscosity on pasting, due to limitation of the leaching of amylose and
17 amylopectin from starch granules. Therefore, *O. anatolica* tuber gum, a complex biopolymer, can
18 provide interesting and unique functionality to the food industry in the development of novel
19 food structures.

20

21 **Keywords:** Gelatinisation, Gum, *Orchis anatolica*, Orchidaceae, Pasting, Salep.

22

23 1.0. Introduction

24
25 *Orchis anatolica* (*O. anatolica*) is one of the terrestrial species of the *Orchis* genus in the
26 Orchidaceae family, which has a larger number of species than any other family of flowering
27 plants. *O. anatolica* generally grows in pine or macchie forests from the Southern Aegean Islands
28 to Iran, especially in parts of Anatolia. Salep gum or *OaG* is obtained from dried tubers of
29 terrestrial orchids. It is used either as a powder or in the form of a hot traditional beverage with
30 pleasant taste since the time of the Ottoman Empire (Tamer, Karaman, & Copur, 2006; Hossain,
31 2011; Georgiadis et al., 2012). Salep gum has also been used in medicines and ice-creams
32 (Lange, 1998).

33
34 Salep gum is a non-toxic, water-soluble and multi-component polysaccharide (e.g., glucomannan,
35 starch) with a high molecular weight (Pourjavadi, Doulabi, Soleyman, Sharif, & Eghtesadi, 2012)
36 compared with other polysaccharides, such as high-amylose starch (Salemis & Rinaudo, 1984),
37 chitosan and alginate (Harding, Abdelhameed, & Morris, 2010). It contains starch, mucilage,
38 sugar, nitrogenous substances, ash, particularly calcium, potassium, iron, chlorides and
39 phosphates, and some trace levels of volatile oils (Hossain, 2011; Lalika et al., 2013). It is a good
40 source of glucomannan that is considered a dietary fiber (Tekinsen & Guner, 2010; Nikjooy, Joo,
41 & Jahanshahi, 2014). The composition of orchid tuber also varies considerably, depending on
42 botanical origin, environmental and biological factors, such as the species, the development state
43 of the perennial plant, the stage of flowering, the time of harvesting, season and the site
44 composition (Tekinsen & Guner, 2010; Lalika et al., 2013).

45 Salep gum also confers many important functional properties such as thickening, stabilizing,
46 flavouring and gelling properties in food formulations, as well as medical formulations. It is one

47 of the most thermally-stable hydrocolloids and produces hydrogels for biomedical applications
48 due to its high capacity for water storage. In addition, Salep biopolymer modified by graft
49 copolymerization have higher swelling rates, which makes it a good candidate for drug delivery
50 systems (Bardajee, Hooshyar, & Kabiri, 2012; Pourjavadi et al., 2012). Therefore, Salep gum has
51 attracted increasing research interest in the last decade, because of its unique functional
52 properties and nutritional value in food and medical formulations (Alonso-Sande, Teijeiro-
53 Osorio, Remunan-Lopez, & Alonso, 2009; Bardajee, Hooshyar, Asli, Shahidi, & Dianatnejad,
54 2013).

55 Starch, which abounds in nature as an energy source in tubers, roots, cereal grains and legumes,
56 is widely used in the food industry. The starch content of plants is variable depending on
57 botanical origin, environmental and biological factors, such as the species, site composition,
58 harvesting etc. It consists primarily of two polysaccharides; amylose (AM) and amylopectin
59 (AP). These polysaccharides are homopolymers of α -D-glucose, and occur together in compact
60 assemblies declared as starch granules (Miles, Morris, & Ring, 1985; Kett et al., 2013). AM is
61 basically a linear polymer, with the formation of uncharged coaxial double helices being
62 promoted by its (1 \rightarrow 4)-axial-equatorial linkage geometry. On the other hand, the α -D-glucose
63 residues in AP are also extensively (1 \rightarrow 4)-linked, but with branches through 1 \rightarrow 6 linkages
64 (Considine et al., 2011).

65 When starch granules are heated in excess water, they hydrate and swell, because hydrogen
66 bonds in amorphous regions are disrupted, leading to absorption of water which acts as a
67 plasticizer. Greater hydration and swelling occur in amorphous regions, pulling apart crystallites,
68 with these regions ultimately undergoing hydration and melting (Considine et al., 2011). This
69 process is known as gelatinisation and is characterised by changes in viscosity. The gelatinisation

70 process includes “pasting temperature” which is the temperature coinciding with the rapid
71 increase in viscosity which ensues the onset of gelatinisation, and the “peak viscosity”, which is
72 the maximum viscosity value during pasting (Ratnayake & Jackson, 2009; Kett et al., 2013).
73 After the process of gelatinisation or pasting, the resultant viscoelastic mass called a “paste”
74 consists of a continuous phase with a molecular dispersion containing dissolved starch polymer
75 molecules forming a network, and discontinuous phase of swollen granules, granule ghosts and
76 granule fragments, depending on the origin of starches (Atkin, Abeysekera, & Robards, 1998;
77 Baldwin, 2001; Debet & Gidley, 2006; Kett et al., 2013).

78 Sezik (1967) and Tekinsen and Guner (2010) in previous works also researched the
79 compositional properties of *OaG* which would be expected to vary significantly due to biological
80 and environmental factors. Moreover, there is not much information in literature about the
81 chemical properties of *OaG*. Nevertheless, the physical, morphology and specifically pasting
82 properties of *OaG* have not been determined to date yet. There are also no recorded physical,
83 morphology and gelatinisation data available in literature about these properties of *OaG*.
84 Therefore, the objective in this study was to determine the chemical composition, physical,
85 morphological and pasting properties of *OaG* to better understand and predict its
86 physicochemical properties to aid its use as a functional ingredient in the food industry.

87

88 **2.0. Materials and Methods**

89

90 *2.1. Materials*

91 *O. anatolica*, which is a delicate perennial plant with pink to violet flowers, grows mostly in
92 lightly-shaded pine forests or mountains in several parts of Anatolia, such as in the Taurus
93 Mountains (Fig. 1 a, b). *O. anatolica* is similar to *O. quadripunctata* but spike lax of *O. anatolica*
94 has 5-10 large flowers with bracts. The colour of *O. anatolica*'s flowers is rose purple and/or
95 seldom white, with purple dots and lines in centre of labellum. Sepal of *O. anatolica*, with
96 spreading or \pm reflexed ovate-obtuse, is 7-11 x 3.5-4 mm. Labellum ovate of *O. anatolica* is also
97 nearly 10-15 mm with 3-lobed. Flowering time of *O. anatolica* is between March and May
98 (Altundag et al. 2012).

99 In this study, the fully formed orchid tubers, at the end of the flowering stage, were harvested
100 from several defined locations in the Taurus Mountains near Tatlicak, Konya (Turkey) between
101 June 1st and July 7th of 2013. This harvesting period was selected because the seed vessels of *O.*
102 *anatolica* would be fully formed at the end of flowering stage, and a randomized sampling
103 technique was used for tuber harvesting. The orchid plants at the flowering stage were identified
104 as being *O. anatolica* by the Herbarium Laboratory in the Biology Department at Selcuk
105 University, Konya, Turkey, as described by Altundag et al. (2012). All chemicals, reagents and
106 solvents used were of analytical grade and purchased from Sigma-Aldrich (Vale Road, Arklow,
107 Wicklow, Ireland).

108 109 2.2. Powder preparation

110 The round-edged appearances of *O. anatolica* freshly harvested from Taurus Mountains are
111 shown in Fig. 1 (c). After harvesting, *O. anatolica* terrestrial tubers were washed, followed by
112 poaching in boiling water for 12 min to remove the bitterness of their fresh state, with their
113 epidermis removed, after which they were dried in the shade for 7 d (Fig. 1 d). They were milled

114 at 600 rpm using a rotary mill (Break Mill SM 3, Brabender, Germany) until a fine powder was
115 finally obtained as described by Bulut-Solak and O'Mahony (2015).

116

117 *2.3. Preparation of dispersions*

118 Dispersions of *O. anatolica* tuber gum (*OaG*) were prepared at concentrations of 0.5%, 1.5%,
119 2.0% and 2.5% w/v by dispersing finely-ground *OaG* gum in deionised water, and stirred at low
120 speed using a magnetic stirrer for 2 h at 22°C until completely dispersed. All dispersions were
121 held for 18 h at 4°C to ensure complete hydration prior to assessment. Each dispersion was
122 removed from storage at 4°C and equilibrated at room temperature (22°C) for 15 min before the
123 pasting treatment.

124

125 *2.4. Analytical determinations*

126 *2.4.1. Particle size distribution*

127 The particle size distribution of the milled powders (Section 2.2) was analysed using a Malvern
128 Mastersizer 3000 with Aero S dry dispersion unit (Malvern Instruments, Worcestershire, UK) as
129 described by Amagliani, O'Regan, Kelly and O'Mahony (2016).

130

131 *2.4.2. Chemical composition*

132 Moisture, ash, fat, protein and crude fiber contents of the samples were determined according to
133 the standard methods of the Association of Analytical Chemists (AOAC, 2010). The samples
134 were analysed for moisture and ash contents using Gravimetric methods, fat content using the
135 Soxhlet method, and protein content using the Kjeldahl method using a total nitrogen to protein

136 conversion factor of 6.25 (AOAC, 2010). Total dietary fibre content was determined using a
137 commercial test kit from Megazyme International (K-TDRF-12/15, Bray, Ireland) based on
138 AOAC Method 991.43 and AACC Method 32-07.01. Samples were enzymatically treated with
139 heat-stable α -amylase, protease and amyloglucosidase, followed by a treatment with four
140 volumes of 95% ethanol to precipitate the fibre and remove depolymerised protein and glucose
141 from starch. The residue was filtered, washed, dried overnight and weighed (Megazyme
142 International Ireland Limited, 2015). Total reducing sugars (glucose, fructose and saccharose)
143 were analyzed with HPLC as described by Makila et al. (2014). pH values of the *OaG*
144 dispersions were measured at ambient temperature using a WTW digital pH meter equipped with
145 a glass electrode (Hach, H-Series H260G Bench-top pH and ISE Meter, Canada).

146

147 2.4.3. Starch content

148 The samples were prepared according to the AOAC procedure specified for total starch (AOAC
149 Method 996.11 and AACC Method 76-13.01) using the enzyme-based assay kit from Megazyme
150 International (Catalogue Number: AMG/AA, 05/16, K-TSTA-50A/K-TSTA-100A). The
151 absorbance of all samples and standard solutions was determined at 510 nm (Megazyme
152 International Ireland Limited, 2016) and the starch content of the samples was estimated using
153 the following formula:

$$154 \text{ Starch} = \Delta A \times (F/W) \times FV \times 0.9 \text{ [g/100 g]}$$

155 ΔA = Absorbance (reaction) read against the reagent blank

156 F = 100 (μg of D-glucose control) / absorbance value for 100 μg glucose (Conversion from
157 absorbance to μg)

158 W = weight of the sample (100 mg)

159 **FV**= Final volume

160

161 2.4.4. Glucomannan content

162 The glucomannan content of *OaG* was determined using the method of Chua et al. (2012). The
163 *OaG* was processed with physical procedures, such as removing epidermis, in order to reduce the
164 levels of impurities prior to drying (Bulut-Solak and O'Mahony 2015). A sample of *OaG* (200
165 mg wet basis) was weighed and added to formic acid-sodium hydroxide buffer (0.1 mol/L; 50
166 mL) and stirred magnetically at pH 3.13 for 4 h at 22 °C in order to separate the free sugars of the
167 *OaG* sample. The mixture was diluted with deionised water to 100 mL and then centrifuged at
168 4500 g for 40 min at 25°C. One sample was prepared as a blank for the tuber glucomannan
169 dispersion extract (TGE). Sulphuric acid (3 mol/L; 2.5 mL) was added to a flask containing 5 mL
170 of supernatant of TGE. The resultant dispersions were stirred and hydrolyzed for 90 min in a
171 boiling water bath, and cooled quickly to 22°C, and sodium hydroxide (6 mol/L; 2.5 mL) was
172 added. The solution was made up to 25 mL with deionised water to form the TGE hydrolysate
173 (TGEH). One sample was prepared as a blank for the TGEH. 3,5-dinitro salicylic acid (3,5-DNS;
174 1.5 mL of a 1% w/w reagent) was added to TGE and TGEH solutions (2 mL) and heated in a
175 boiling water bath for 5 min before being cooled quickly to 22°C. Deionised water was added to
176 dilute the samples to a volume of 25 mL. The absorbances of the TGE and the TGEH samples
177 were measured at 550 nm with deionised water used as a blank (Chua et al. 2012). D-glucose and
178 D-mannose stock solutions (1 mg/mL; 0.4 mL, 0.8 mL, 1.2 mL, 1.6 mL, 2.0 mL) were
179 transferred into eleven 25 mL volumetric flasks, as well as 2 mL of deionised water as a blank.
180 Deionised water was added to a volume of 2 mL, followed by the addition of 3,5-DNS (1% w/w;
181 1.5 mL) to each volumetric flask. This mixture was heated for 5 min in a boiling water bath and

182 cooled to 22°C quickly before diluting to 25 mL with deionised water. The absorbance of the
183 standard solutions was read at 550 nm (Chua et al. 2012). A D-mannose standard curve was
184 constructed using the procedure as described for glucose, and used as a correction factor in Eq. 1.
185 The glucomannan content of the samples was determined according to Eq. (1),

$$\text{Glucomannan (\%)} = \frac{5000 f \times (5T - T_0)}{m} \quad (\text{Eq. 1})$$

186 **f** is the correction factor obtained from the glucose and mannose standards

187 **T** is the glucose content (mg) in the tuber glucomannan dispersion extract hydrolysate (TGEH)
188 obtained from the standard curve

189 **T₀** is the glucose content (mg) in the tuber glucomannan dispersion extract (TGE) obtained from
190 the standard curve

191 **m** is the mass of the *OaG* sample (200 mg wet basis).

192

193 2.4.5. Pasting behaviour

194 The pasting properties of all *OaG* dispersions were determined using an AR-G2 controlled-stress
195 rheometer equipped with a starch pasting cell (AR-G2; TA Instruments Ltd., Waters LLC,
196 Leatherhead, Surrey, UK); the internal diameter of the cell was 36.0 mm, the diameter of the
197 rotor was 32.4 mm, and the gap between the two elements at the base of the geometry was 0.55
198 mm. All measurements of viscosity were carried out at a fixed shear rate of 16.8 s⁻¹. Analysis was
199 conducted as described by Kett et al. (2013), whereby, after loading onto the rheometer, the
200 sample was equilibrated at 35°C for 1 min, heated from 35 to 95°C over 4 min, held at 95°C for 6
201 min, cooled from 95 to 35°C within 4 min and held for 5 min at 35°C.

202

203 *2.4.6. Microstructural analyses*204 *2.4.6.1. Scanning electron microscopy*

205 Scanning electron microscopy (SEM) analysis was carried out in the Biosciences Imaging
206 Centre, Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland. *OaG*
207 powder was mounted on aluminium stubs using double-sided adhesive carbon tape, and sputter
208 coated with a 5 nm layer of gold/palladium (Au:Pd = 80:20) using a Quorum Q150R ES Sputter
209 Coating Unit (Quorum Technologies Ltd., Sussex, U.K.) to prevent surface charging by the
210 electron beam. Subsequently, the *OaG* sample was loaded into a sample tube and then examined
211 using a JSM-5510 scanning electron microscope (JEOL Ltd, Tokyo, Japan), operated at an
212 accelerating voltage of 5 kV.

213

214 *2.4.6.2. Light microscopy*

215 The *OaG* dispersions were monitored using polarised light on an Olympus B×51 light
216 microscope (LM) (Olympus Corporation, Tokyo, Japan) fitted with a heating stage (CO 102;
217 Linkam Scientific Instruments, Tadworth, Surrey, UK). After placing on the microscope, all
218 samples were equilibrated at 35°C for 2 min, heated from 35 to 95°C at 10°C min⁻¹ and tempered
219 at 95°C over a period of 6 min before cooling from 95 to 35°C for 5 min. A magnification of 10×
220 was used for microscopy analysis.

221

222 *2.5. Statistical data analysis*

223 All samples were prepared freshly on three separate occasions, with all analytical measurements
224 conducted in triplicate. The results are given as mean values ± standard deviations. One-way

225 analysis of variance (ANOVA) was performed for statistical analysis. SAS 9.1.3 was used for all
226 analyses (SAS Institute Inc., Cary, NC, USA). Duncan's multiple comparison test was used to
227 determine significant differences between the various treatments and results with $P < 0.05$ were
228 considered significantly different.

229

230 **3.0. Results and Discussion**

231

232 *3.1. Physical and chemical properties*

233 The volume mean diameter ($D [4,3]$) of the milled *OaG* powder was $180 \pm 1.25 \mu\text{m}$. While 10% of
234 the powder particles ($D_v [10]$) had diameter less than $18.6 \pm 0.22 \mu\text{m}$, 90% of the powder
235 particles ($D_v [90]$) had diameter less than $406 \pm 1.63 \mu\text{m}$. Surface weighted mean particle
236 diameter of *OaG* powder ($D[3,2]$) was $37 \pm 0.76 \mu\text{m}$. Specific surface area (SSA) of the *OaG* was
237 also $161.8 \pm 0.00 (\text{m}^2\text{kg}^{-1})^{\text{g}}$. The chemical composition of *OaG* is presented in Table 1. *OaG*
238 contained 41.6% starch, 32.3% dietary fiber, 18.5% glucomannan, 1.52% crude fiber, 0.19%
239 glucose, 2.34% saccharose, 6.73% protein, 7.73% moisture, 2.25% ash and 0.28% fat; no
240 fructose was found in *OaG*. The mean pH value of the *OaG* dispersions was 6.11 ± 0.02 . The other
241 undetermined compounds in *OaG* would possibly be other minor reducing sugars which are
242 related to the raffinose series frequently found in photosynthetic tissues (Avigad & Dey, 1997).
243 These compounds also represent low molecular weight heteropolymers associated with the
244 metabolism of mucilaginous polysaccharides, as noted in some temperate orchid species
245 (Buchala, Franz, & Meier, 1974; Franz, 1979; Meier & Reid, 1982).

246

247 The compositional profile of *OaG* is similar to some previous results published by Farhoosh and
248 Rizai (2007), and Tekinsen and Guner (2010). Farhoosh and Rizai (2007) reported that round-
249 edged tuber gum contained 19.3% glucomannan, 6.85% starch, 13.2% moisture, 7.35% protein
250 and 2.8% ash whereas, Tekinsen and Guner (2010) reported that *OaG* contained 43.5%
251 glucomannan, 14.7% starch, 10.7% moisture, 3.20% protein, 1.94% ash and had a pH of 5.71.
252 Lalika et al. (2013) reported that edible orchids had 5.36% protein, 2.7% crude fiber, 2.2% ash,
253 1.57% fat and 0.09 mg/100g vitamin C. Moreover, Tekinsen and Guner (2010) also reported that
254 the mean values of compositional contents of Salep gum, depending on the different species in
255 the Orchidaceae family, were 17.7-54.6% glucomannan, 5.4-38.7% starch, 0.95-2.83% ash, 9.35-
256 12.4% moisture with pH ranging from 5.61-6.20.

257 Starch serves as an energy reserve in tubers (Buckeridge, 2010). When orchid tubers are fully
258 formed, they contain maximal levels of starchy matter to supply energy to the perennial plant
259 during winter (Hossain, 2011). As indicated in Table 1, the main component of *OaG* was starch.
260 Additionally, the high level of starch (41.6%) in the *OaG* could be possibly due to the fact that
261 the plants were harvested in the period June-July. *OaG* had high levels of dietary fiber (32.3%),
262 glucomannan (18.5%) and crude fiber (1.52%), which are considered to have positive health
263 benefits (Nishinari, 2000). Previous work has also shown that Turkish orchid tubers generally
264 had total dietary fiber content ranging from 11.6-40.1% (Gumus, 2009). When compared to some
265 previous results, if the tubers are harvested early at the flowering stage, the tubers contain less
266 starch and higher glucomannan, because the formation of starch in the tubers takes place mainly
267 between March to July (Buchala et al., 1974); it should be noted that there is a negative
268 correlation between starch and glucomannan contents in *OaG* (Tekinsen and Guner 2010).

269 *OaG* had relatively high protein content (6.73%), but there is a considerable variation for protein
270 values reported previously. For example, Tekinsen and Guner (2010) reported that the protein
271 content in *OaG* was 3.20%. The low protein content reported in previous study may arise from
272 using the generic nitrogen to protein conversion factor of 5.70 (Tkachuk, 1969). Additionally,
273 *OaG* had 7.73% moisture and 2.25% ash (Table 1). Even if depending on the period of the drying
274 process during summer, the ash and moisture contents were similar to other previous results
275 because all dried tubers were held until their milky appearance changed to almost a semi-
276 transparent yellowish rough state. Sezik (1967) also reported that if the moisture content of Salep
277 gum was lower than 10%, it could result in enhanced stability and a longer shelf life. Moreover,
278 *OaG* contained trace levels of fat. However, Citil and Tekinsen (2011) also reported that the
279 mean fat content in Salep gum was 2.02%, but this could be due to boiling the tubers in milk
280 before drying, and the presence of milk fat may possibly have increased the fat content in Salep
281 gum.

282

283 3.2. Morphological structures of native *OaG*

284 The morphological properties of native *OaG* milled powder particles are presented in Fig. 2. It
285 was found that the *OaG* particles had irregular shapes. Some fracture shapes of the *OaG* particles
286 were probably caused by mechanical breakdown of *OaG* during milling. When the surface of the
287 *OaG* particles was focussed and magnified at the highest level, multilayered surfaces of the *OaG*
288 were not well observed. Moreover, the smooth surface of *OaG* with round edges was clearly
289 observed in Fig. 2, even if a disadvantage of SEM for the sample preparation, especially drying
290 and metal coating, could slightly limit the visualisation to the *OaG* particles in their original
291 environment. However, some irregular shaped particles with partly roughened surfaces were also

292 observed at the highest magnification level. Rough surfaces of the particles may have been
293 generated by partial gelatinisation of the starch granules/containing particles of *OaG* during pre-
294 treatment (i.e., poaching in water prior to drying) of the tubers. Similarly, lamellar surfaces of
295 konjac glucomannan have also been reported by Cheng, Abd Karim and Seow (2011). Razavi,
296 Nyamathulla, Karimian and Noordin (2014) also observed the morphological properties of
297 palmate tubers of *Orchis morio var mascula*, which were reported to be irregular rod-shaped
298 particles with roughened surfaces.

299

300 3.3. Pasting properties of *OaG*

301 The pasting treatment was applied to the dispersions of *OaG* at varying concentrations. The mean
302 values of the initial viscosity, peak viscosity, and viscosity recorded at the end of the holding
303 period at 95°C, on completion of cooling to 35°C, and at the end of the final holding period at
304 35°C are presented in Table 2. The pasting curves obtained from the dispersions of *OaG* are also
305 shown in Fig. 3. The pH values of all dispersions were similar (6.11 ± 0.01) ($P > 0.05$).

306 As seen in Table 2 and Fig 3, the values of initial viscosity of the dispersions at varying
307 concentrations ranged from 33.7 ± 0.24 to 1193 ± 92.0 mPa.s, whereas the values of peak viscosity
308 ranged from 23.0 ± 0.68 to 702 ± 59.1 mPa.s ($P < 0.05$). After the cooling period at 35°C, the final
309 viscosity values of the *OaG* dispersions ranged from 34.3 ± 0.45 to 1437 ± 83.3 mPa.s, depending
310 on the concentration of the dispersions (Table 2). At the end of the starch pasting treatment, the
311 viscosity values of the final pastes at 35°C were higher compared to the initial viscosity values
312 (Table 2 and Fig. 3). These higher viscosity values were due to the swelling of starch granules
313 during pasting. It should also be noted that while fresh orchid tubers were being poached in

314 boiling water, partial, limited gelatinisation/pasting of starch may have taken place as part of this
315 pre-treatment process.

316 Moreover, the temperatures of the peak viscosity of the dispersions ranged from $94.2\pm 0.64^{\circ}\text{C}$ to
317 $97.4\pm 0.35^{\circ}\text{C}$. At the lowest *OaG* concentration (0.5%), the starch granules were more free to
318 expand during the gelatinisation process, so that the peak viscosity was seen at a lower
319 temperature ($94.2\pm 0.64^{\circ}\text{C}$) while the temperature of the peak viscosity of the dispersion at the
320 highest concentration (2.5%) occurred at a significantly higher temperature ($95.4\pm 0.32^{\circ}\text{C}$) ($P <$
321 0.05). For the *OaG* dispersion at 2.5%, a possible reason for lower peak temperature compared to
322 the 1.5 and 2.0% *OaG* samples could also be an increase in the effective concentration of dietary
323 fiber and glucomannan which has a relatively high water holding capacity, which could limit the
324 swelling of starch granules because of reducing amount of water available for gelatinisation. It
325 could likely prevent both dissolution and association of primarily AM, as well as AP. These
326 findings hypothesise that paste characteristics of *OaG* could be affected by glucomannan in the
327 continuous phase. Interactions between starch and glucomannan, as well as dietary fiber could
328 also possibly have influenced pasting properties, with respect to changes in paste viscosity and
329 gelatinisation temperature. Moreover, a delay starch gelatinization could be also due to the
330 interaction between dietary fiber and starch, because dietary fiber with the presence of high
331 amount of hydroxyl groups has a great water binding capacity, and can be attributed to a
332 reduction in water availability which causes partial gelatinization of crystalline regions in the
333 starch granules (Funami et al., 2005). This interaction could also lower viscosity and reduce
334 pasting. Jiang and Ramsden also reported that effect of this interaction is dependent upon starch
335 and dietary fiber concentration (1999). Nagano et al. (2008) reported that guar gum in a maize
336 starch suspension (5%) inhibited starch components from leaching out of the granules to the

337 continuous phase of starch pastes during gelatinisation, to an extent dependant on the
338 concentration of added guar gum. However, Tester and Sommerville (2003) reported that the
339 presence of polymers and/or other compounds in the liquid phase has no effect on swelling
340 volume until the amount of available water becomes a limiting factor. The swelling properties of
341 the starch granules in *OaG* were probably offset to higher temperatures as a result of the chemical
342 composition of *OaG*.

343

344 *3.4. Visual assessment of pasting properties of OaG using light microscopy*

345 The effect of gelatinisation on heating of starch granules in *OaG* was explored further using a
346 polarised LM fitted with a heating stage. Several micrographs of the dispersions of *OaG* prepared
347 freshly at varying concentrations from 0.5 to 2.5% were taken at different temperatures using LM
348 and these micrographs were recorded from the middle of the sample in each slide (Fig. 4). The
349 arrows in the each micrograph clearly show selected starch granules/particles in the *OaG*
350 dispersions during pasting (Fig. 4).

351 As seen in the micrographs, starch granules in *OaG* particles began to swell gradually during
352 heating and the swelling of the starch granules greatly increased between 65 and 75°C. Moreover,
353 the changes in the size of *OaG* particles increased when the temperature reached 65°C,
354 demonstrating that gelatinisation had already begun. However, the remaining intact granule
355 ghosts in *OaG* were further swollen at the end of the holding period of the 95°C, when compared
356 to 75°C (Fig. 4). Other important factors affecting the gelatinisation could be the amount of
357 available water in the dispersions, the nature of starch and the starch: glucomannan ratio in *OaG*
358 as they could affect the relative importance of swelling of starch granules in the discontinuous
359 and continuous phases of pastes/gels.

360 In addition, lipids and proteins in plants, as well as tubers, are known to be associated with both
361 the surface layer and the interior of starch granules. The protein and lipid in the surface layer of
362 starch granules also prevent the swelling of starch granules, due to the fracturing of a restricting
363 layer of lipid and protein at the surface of the starch granules (Debet & Gidley, 2007). The
364 protein content of starch granules in the *OaG* particles could also slightly influence the swelling
365 property of the granules during gelatinisation.

366

367 **4.0. Conclusion**

368

369 *OaG* has a unique and interesting chemical composition and is mainly composed of starch,
370 dietary fiber and glucomannan. The pasting behaviour of *OaG* appears to be influenced by its
371 chemical composition, in particular the content of the high molecular weight polysaccharide,
372 glucomannan and dietary fiber. These glucomannan and dietary fiber components, and their
373 possible interactions with starch, in *OaG* may have resulted in higher peak pasting temperatures
374 and lower peak viscosity, due to limitation of the leaching of AM and AP from starch granules.
375 These findings provide novel practical information on the role and potential usefulness of *OaG* as
376 an ingredient in controlling rheology and modifying texture of foodstuffs. Consequently, *OaG*-
377 based ingredients may be used for enhancing the functional properties of food formulations,
378 which may provide interesting opportunities to the food industry in the development of new
379 product structures.

380

381 **5.0. Acknowledgements**

382

383 This research was carried out at University College Cork, supported by The Turkish Scientific
384 and Technological Research Council (TUBITAK) (B.14.2.TBT.0.06.01-219-84, 2013) and the
385 Turkish Ministry of Food, Agriculture and Livestock (53231444-110.05-17969, 2013). The
386 authors appreciate the assistance of Dr. Ozlem Cetin at the Biotechnology Department in Selcuk
387 University for identification of *Orchis anatolica*. The authors are grateful to Suzanne Crotty at
388 the BioSciences Imaging Centre, Department of Anatomy and Neuroscience, University College
389 Cork for assistance with the scanning electron microscopy analysis and to David Waldron at the
390 School of Food and Nutritional Sciences, University College Cork for assistance with the light
391 microscopy analysis.

392

393 **6.0. References**

394

395 Alonso-Sande, M., Teijeiro-Osorio, D., Remuñán-López, C., & Alonso, M. J. (2009).
396 Glucomannan, a promising polysaccharide for biopharmaceutical purposes. *European Journal of*
397 *Pharmaceutics and Biopharmaceutics*, 72, 453-462.

398

399 Altundag, E., Sevgi, E., Kara, O., Sevgi, O., Tecimen, H. B., & Bolat, I. (2012). Studies on the
400 chorology, ecology, morphology and conservation strategies of *Orchis anatolica* Boiss
401 (Orchidaceae). *Journal of Environmental Biology*, 33, 325-333.

402

403 AOAC (2010). Official methods of analysis of analysis of the association of official analytical
404 chemists (18th ed.). *Association of Official Analytical Chemist AOAC*, Washington, DC, USA.

405

406 Amagliani L., O'Regan J., Kelly A. L., & O'Mahony J. A. (2016). Physical and flow properties
407 of rice protein powders. *Journal of Food Engineering* 190, 1-9.

408

- 409 Atkin, N. J., Abeysekera, R. M., & Robards, A. W. (1998). The events leading to the formation
410 of ghost remnants from the starch granule surface and the contribution of the granule surface to
411 the gelatinization endotherm. *Carbohydrate Polymers*, *36*, 193-204.
- 412
413 Avigad, G., & Dey, P. M. (1997). Carbohydrate Metabolism: Storage Carbohydrates. In P. M.
414 Dey, & J. B. Harborne (Eds). *Plant Biochemistry* (pp. 143-204), London: Academic Press.
- 415
416 Baldwin, P. M. (2001). Starch granule-associated proteins and polypeptides: A review
417 *Starch/Staerke*, *53*, 475-503.
- 418
419 Bardajee, G. R., Hooshyar, Z., & Kabiri, F. (2012). Preparation and investigation on swelling and
420 drug delivery properties of a novel silver/Salep-g-poly(acrylic acid) nanocomposite hydrogel.
421 *Bulletin of the Korean Chemical Society*, *33*, 2635-2641.
- 422
423 Bardajee, G. R., Hooshyar, Z., Asli, M. J., Shahidi, F. E., & Dianatnejad, N. (2013). Synthesis of
424 a novel supermagnetic iron oxide nanocomposite hydrogel based on graft copolymerization of
425 poly [(2-dimethylamino) ethyl methacrylate] onto salep for controlled release of drug. *Materials*
426 *Science & Engineering C*, doi: 10.1016/j.msec.2013.11.022.
- 427
428 Bulut-Solak, B., & O'Mahony, J. A. (2015). Rheological and microstructural properties of
429 rennet-induced milk protein gels formed with *Orchis anatolica* gum. *International Dairy*
430 *Journal*, *51*, 59-64.
- 431
432 Buchala, A. J., Franz, J., & Meier, H. (1974). A glucomannan from the tubers of *Orchis morio*.
433 *Phytochemistry*, *13*, 163-166.
- 434
435 Buckeridge, M. S. (2010). Seed cell wall storage polysaccharides: Models to understand cell wall
436 biosynthesis and degradation. *Plant Physiology*, *154*, 1017-1023.
- 437
438 Cheng, L. H., Abd Karim, A., & Seow, C. C. (2007). Effects of acid modification on physical
439 properties of konjac glucomannan (KGM) films. *Food Chemistry*, *103*, 994-1002.

- 440
441 Chua, M., Chan, K., Hocking, T. J., Williams, P. A., Perry, C. J., & Baldwin, T. C. (2012).
442 Methodologies for the extraction and analysis of konjac glucomannan from corms of
443 *Amorphophallus konjac* K. Koch. *Carbohydrate Polymers*, 87, 2202-2210.
444
445 Cital, O. B., & Tekinsen, K. K. (2011). A comparative study on fatty-acid composition of Salep
446 obtained from some Orchidaceae species. *Chemistry of Natural Compounds*, 46, 943-945.
447
448 Considine, T., Noisuwan, A., Hemar, Y., Wilkinson, B., Bronlund, J. & Kasapis, S. (2011).
449 Rheological investigations of the interactions between starch and milk proteins in model dairy
450 systems: A review. *Food Hydrocolloids*, 25, 2008-2017.
451
452 Debet, M. R., & Gidley, M. J. (2006). Three classes of starch granule swelling: Influence of
453 surface proteins and lipids. *Carbohydrate Polymers*, 64, 452-465.
454
455 Debet, M. R., & Gidley, M. J. (2007). Why do gelatinised starch granules not dissolve
456 completely? Roles for amylose, protein and lipid in granule “ghost” integrity. *Journal of*
457 *Agricultural and Food Chemistry*, 55, 4752-4760.
458
459 Farhoosh, R., & Riazi, A. (2007). A compositional study on two current types of Salep in Iran
460 and their rheological properties as a function of concentration and temperature. *Food*
461 *Hydrocolloids*, 21, 660-666.
462
463 Franz, G. (1979). Metabolism of reserve polysaccharides in tubers of *Orchis morio*. *Planta*
464 *Medica*, 36, 68-73.
465
466 Funami, T., Kataoka, Y., Omoto, T., Gota, Y., Asai, I., & Nishinari, K. (2005). Effects of non
467 ionic polysaccharides on the gelatinization and retrogradation behavior of wheat starch. *Food*
468 *Hydrocolloids*, 19, 1-13.
469

- 470 Georgiadis, N., Ritzoulis C., Charchari, E., Koukiotis, C., Tsiptsias, C., & Vasiliadou, C.
471 (2012). Isolation, characterization and emulsion stabilizing properties of polysaccharides from
472 orchid roots (Salep). *Food Hydrocolloids*, 28, 68-74.
- 473
- 474 Gumus, C. (2009). Bati Karadeniz Bolgesinde Salep elde edilmesinde kullanılan bazı orkide
475 türlerinin (*Orchidaceae*) cogaltım yöntemleri üzerinde araştırmalar. PhD Thesis, Natural Science
476 Institute, Ankara University, 1-229.
- 477
- 478 Harding, S. E., Abdelhameed, A. S., & Morris, G. A. (2010). Molecular weight distribution
479 evaluation of polysaccharides and glycoconjugates using analytical ultracentrifugation.
480 *Macromolecular Bioscience*, 10, 714-720.
- 481
- 482 Hossain, M. M. (2011). Therapeutic orchids: Traditional uses and recent advances-An overview-
483 A review. *Fitoterapi*, 82, 102-140.
- 484
- 485 Kett, A. P., Chaurin V., Fitzsimons, S. M., Morris, E. R., O'Mahony, J. A., & Fenelon, M. A.
486 (2013). Influence of milk proteins on the pasting behaviour and microstructural characteristics of
487 waxy maize starch. *Food Hydrocolloids*, 30, 661-671.
- 488
- 489 Jiang, G., & Ramsden, L. (1999). Characterization and yield of the arabinogalactanprotein
490 mucilage of taro corms. *Journal of the Science of Food and Agriculture*, 79, 671-674.
- 491
- 492 Lalika, M. C. S., Mende D. H., Urio, P., Gimbi, D. M., Mwanyika, S. J., & Donati, G. (2013)
493 Domestication potential and nutrient composition of wild orchids from two southern regions in
494 Tanzania. *Time Journals of Biological Sciences and Technology*, 1 (1) 1-11.
- 495
- 496 Lange, D. (1998). Europe's medicinal and aromatic plants: Their use, trade and conservation.
497 Cambridge. Traffic Europe/International, Cambridge, United Kingdom, 10.
- 498

- 499 Makila, L., Laaksonen, O., Diaz, J. M. R., Vahvaselka, M., Myllymaki, O., Lehtomaki, I.,
500 Laakso, S., Jahreis, G., Jouppila, K., Larmo, P., Yang, B., & Kallio, H. (2014). Exploiting
501 blackcurrant juice press residue in extruded snacks. *LWT - Food Science and Technology*, 57,
502 618-627.
- 503
- 504 Megazyme International Ireland Limited. (2015). Total dietary fiber, 12/15, K-TDFR-100A/K-
505 TDFR-200A. Ireland.
- 506
- 507 Megazyme International Ireland Limited. (2016). Total starch assay procedure
508 (amyloglucosidase/ α -Amylase Method) AA/AMG, 05/16, K-TSTA-50A/K-TSTA-100A. Ireland.
- 509
- 510 Meier, H., & Reid, J. S. G. (1982). Reserve polysaccharides other than starch in higher plants. In
511 F. A. Loewus, & W. Tanner (Eds). *Encyclopedia of Plant Physiology* (pp. 418-471), Springer-
512 Verlag, Berlin, 13A.
- 513
- 514 Nagano, T., Tamaki, E., & Funami, T. (2008). Influence of guar gum on granule morphologies
515 and rheological properties of maize starch. *Carbohydrate Polymers*, 72, 95-101.
- 516
- 517 Nikjooy, S., Joo, M. G., & Jahanshahi, S. S. (2014). The effect of various concentrations of
518 Salep gum on physicochemical characteristics of low-fat white cheese, *International Journal of*
519 *Agriculture and Crop Sciences*, 8, 5, 136-141.
- 520
- 521 Nishinari, K. (2000). Konjac glucomannan, *Developments in Food Science*, 41, 309-330.
- 522
- 523 Pourjavadi, A., Doulabi, M., Soleyman, R., Sharif, S., & Egtesadi, S. A. (2012). Synthesis and
524 characterization of a novel (Salep phosphate)-based hydrogel as a carrier matrix for fertilizer
525 release. *Reactive and Functional Polymers*, 72, 667-672.
- 526
- 527 Ratnayake, W. S., & Jackson, D. S. (2009). Starch gelatinization. *Advances in Food and*
528 *Nutrition Research*, 55, 221-268.
- 529

- 530 Razavi, M., Nyamathulla, S., Karimian, H., & Noordin, M. I. (2014). Novel swellable polymer of
531 Orchidaceae family for gastroretentive drug delivery of famotidine. *Drug Design, Development
532 and Therapy*, 8, 1315-1329.
- 533
- 534 Salemis, P., & Rinaudo, M. (1984). Gel permeation chromatography of starch and other
535 uncharged polysaccharides. *Polymer Bulletin*, 11, 397-400.
- 536
- 537 Sezik, E. (1967). Turkiye'nin Salepgilleri ticari Salep cesitleri ve ozellikle Mugla Salebi uzerinde
538 arastirmalar. PhD Thesis, No 34, Istanbul University, Istanbul.
- 539
- 540 Tamer, C. E., Karaman, B., & Copur, O.U. (2006). A traditional Turkish Beverage: Salep. *Food
541 Reviews International*, 22, 43-50.
- 542
- 543 Tekinsen, K. K., & Guner, A. (2010). Chemical composition and physicochemical properties of
544 tubera Salep produced from some Orchidaceae species. *Food Chemistry*, 121, 468-471.
- 545
- 546 Tester, R. F., & Sommerville, M. D. (2003). The effects of non-starch polysaccharides on the
547 extent of gelatinisation, swelling and amylase hydrolysis of maize and wheat starches. *Food
548 Hydrocolloids*, 17, 41-54.
- 549
- 550 Tkachuk, R. (1969). Nitrogen to protein conversion factors for cereals and oilseed meals. *Cereal
551 Chemistry*, 46, 419-423.
- 552
- 553
- 554
- 555
- 556
- 557
- 558
- 559
- 560
- 561
- 562

563 **Figure Legends**

564
565 **Figure 1 (a-d).** Photographs of *Orchis anatolica* harvested from the Taurus Mountains
566 All : (a-b) *Orchis anatolica*; (c) fresh round-edged tubers of *O. antolica*; dried tubers of *O.*
567 *antolica*.

568 **Figure 2.** Scanning electron micrographs of native *OaG* granules at magnifications of
569 100 x (a), 250 x (b), 1000 x (c) and 2000 x (d).

570
571 **Figure 3.** Pasting curves for *Orchis anatolica* gum at concentrations of 0.5 % (—), 1.5% (Δ), 2%
572 (o) and 2.5% (*) at pH 6.11 ± 0.01 . The dotted lines without symbols show temperature.

573
574 **Figure 4** The micrographs of *OaG* at varying concentrations. All: (a) 2.0% *OaG*; (b) 2.5% *OaG*;
575 at different temperatures viewed under polarised light. Left-hand column at 35 °C, middle-
576 columns from 55 to 95 °C on heating, right-hand column at 47 °C on cooling. Magnification 10 \times .
577 Scale bar 200 μm .

1 **Table 1.** Chemical composition and pH value of *Orchis anatolica* tuber gum

Composition	<i>O. anatolica</i> tuber gum (g/100 g)	Range*
Starch	41.6±1.52	5.40-38.7
Total dietary fibre	32.3±1.44	11.6-40.1
Glucomannan	18.5±0.33	17.7-54.6
Crude fibre	1.52±0.03	2.70
Protein	6.73±0.22	3.20-7.35
Ash	2.25±0.04	0.95-2.83
Fat	0.28±0.08	2.02
Moisture	7.73±0.51	9.35-13.2
Glucose	0.19±0.01	-
Saccharose	2.34±0.02	-
Fructose	Not founded	-
pH	6.11±0.02	5.71-6.20

2
3 Data presented are means of three replicates ± the standard deviations.

4 *Range values based on previous published information from Farhoosh and Rizai (2007), Gumus (2009), Tekinsen
5 and Guner (2010) and Lalika et al. (2013).
6

7
8
9
10
11
12
13
14
15

16 **Table 2.** Viscosity (mPa.s) of *O. anatolica* tuber gum dispersions at various stages of the
 17 pasting regime

Stage of pasting	0.5%	1.5%	2.0%	2.5%
Initial viscosity (mPa.s)	33.7±0.24 ^d	306±8.91 ^c	622±40.5 ^b	1193±92.0 ^a
Peak viscosity (mPa.s)	23.0±0.68 ^c	177±97.5 ^c	481±178 ^b	702±59.1 ^a
End of holding at 95 °C (mPa.s)	16.4±0.09 ^d	93.0±4.12 ^c	217±24.5 ^b	494±47.8 ^a
End of cooling to 35 °C (mPa.s)	32.0±0.38 ^d	256±11.6 ^c	566±43.9 ^b	1130±72.9 ^a
Final paste at 35 °C (mPa.s)	34.3±0.45 ^d	328±13.5 ^c	751±58.1 ^b	1437±83.3 ^a
Temperature of peak viscosity (°C)	94.2 ±0.64 ^c	97.4±0.35 ^a	96.8±0.14 ^a	95.4±0.32 ^b

18
 19 Data presented are means of three replicates ± the standard deviations.

20 Means (n=3) within the same column with different superscript letters differ significantly (P < 0.05).

21
 22
 23

1

2 **Figure 1 (a-d).**

3

4

5

6

7

8

9

10

11

12

13

14


15

16

17

18

19



20 **Figure 2.**

21

22

23


24

25

26

27

28



29


30 **Figure 3.**

31

32

33

34



35

36

37

Figure 4.

1 **Research Highlights**

2

3 Milled *OaG* had irregular shaped particles with round edges

4 The main components of *OaG* were starch, dietary fiber and glucomannan

5 The starch pasting behaviour of *OaG* was influenced by the glucomannan and dietary fiber
6 components

7 *OaG* is a potent viscosity modifier and thickener for food applications