CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Temporal sequence of muscle activation in the glute bridge exercise
Authors	Scheurer, Sebastian;Tedesco, Salvatore;O'Flynn, Brendan;Brown, Kenneth
Publication date	2025-09-04
Original Citation	Scheurer, S., Tedesco, S., O'Flynn, B. and Brown, K. (2025) 'Temporal sequence of muscle activation in the glute bridge exercise', Gait & Posture, 121, pp.220-221. https://doi.org/10.1016/j.gaitpost.2025.07.236
Type of publication	Article (peer-reviewed);journal-article
Link to publisher's version	10.1016/j.gaitpost.2025.07.236
Rights	© 2025, Elsevier B.V. For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. - https://creativecommons.org/licenses/by/4.0/
Download date	2026-10-09 11:16:42
Item downloaded from	https://hdl.handle.net/10468/18200

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Temporal sequence of muscle activation in the glute bridge exercise

Sebastian Scheurer[1] Salvatore Tedesco[2]
Brendan O'Flynn[2] Kenneth N. Brown[1]

[1] Insight Centre for Data Analytics, School of Computer Science and Information Technology, University College Cork, T12 XF62 Cork, Ireland

[2] Tyndall National Institute, University College Cork, T12 R5CP Cork, Ireland

Introduction

The glute bridge (GB) is a foundational exercise strengthening lower posterior muscles and valuable for fitness, strength training, rehabilitation, and physical therapy. While the contributions of the muscles involved in GB have been investigated (Gasibat et al., 2023; Lehecka et al., 2017; Youdas et al., 2017), the temporal sequence of their activation remains unexplored.

Research Question

To address this gap, we conducted a statistical analysis of 1,272 muscle onsets from surface electromyography (EMG) data.

The central research question is: What insights can be gained from the statistical analysis of muscle onset timings in lower posterior muscles during glute bridges?

Methods

We recruited five healthy participants (age 29–38, three male, two female). EMG data were collected using a FREEEMG system (BTS Bioengineering) at 1,000 Hz. Electrodes were attached to left and right rectus femoris (RF), gluteus maximus (GM), erector spinae (ES), and biceps femoris (BF), following the SENIAM guidelines (http://seniam.org/). Participants completed 1–4 cycles of 20 GBs, contributing a total of 159 GBs. The study was approved by the Ethics Committee at Setanta College Ltd.

We processed EMG data and determined muscle onsets following Schuermans et al. (2017). Noise levels for each test cycle were estimated as the standard deviation of a three-second resting period with all eight muscles at rest. Muscle onsets were defined as the first 0.1 s with signal consistently above 3× noise level. The resulting 1,272 muscle onsets, shown in Fig. 1, were analysed with a linear

mixed effects model (Bates et al., 2015) of the form `Onset ~ Muscle * Side + (1|Participant/Rep)`.

Results

The fixed effects from Table 1 with confidence intervals (CIs) that do not span zero show that the typical GB is initiated by the left BF and both ESs and GMs, all of which activate in the first 0.1 s. The right BF activates 0.1 s later, followed within 0.25 s by the left and finally, <0.1 s later, the right RF. Random effects (Table 2) revealed that 13.4% of variance in onset timing was due to participants, 22.1% to repetitions, and 64.5% remained as residual variance (SD = 0.127 s).

Discussion

Findings indicate that GB muscle activation sequence is remarkably consistent both within and across individuals. This contrasts with other posterior muscle exercises, such as prone hip extension (Bonacci et al., 2009; Carlos et al., 2024; Kiseljak & Medved, 2022; Lirio-Romero et al., 2021; Suehiro et al., 2014) where onset timing varies and has been controversially linked to low back pain or hamstring injury risk (Emami et al., 2014; Ryan & Bruno, 2024; Schuermans et al., 2017). The consistency in GB activation timing suggests it may provide a more stable basis for injury risk modelling and clinical assessment compared to other posterior chain exercises.

Funding

This publication has emanated from research supported by a research grant from the Disruptive Technologies Innovation Fund (DTIF) project HOLISTICS funded by Enterprise Ireland (EI). Aspects of this work have been supported by Science Foundation Ireland under grant 12/RC/2289-P2-INSIGHT, and 13/RC/2077-CONNECT, which are cofounded by the European Regional Development Fund (ER 700 DF) under Ireland's European Structural and Investment Funds Programme 2014–2020. For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

References

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Bonacci, J., Chapman, A., Blanch, P., & Vicenzino, B. (2009). Neuromuscular adaptations to training, injury and passive interventions. *Sports Medicine*, *39*(11), 903–921. https://doi.org/10.2165/11317850-000000000-00000

Carlos, C. Q., Pessina, A. L. G., & Rodrigues Martinho Fernandes, L. F. (2024). Women with low back pain do not show a pattern in the lumbopelvic muscle activation sequence that differentiates from women without low back

pain. *Journal of Bodywork and Movement Therapies*, *40*, 2218–2226. https://doi.org/10.1016/j.jbmt.2023.07.003

Emami, M., Arab, A. M., & Ghamkhar, L. (2014). The activity pattern of the lumbo-pelvic muscles during prone hip extension in athletes with and without hamstring strain injury. *International Journal of Sports Physical Therapy*, *9*(3), 312–319.

Gasibat, Q., Alexe, C. I., Raveica, G., Tohănean, D. I., Vasilios, K., & Alexe, D. I. (2023). Decoding hip muscle activation: A comparative electromyographic analysis of turn-out bent knee pulse and single-leg banded glute bridge exercises in healthy female subjects. *European Journal of Investigation in Health, Psychology and Education*, *13*(9), 1612–1623. https://doi.org/10.3390/ejihpe13090117

Kiseljak, D., & Medved, V. (2022). Motor recruitment pattern during the prone hip extension test: Is hip extension initiated by the hip or the lumbar extensor muscles? *International Journal of Therapy and Rehabilitation*, *29*(5), 1–11. https://doi.org/10.12968/ijtr.2020.0139

Lehecka, B. J., Edwards, M., Haverkamp, R., Martin, L., Porter, K., Thach, K., Sack, R. J., & Hakansson, N. A. (2017). Building a better gluteal bridge: Electromyographic analysis of hip muscle activity during modified single-leg bridges. *International Journal of Sports Physical Therapy*, *12*(4), 543–549.

Lirio-Romero, C., Palomo-Carrión, R., Romay-Barrero, H., Ferri-Morales, A., Prieto-Gómez, V., & Torres-Lacomba, M. (2021). Age differences in motor recruitment patterns of the shoulder in dynamic and isometric contractions. A cross-sectional study. *Journal of Clinical Medicine*, *10*(3). https://doi.org/10.3390/jcm10030525

Ryan, N., & Bruno, P. (2024). The clinical utility of the prone hip extension test in the diagnosis of motor control impairments associated with low back pain: A cross-sectional study using motion capture and electromyography. *Clinical Biomechanics*, *118*, 106317. https://doi.org/10.1016/j.clinbiomech.2024.106317

Schuermans, J., Van Tiggelen, D., & Witvrouw, E. (2017). Prone hip extension muscle recruitment is associated with hamstring injury risk in amateur soccer. *International Journal of Sports Medicine*, *38*, 696–706. https://doi.org/10.1055/s-0043-103016

Suehiro, T., Mizutani, M., Okamoto, M., Ishida, H., Kobara, K., Fujita, D., Osaka, H., Takahashi, H., & Watanabe, S. (2014). Influence of hip joint position on muscle activity during prone hip extension with knee flexion. *Journal of Physical Therapy Science*, *26*(12), 1895–1898. https://doi.org/10.1589/jpts.26.1895

Youdas, J. W., Hartman, J. P., Murphy, B. A., Rundle, A. M., Ugorowski, J. M., & Hollman, J. H. (2017). Electromyographic analysis of gluteus maximus and hamstring activity during the supine resisted hip extension exercise versus supine unilateral bridge to neutral. *Physiotherapy Theory and Practice*, *33*(2), 124–130. https://doi.org/10.1080/09593985.2016.1271848



Figure 1: Muscle onset time (s) by participant, muscle, and side

Table 1: Fixed effect estimates, standard errors (SE), t statistic, and 95% confidence intervals (CI), rounded to three significant digits

Effect	Estimate	SE	t	95% CI Lower	95% CI Upper
Intercept	0.0734	0.0288	2.55	0.0169	0.13
ES	0.00285	0.0143	0.199	−0.0251	0.0308
BF	0.101	0.0143	7.08	0.0731	0.129
RF	0.296	0.0143	20.7	0.268	0.324
Left	0.0131	0.0143	0.917	−0.0149	0.0411
ES:Left	0.00235	0.0202	0.116	−0.0372	0.0419
BF:Left	−0.146	0.0202	−7.24	−0.186	−0.107
RF:Left	−0.0478	0.0202	−2.37	−0.0874	−0.00824

Table 2: Random effects variance and standard deviation (SD), rounded to three significant digits

Effect	Variance	SD
Participant:Rep	0.00556	0.0746
Participant	0.00338	0.0582
Residual	0.0162	0.127