

Title	Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two o-hydroxyaldehyde groups - The importance of topology
Authors	Pareras, Gerard;Szczepanik, Dariusz Wojciech;Duran, Miquel;Solà, Miquel;Simon, Silvia
Publication date	2019-11-06
Original Citation	Pareras, G., Szczepanik, D. W., Duran, M., Solà, M. and Simon, S. (2019) 'Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two o-hydroxyaldehyde groups - The importance of topology', Journal of Organic Chemistry, 84(23), pp. 15538-15548. doi: 10.1021/acs.joc.9b02526
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1021/acs.joc.9b02526
Rights	© 2019, American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Journal of Organic Chemistry, copyright © American Chemical Society, after peer review and technical editing by the publisher. To access the final edited and published work see https://pubs.acs.org/doi/abs/10.1021/acs.joc.9b02526
Download date	2024-12-05 23:37:06
Item downloaded from	https://hdl.handle.net/10468/9436



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Article

Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two *o*-hydroxyaldehyde groups. The importance of topology

Gerard Pareras, Dariusz Wojciech Szczepanik, Miquel Duran, Miquel Solà, and Silvia Simon

J. Org. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.9b02526 • Publication Date (Web): 06 Nov 2019

Downloaded from pubs.acs.org on November 14, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two *o*-hydroxyaldehyde groups. The importance of topology

Gerard Pareras^{a,‡}, Dariusz W. Szczepanik^{b,‡}, Miquel Duran^b, Miquel Solà^{b,*} and Sílvia Simon^{b,*}


^a Department of Chemistry, University College Cork, College Rd, Cork, Ireland

^b Institut de Química Computacional i Catàlisi and Departament de Química, Universitat de Girona, C/ Maria Aurèlia Capmany 69, 17003 Girona, Catalonia, Spain

‡ These two authors contributed equally to this work.

Corresponding authors: silvia.simon@udg.edu, miquel.sola@udg.edu

TOC Graphic




Abstract

The fact that intramolecular resonance-assisted hydrogen bonds (RAHBs) are stronger than conventional ones is attributed to the partial delocalization of the π -electrons within the hydrogen bond (HB) motif, the so-called quasi-ring. If an aromatic ring is involved in the formation of the RAHB, previous studies have shown that there is an interplay between aromaticity and HB strength. Moreover, in 1,3-dihydroxyaryl-2-aldehydes, some of us found that the position of the quasi-ring formed by the substituents interacting through RAHB influences the strength of the H-bonding, the HBs being stronger when a kinked-like structure is generated by formation of the *quasi*-ring. In this work, we explore this concept further by considering a set of acenes and phenacenes of different sizes with two *o*-hydroxyaldehyde substituents. Calculations with the CAM-B3LYP/6-311+G(d,p)+GD3B method show that for long acenes or phenacenes, once the substituent effect loses importance because quasi-rings are pull apart far from each other, the different topologies rule the HB distances. This fact can be explained in most cases using an extended Clar's aromatic π -sextet model. In some kinked systems, however, the justification from the Clar model has to be complemented by taking into account the repulsion between hydrogen atoms. Triphenylene-like compounds with different number of benzene rings have been studied finding out a very good relationship between aromaticity of the *ipso*- and *quasi*-rings with the RAHB distances. This result confirms the importance of the communication of the π -systems of the *ipso*- and *quasi*-rings.

Introduction

Hydrogen bonds (HBs) are one of the strongest non-covalent interactions. They play an important role in many chemical and biological processes; for instance, HBs determine the crystal structure as well as the stabilization of the second-order structure of proteins.¹⁻⁴ Because of their directionality and the fact of being strong enough to assemble molecular systems in aggregates, HBs are one of the most important interactions in supramolecular chemistry and molecular engineering.⁵ Depending on its bond strength, which can range from 0.2 kcal/mol to 40 kcal/mol, HBs can be classified as weak, moderate or strong.²

HBs assisted by additional effects, e.g., resonance-assisted or charge-assisted,⁶⁻⁹ are among the strongest HBs known. Intramolecular resonance-assisted hydrogen bonds (RAHB) introduced by Gilli *et al.*⁶⁻⁹ are stronger than conventional HBs because of the extra stabilization connected with the partial delocalization of the π -electrons within the so-called *quasi*-ring containing conjugated formally single and double bonds. A schematic representation of electronic effects proceeding within such cyclic RAHB is shown in Scheme 1 for malonaldehyde. The characteristic elongation of formally double bond and shortening of formally single bonds in the *quasi*-ring together with a strengthening of the HB are fingerprints of RAHBs.




Scheme 1. The two main resonance structures involved in the resonance-assisted hydrogen bond in malonaldehyde.

Many computational studies¹⁰⁻²⁷ and experimental works²⁸⁻³¹ can be found related to intramolecular and intermolecular RAHBs. The origin of the additional strength of RAHB as compared to conventional HBs is still subject of an ongoing debate.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


Gilli *et al.*⁶⁻⁹ attributed it to the delocalization of the π -electrons. Mo *et al.*^{16,18} provided further evidences that the enhanced HB comes from the charge flow from the HB donor to the acceptor through the π -conjugation. Recently, Grosch *et al.*¹¹ arrived to the conclusion that π polarization and σ charge transfer are the responsible for enhancing RAHB in line with the Gilli *et al.* proposal, although they emphasized that there is no resonance assistance in the sense of an interplay between σ charge transfer and π polarization. Moreover, other researchers, like Sanz *et al.*,^{25,26} argued that the characteristics of the σ -skeleton, and not the resonance assistance phenomenon, is the main responsible for the extra stability of RAHB. Also, Guevara-Vela *et al.*^{13,14,24} analyzed the nature of the RAHB to conclude that there is an increase in electron localization of π -electrons (not delocalization as expected from Gilli's interpretation) in the *quasi*-ring of RAHB. This localization leads to greater electrostatic, polarization, and charge transfer effects that result in stronger H-bonds. Thus, we can find different interpretations of the basis of the "assistance" in RAHBs.



Scheme 2. The two main resonance structures involved in the resonance-assisted hydrogen bond in *o*-hydroxybenzaldehyde.

The first relation between HB formation and aromaticity was reported in a landmark paper by Dewar already in 1945.³² In his study, Dewar³² showed that presence of an intramolecular hydrogen bond induces aromatic character at the seven-membered ring (7-MR) of the stipatic acid, thus providing an explanation to the physicochemical properties of this molecule. More recently, studies by our group^{33,34} described the interplay between RAHB strength and aromaticity of the *ipso*-ring in a series of *o*-hydroxyarylaldehydes (see Scheme 2). These compounds can separate the π -resonance effect in three different individual


1
2
3 effects, namely, (a) the delocalization within the *quasi*-ring, identified by Gilli *et al.*⁸
4 and already present in malonaldehyde, (b) the effect of the π -electron delocalized
5 within the benzene ring (structure **I** in Scheme 2), and (c) the substituent effect of
6 hydroxyl group (electron-donating group) and carbonyl (electron-withdrawing) that
7 favors the canonical structure with charge separation (structure **II** in Scheme 2).
8 Compared to malonaldehyde, with C=C localized double bond, effects (a) and (c)
9 cooperate with each other, so reinforcing the HB, while (b) goes against HB
10 formation. From the data found in these previous studies, it is clear that RAHB
11 enhancement or weakening can be controlled by changing the substituents³⁵ and
12 the aromaticity of *ipso*-ring.^{33,34} Indeed, in 1,3-dihydroxyaryl-2-aldehydes,³³ we
13 found that the position of the *quasi*-ring formed by the substituents interacting
14 through RAHB influences the strength of the H-bonding, the HBs being stronger
15 when a kinked-like structure is generated by formation of the *quasi*-ring. Similar
16 results were obtained by Houjou *et al.*^{28,29} from the analysis of the proton transfer
17 processes in the two systems represented in Scheme 3. These authors showed
18 that HBs are stronger for **I** than **II** irrespective of solvent polarity or crystal packing.
19 According to the authors, the reason is the low distortion of the aromatic rings in **I**
20 as compared to **II**. For the latter, the loss of aromaticity is higher reducing the
21 stability of the HBs formed.
22
23
24
25
26
27
28
29
30
31
32
33
34
35



47 **Scheme 3.** Structure of the α,α -diimine (**I**) and β,β -diimine (**II**) species.
48
49

50 It is very well-known that aromaticity of condensed benzene rings, the so-called
51 benzenoid hydrocarbons, depends on their number of rings as well as their relative
52 position, that is, their isomeric structure. $[n]$ Acenes and $[n]$ phenacenes (n being
53 the number of rings) are two series of isomeric aligned benzenoid compounds
54
55
56
57
58
59
60

(Scheme 4). The aromaticity of these compounds has been widely analyzed using different indices.³⁶ This aromaticity can be explained using Clar's π -sextet model,³⁷ which states the Kekulé resonance structure with the largest number of disjoint aromatic π -sextets is the most important. Following this rule, the Clar's structures for [3]acene (anthracene) and [3]phenacene (phenanthrene) are the ones represented in Scheme 4. [n]Acenes with a migrating π -sextet have similar aromaticity in all rings,³⁸ whereas in [n]phenacenes the outer rings are more aromatic than the central one. Moreover, [n]phenacenes have a larger number of aromatic π -sextets than the corresponding [n]acenes and, therefore, are more stable and more aromatic as a result of more stabilizing π -interactions.³⁹



Scheme 4. Clar structures for (a) anthracene and (b) phenanthrene.

Following this line of reasoning, one could consider system **II** in Scheme 3 as a part of a *pseudo*-tetracene species formed by a naphthalene attached to two *quasi*-rings in a linear arrangement, whereas compound **I** as a part of *pseudo*-chrysene with two *quasi*-rings in a kinked topology. Given that [n]phenacenes are more stable than [n]acenes, one can make the hypothesis that **I** is more stable than **II** due to the better π interactions found in kinked structures as compared to linear ones. The main goal of this work is to prove this hypothesis by analyzing a set of acenes and phenacenes of different sizes with two *o*-hydroxyaldehyde substituents at the outermost rings. We anticipate here that the arrangement of the aromatic rings has a great influence in the RAHB strength and that kinked distribution of rings results in stronger RAHBs.

Results and Discussion

In this section, we will discuss first the changes in RAHB distances when increasing the chain of polycyclic aromatic hydrocarbons (PAHs) in two different topologies, linear and kinked structures. Then we will analyze the effect of including sequentially a new kink starting from a linear topology to finally yield the fully kinked structure. Finally, the design of triphenylene-like compounds will be used to discuss the direct relation between aromaticity and HB strength.

By adding an extra *quasi*-ring to *o*-hydroxybenzaldehyde (Scheme 2) formed by two adjacent OH and HCO substituents, one can generate the five different isomers of dihydroxydibenzaldehyde shown in Figure 1. Two of them present a linear arrangement and resemble anthracene, whereas three of the isomers have a kinked structure and bear a resemblance to phenanthrene.





Figure 1. Dihydroxydibenzaldehyde isomers with two *quasi*-rings. RAHB distances in Å.

When the new substituents to build the *quasi*-ring (OH and HCO) are added in positions 4 and 5 of *o*-hydroxybenzaldehyde (see Figure 1 for numbering), one can have two linear conformers, **LM-1** and **LP-1** (L for linear and M and P respect to the relative position of both OH and HCO in the *ipso*-ring, *meta* and *para*

respectively). In a previous work,³⁵ we dealt with the substituent effect on RAHB distances for substituted *o*-hydroxybenzaldehyde. It was found that depending on the electronic character of the substituents the RAHB is strengthened or weakened. As can be seen in Scheme 5, EDG and EWG in *para* an *ortho* position to each other helps to activate the RAHB. For that reason **LM-1** presents a shorter RAHB distance (1.736 Å) as compared to *o*-hydroxybenzaldehyde (1.758 Å), whereas the RAHB in **LP-1** is deactivated (1.790 Å), as an EWG (CHO) is added to position 4 and an EDG (OH) is attached to position 5. The difference of 0.054 Å in the RAHB bond length of **LM-1** and **LP-1** reflects the substituent effect. As a result of stronger RAHBs, **LM-1** is more stable than its **LP-1** isomer by 8.15 kcal mol⁻¹ (see Table 1).



Scheme 5. The resonance structures responsible for the HB strengthening. EWG in positions 3 and 5 and EDG in positions 4 and 6 activate the RAHB (see Figure 1 for numbering). For further details see ref. 35.


When the second *quasi*-ring is added at the *ortho* positions, we can get three different kinked conformers, named **KP-1a**, **KP-1b**, and **KM-1** (**KP-1a** for OH and **KP-1b** for HCO in *para* positions). As shown in Figure 1, RAHBs in **KP-1a** and **KM-1** become shortened as compared to the *o*-hydroxybenzaldehyde reference, whereas **KP-1b** remains almost unaffected. ~~**KP-1a** is activated because of the presence of an EWG in position 6 (Scheme 5a) and an EDG in position 5 (Scheme 5f) and also because of H...H repulsion between *ortho* substituents.~~ **KM-1** is activated because of the presence of an EWG in position 3 (Scheme 5b) and an

1
2
3 EDG in position 4 (Scheme 5d). **KP-1a** and **KP-1b** are not activated by
4 substituents when forming the RAHB, as EDG and EWG of the *quasi*-rings are,
5 respectively, in positions 5 and 6 in **KP-1a** and 3 and 4 in **KP-1b**. The slight
6 shortening in **KP-1a** can be explained because of a H...H repulsion interaction.
7 Indeed, when one of the HCO groups in **KP-1a** is rotated by 90° through the C_{ring}-
8 C_{co} bond to place it perpendicular to the 6-MR in such a way that both the RAHB
9 and the H...H repulsion interaction are broken, the RAHB that remains increases
10 its bond length by 0.048 Å. Moreover, we attributed the small deactivating effect in
11 **KP-1b** to the O...O lone pair repulsions between the two OH groups. Table 2
12 shows that **KM-1** is much more stable than **KP-1a** and **KP-1b** (10.86 kcal mol⁻¹
13 and 9.09 kcal mol⁻¹ respectively). The local dipole-dipole repulsions together with
14 H...H and O...O repulsions are the main causes for the destabilization of **KP-1a**
15 and **KP-1b** with respect to **KM-1**. It is worth mentioning that a weak HB (2.423 Å)
16 between aldehyde group and oxygen of the hydroxyl group may also slightly favor
17 the stabilization of the **KM-1** isomer.
18
19
20
21
22
23
24
25
26
27
28
29

30
31 If we now focus on aromaticities of the *ipso*-rings, which are collected in Table 1
32 and Table 2 for linear and kinked compounds respectively, we can state that there
33 is not a direct relation between the aromaticity of the *ipso*-ring and the HB
34 distances, although the systems with stronger RAHBs are the ones having the less
35 aromatic *ipso*-rings. Thus, from these data one can observe that different effects
36 are the responsible of the RAHB length in the *quasi*-ring, being not always very
37 easy to be predicted the main factor ruling the change in the HB distance. To
38 decrease substituent effects, one can increase the distance between them by
39 adding extra benzene rings and building PAHs with two o-hydroxyaldehyde
40 substituents with two different topologies, namely, linear (named **LM-n** and **LP-n**,
41 where **n** is the number of benzene rings) and kinked (**KM-na**, **KM-nb**, and **KP-n**
42 following the nomenclature of Figure 1 and being **n** the number of benzene rings).
43 All these new compounds resemble [*n*]acenes (linear) and [*n*]phenanthrenes
44 (kinked). It is worth mentioning that the maximum number of added benzene rings
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

in this work is 5, as it was found that for $[n]$ acenes with larger n values, open-shell biradical singlet state can be more stable than closed-shell singlet state.^{40,41}

Figure 2 depicts the linear structures studied with the corresponding RAHB distances and the labels of the 6-MRs. Aromaticity of the different 6-MRs and *quasi*-rings are collected in Table 1. Let's start adding one ring to both **LP-1** and **LM-1** systems, so giving **LP-2** and **LM-2**. The addition of a first ring brings to a reduction of the strength of RAHBs that have longer HB distances. There are two electronic effects that explain the weakening of RAHBs in **LP-2** and **LM-2**. First, the substituent effect decreases as the number of benzene rings increases. This is especially important in the case of **LP-2** for which it is not possible to find monoionic resonance structures that favor the RAHB. For **LM-2**, the monoionic resonance structure in Scheme 6 (and its symmetric counterpart not depicted) indicates that substituent effects favor both RAHBs as shown already by Krygowski *et al.*,⁴² but the effect is not as important as in **LM-1** due to the increased distance between substituents.



Scheme 6. Substituted naphthalene resonance structure responsible for the HB strengthening in **LM-2**.

Second, the aromaticity of the 6-MRs when going from **LP-1** or **LM-1** to **LP-2** or **LM-2** compounds decreases and the π -density is partially localized in the C2–C3, C4–C5, and C6–C1 bonds (see Figure 1 for labels). Then, the C1–C2 bond having the two substituents has less π -electrons available for resonance assistance (C1–C2 bond distances are 1.425 Å for **LM-1** and 1.435 Å for **LM-2**) and, consequently, the HB is weakened. The same trend is also followed when adding more benzene rings, showing a convergence of hydrogen bond distances and aromaticities for $n > 3$. Linear PAHs present a migrating π -sextet (see Scheme 4), with similar

aromaticities for the different rings. The convergence between **LP** and **LM** aromaticities can be seen in Table 1. **LM-1** and **LP-1** start with very different aromaticities (0.0563 and 0.0817 respectively) and they converge to similar values (0.0435 and 0.0442 for *ipso*-rings in **LM-5** and **LP-5** respectively).

Table 1. PDI values (in e) for the benzene rings and *quasi*-rings (PDI_q) of the linear systems. ΔE_{rel} is the relative energy between **LM_n** and **LP_n** isomers (kcal mol⁻¹). See Figure 2 for labels of the rings.

Systems	PDI			PDI _q		ΔE_{rel}
	A	B	C	A'	B'	
LM-1	0.0563			0.0107	0.0107	-
LP-1	0.0817			0.0134	0.0134	8.15
LM-2	0.0553	0.0553		0.0080	0.0080	-
LP-2	0.0649	0.0649		0.0086	0.0086	3.20
LM-3	0.0503	0.0616	0.0503	0.0064	0.0064	-
LP-3	0.0540	0.0670	0.0540	0.0066	0.0066	1.37
LM-4	0.0462	0.0599	0.0599	0.0055	0.0055	-
LP-4	0.0477	0.0619	0.0619	0.0056	0.0056	0.62
LM-5	0.0435	0.0577	0.0631	0.0050	0.0050	-
LP-5	0.0442	0.0588	0.0648	0.0050	0.0050	0.28

Table 1 also collects the relative energy between **LM** and **LP** isomers, with **LM** being always the most stable one. When only one benzene ring is separating both *quasi*-rings (**LP-1** and **LM-1**) the energy difference is quite large (8.15 kcal mol⁻¹). When the number of 6-MR is increasing, the relative stability of **LP-n** with respect to **LM-n** decreases to 0.28 kcal mol⁻¹, the two isomers being almost degenerated for $n = 5$. Not only relative energies and aromaticities converge with the increase of n , but also HB lengths as shown in Figure 2. Then, despite the substituent effect favors the RAHB in **LM-n** systems and disfavors it in **LP-n** species, there is a convergence in the relative energies, aromaticities, and HB lengths of this two series of compounds, indicating that substituent effects becomes negligible for systems with $n > 3$.




Figure 2. $[n]$ Acene-like isomers ($n = 1-5$). Bond lengths in Å.

Now let's focus on kinked systems, which are built from **KP-1a**, **KP-1b**, and **KM-1**, by adding extra 6-MRs to generate the different $[n]$ phenacene-like compounds depicted in Figure 3. **KM** has been used to name systems with the same substituent building the *quasi*-rings located in “*meta*” position one with respect the other, i.e., not connected by a straight axis containing the two most separated C atoms. Similar to the linear set, we build kinked PAHs up to five benzene rings. The main difference with respect to linear conformers is that, for each system, we can have two different *quasi*-rings with RAHB distance. When one ring is added, the hydrogen bond distances increase or decrease depending on aromaticity of the *ipso*-ring and the substituent effect. However, again, the substituent effect is partially neutralized due to the existence of different π -delocalization paths. As before, the RAHB distances tend to converge while increasing the number of benzene rings. Relative energies between kinked and linear (**LM-n** as reference) are collected in Table 2, ΔE_{K-L} . As the number of kinked rings increase, they are much more stable than the linear isomers, in agreement with previous studies of

Poater *et al.*³⁸ It is worth noting that the relative energy of the different kinked isomers for $n \geq 2$ depends mainly on the number of H...H repulsion interactions (2HRIs) between the H atoms located in the bay regions of the kinked species. The higher the number of 2HRI, the lower the stability of the isomer. For instance, **KM-2b** (two 2HRI) is less stable than **KP-2** (one 2HRI) and this in turn is less stable than **KM-2a** (no 2HRI). Or the order of stability for $n = 3$ is **KP-3a** (three 3HRI) < **KM-3** (two 2HRI) < **KP-3b** (one 2HRI). And the same for $n = 4$ and 5 (see Figure 4).

Table 2. PDI values (in e) for the benzene rings and *quasi*-rings (PDI_Q) of the kinked systems. $E_{(K-L)}$ is the relative energy to LM- n while E_{rel} is the relative energy between **KM- n** and **KP- n** isomers (kcal mol⁻¹)

Systems	PDI					PDI _Q		ΔE_{K-L}^a	ΔE_{rel}^b
	A	B	C	D	E	A'	B'		
KM-1	0.0586					0.0149	0.0138	0.00	
KP-1a	0.0668					0.0202	0.0202	10.86	10.86
KP-1b	0.0704					0.0170	0.0170	9.09	9.09
KM-2a	0.0609	0.0609				0.0192	0.0192	-3.56	
KM-2b	0.0592	0.0592				0.0195	0.0195	2.75	6.30
KP-2	0.0539	0.0575				0.0179	0.0178	-3.13	0.42
KM-3	0.0611	0.0509	0.0599			0.0179	0.0179	-9.15	2.21
KP-3a	0.0608	0.0540	0.0608			0.0187	0.0187	-5.00	6.36
KP-3b	0.0632	0.0519	0.0632			0.0182	0.0182	-11.36	
KM-4a	0.0613	0.0551	0.0551	0.0613		0.0840	0.0184	-19.39	
KM-4b	0.0600	0.0557	0.0557	0.0600		0.0186	0.0186	-13.02	6.37
KP-4	0.0591	0.0563	0.0542	0.0609		0.0184	0.0182	-16.53	2.86
KM-5	0.0610	0.0540	0.0585	0.0546	0.0597	0.0182	0.0184	-24.79	3.00
KP-5a	0.0599	0.0553	0.0589	0.0553	0.0599	0.0185	0.0185	-21.47	6.33
KP-5b	0.0614	0.0535	0.0600	0.0535	0.0614	0.0183	0.0183	-27.79	

^a LM- n is the reference system for energy difference

^b The most stable systems between **KM- n** and **KP- n** is taken as the reference system




Figure 3. [*n*]Phenacene-like isomers (*n* = 1–5). Bond lengths in Å.




Figure 4. The four H...H repulsion interactions in **KM-5** molecule indicated in red color. Bond lengths in Å.

Previous studies³⁶ reported that, in [*n*]phenacenes, the aromaticity is larger in the external rings than in the central ones, while in [*n*]acenes external rings are slightly

1
2
3 aromatic than inner ones. For our systems, this fact should be translated in more
4 aromatic *quasi*-rings for **KP** and **KM** compounds as compared to **LP** and **LM**
5 systems. Indeed, this is what is found and, for instance, PDI_Q is 0.005 e in **LM-5**
6 and four time higher (0.0182 e) in **KM-5**. This is in agreement with the HB strength
7 of *quasi*-rings, being stronger, with shorter RAHB distance, in kinked compounds
8 (with larger aromaticity for the *quasi*-ring) than in their linear counterparts. In
9 addition, kinked PAHs have an alternation of the aromaticity from the external to
10 the central rings. This alternation can easily be predicted by Clar's aromatic π -
11 sextet model (Scheme 4).
12
13
14
15
16
17
18
19

20
21 When adding one more ring to **KP-1a** to build **KP-2**, we can see that *quasi*-rings
22 in **KP-2** have quite different HB distances (1.650 Å and 1.702 Å for **A'** and **B'**
23 respectively) but similar aromaticities (0.0179 and 0.0178 e). This change on the
24 HB distances is neither related to the aromaticity of the *ipso*-ring (0.0539 e for **A**
25 and 0.0575 e for **B**) nor to the double bond character of the shared C1–C2 (1.403
26 Å for **A'** and 1.395 Å for **B'** ring). The difference in these distances can be
27 understood if we take into account that *quasi*-ring **A'** is affected by a 2HRI that
28 “artificially” makes the RAHB shorter, whereas *quasi*-ring **B'** is free of this type of
29 interactions. The same behavior is found for the rest of isomers (e.g. 1.713 Å for
30 **KM-2a** with no 2HRI and 1.669 Å for **KM-2b** with two 2HRI). To validate the
31 decrease of the HB distance due to the presence of 2HRI, we have removed one
32 of the hydrogen atoms involved in the repulsion to generate a radical structure
33 without 2HRI. In the case of radical **KM-5** generated after removal of a single
34 hydrogen (the one close to the CHO group), the HB length increases from 1.662
35 Å to 1.719 Å, which is very close to the HB length in the second *quasi*-ring (1.715
36 Å) i.e. the one without H...H repulsion. So, indeed, the repulsion seems to be
37 responsible for shortening the HB distance for about 0.06 Å. Moreover, in the case
38 of **KP-5a**, removal of the repulsion between two H on both sites (the ones on
39 phenyl rings close to the *quasi*-ring) leads to the increase of HB length from 1.663
40 Å in **KP-5a** (with H...H repulsion) to 1.720 Å in the biradical system with no H...H
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 repulsion affecting the RAHB. Finally, let us note that at variance with the linear
4 species, the RAHB distances converge already for $n = 2$.
5
6
7

8 From our results, we can see that RAHB distance can be tuned by adding a set of
9 6-MRs between *quasi*-rings. When a linear chain is built, the RAHB distance
10 increases, whereas for kinked chains the length of RAHB remains quite constant.
11 Clar's aromatic π -sextet model extended to *quasi*-rings helps to explain the
12 different RAHB length and aromaticity of the *ipso* and *quasi*-rings in linear and
13 kinked compounds. Nevertheless, the two different RAHB lengths found in some
14 kinked systems and the fact that the most stable isomers have, in some cases,
15 larger RAHB distances cannot be explained with the Clar model. To explain these
16 results one needs to resort to H...H repulsive interactions.
17
18
19
20
21
22
23
24
25

26 **On the way from linear to kinked structures**

27
28
29

30 As said in the previous section, the general trend of the bond lengths in RAHBs of
31 kinked and linear PAHs can be explained using the extended Clar's model that
32 relates aromaticity of the *ipso*- and *quasi*-rings with the RAHB distance. To get
33 deeper insight into the different behavior of linear **LM-5** and kinked **KP-5a** species,
34 we transformed **LM-5** compound to **KP-5a** by kinking one by one the different
35 rings, so going through **LM-5'** (one kink), **KP-5'** (two kinks) and finally **KP-5a** (three
36 kinks). Figure 5 schematically draws the four different compounds that brings to
37 the kinked **KP-5a** from linear **LP-5**. Starting from one kink in right side (**LM-5'**),
38 there is an important decrease of the HB distance (from 1.800 Å to 1.665 Å), while
39 the HB distance of the *quasi*-ring that keeps in a linear topology remains almost
40 unchanged. As it can be seen in Table 3, this change in the topology goes together
41 with important change in the aromaticity of the different rings. Following
42 expectations from Clar's model, there is a decrease of aromaticity in the **D** ring
43 (phenanthrene-like topology) in favor of an increase in **E** ring. As pointed out
44 before, the kinked *quasi*-ring (**B'**) with shorter HB distance becomes much more
45 aromatic.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. PDI values (in e) of the benzene rings and *quasi*-rings (PDI_Q) of the systems studied along the change from linear **LM-5** to kinked **KP-5a** structure. Relative energies referred to **LM-5** (kcal mol⁻¹)

System	PDI					PDI _Q		E _{rel}
	A	B	C	D	E	A'	B'	
LM-5	0.0435	0.0577	0.0631	0.0577	0.0435	0.0050	0.0050	0.00
LM-5'	0.0495	0.0634	0.0613	0.0309	0.0665	0.0060	0.0170	-11.42
KP-5'	0.0654	0.0387	0.0731	0.0401	0.0638	0.0172	0.0176	-23.44
KP-5a	0.0599	0.0553	0.0589	0.0553	0.0599	0.0185	0.0185	-21.47




Figure 5. From linear to kinked PAHs. Bond lengths in Å.


Now we can proceed kinking the other side of the chain, obtaining the **KP-5'** compound. We observed a decrease of the HB distance to 1.719 Å, but not as large as in the previous kink (1.665 Å). The difference again the presence of a 2HRI in the first kink and not in the second one. As seen in the previous kink, there is an important lost of aromaticity in ring **B** and an increase in rings **A** and **A'**. To get the total kinked compound **KP-5a**, we proceed to twist the central part of the chain. Aromaticity of the central rings becomes more similar to each other, with larger values for **A**, **C**, and **E**, as it was expected from Clar's model. *Quasi*-ring aromaticities present no important changes, as their kinked topology do not change when kinking the central part of the chain. The C1–C2 bonds bearing the

substituents in the two outer aromatic rings changes from 1.451 Å in **LM-5** to 1.399 Å in **KP-5a**, thus indicating an increase in the double character of this bond when going from linear to kinked structures. As before, the aromaticity of the *ipso*- and *quasi*-rings increases when going from the linear to the kinked topology.

Finally, Table 3 also collects the relative energies of the different conformers with respect to the linear one (**LM-5**). As it was expected, when kinking only one site, there is an important stabilization (-11.42 kcal mol⁻¹). This agrees with previous work showing that phenanthrene is more stable than anthracene because of better π -interactions.³⁹ When the other site of the chain is kinked, the stabilization energy increases up to -23.44 kcal mol⁻¹, so reinforcing the stabilization because of phenanthrene-like topology being more stable. In the last conformer, where all the chain is kinked, the relative energy decreases to -21.47 kcal mol⁻¹ but because there is an increase of hydrogen repulsion through the whole chain, five 2HRI in **KP-5a** to be compared to three 2HRI for **KP-5**.

Triphenylene-like RAHB

Let us move one step further and consider a set of compounds with two *quasi*-rings that resemble triphenylene (see Scheme 7). Starting from **KP-1a**, **KP-1b**, and **KM-1**, one can add successive benzene rings in the *ipso*-ring keeping the original two *quasi*-rings to generate the compounds depicted in Figure 6 named **KP-Lna**, **KP-Lnb**, and **KM-Ln**, with *n* being the number of benzene rings.



Scheme 7. (a) Clar's structure for triphenylene (b) triphenylene-like double-headed *quasi*-ring compound




Figure 6. Triphenylene-like studied structures. Bond lengths in Å.

For systems with triphenylene-like structures, substituent effects on the RAHB of the *quasi*-rings remain the same in each series of compounds (for each column in Figure 6). After adding one ring (**KP-L2a**, **KP-L2b**, and **KM-L2**) an important shortening of the RAHB distances are observed. According to Clar's π -sextet model (Scheme 6), the central ring in triphenylene is an "empty ring" while the

outer rings are the most aromatic ones. Analyzing data collected in Table 4, one can observe the *ipso*-rings have much smaller aromaticities than the added benzene ring (e.g. for **KP-L2a** 0.0344 e for ring **A** and 0.0860 e for ring **B**). For these compounds it is expected that aromaticity of all external ring being large, so *quasi*-ring also increase their aromaticity when adding a benzene ring (from 0.0170 e for **KP-1b** to 0.0217 e for **KP-L2b**).

Table 4. PDI values (in e) of benzene rings and *quasi*-rings (PDI_Q) of the different triphenylene-like systems. Relative energies respect to KM-Ln (kcal mol⁻¹).

Systems	PDI					PDI _Q		E _{rel}
	A	B	C	D	E	A'	B'	
KP-1a	0.0668					0.0202	0.0202	10.86
KP-1b	0.0704					0.0170	0.0170	9.09
KM-1	0.0586					0.0149	0.0138	
KP-L2a	0.0344	0.0860				0.0242	0.0242	6.22
KP-L2b	0.0383	0.0870				0.0217	0.0217	12.31
KM-L2	0.0329	0.0843				0.0197	0.0175	
KP-L3a	0.0243	0.0682	0.0699			0.0259	0.0259	5.68
KP-L3b	0.0287	0.0703	0.0699			0.0230	0.0230	12.49
KM-L3	0.0238	0.0689	0.0691			0.0212	0.0183	
KP-L4a	0.0206	0.0590	0.0660	0.0613		0.0267	0.0267	5.47
KP-L4b	0.0249	0.0617	0.0670	0.0609		0.0237	0.0237	12.61
KM-L4	0.0201	0.0595	0.0667	0.0610		0.0218	0.0191	
KP-L5a	0.0189	0.0552	0.0633	0.0628	0.0564	0.0268	0.0268	5.47
KP-L5b	0.0227	0.0563	0.0658	0.0627	0.0561	0.0239	0.0239	12.72
KM-L5	0.0181	0.0545	0.0635	0.0634	0.0563	0.0221	0.0193	

The aromaticity of the *ipso*-ring can be tuned by adding extra 6-MRs to it, following a linear topology. As it was commented before, for [*n*]acenes with migrating π -sextet, the aromaticity of all rings decreases while increasing the size of the chain. Keeping this idea in mind, results show that the aromaticity of the *ipso*-ring, which has from the very beginning a low aromaticity because it resembles to the central ring of triphenylene, decreases with the increase in the number of 6-MRs attached to the *ipso*-ring. Because of that, the C=C bond in the ring junction between the *ipso*-ring and the *quasi*-ring becomes more localized for larger systems. The increase in the double character of this bond strengthens the RAHB, which becomes very short for **KM-L5** (going from 1.689 Å in **KM-1** to 1.575 Å in **KM-L5**).

1
2
3 As the *quasi*-ring has more available its π -electron pair for larger systems, its
4 aromaticity increases going from, for instance, 0.0149 e for ring **A'** of **KM-1** to
5 0.0221 e for **KM-L5**. Indeed, for **KM-L5**, the *quasi*-ring is more aromatic than the
6 *ipso*-ring with a PDI of 0.0181 e. Another important issue different from the
7 previous linear and kinked systems studied is that convergence is not fully
8 achieved for $n=5$, neither on the RAHB distance nor on the aromaticity of the RAHB
9 *quasi*-ring.

10
11
12
13
14
15
16
17 As it was pointed out before, substituent effect are not changing while enlarging
18 the chain for each series, so we can analyze the relation between aromaticities of
19 the *ipso*-ring and the *quasi*-ring and the HB distances. The linear representation
20 for the different parameters can be found in the SI for each *quasi*-ring. There are
21 four different *quasi*-rings to take into account; one for **KP-Lna**, system with one
22 2HRI, one for **KP-Lnb**, species with two 2HRI, and two for **KM-Ln**, with one 2HRI
23 and one weak HB, all of them with different HB distances. Figure S1 in SI shows
24 a very good linear relationship between the aromaticity of the *quasi*-ring and the
25 *ipso*-ring for the four series of different RAHB (average R^2 of 0.999); when the
26 aromaticity of the *ipso*-ring decreases, that of the *quasi*-ring increases. Previous
27 works^{33,34} reported that an increase of the PDI in the *quasi*-ring reinforces the
28 RAHB, and the HB distance is shortened. With this idea in mind, we expect to find
29 four linear relationships between both parameters, the aromaticity of the *ipso*-ring
30 and the HB length. This correlation is fulfilled for **A'** and **B'** *quasi*-rings of **KP-1a**
31 and *quasi*-ring **B'** of **KM-1**. For the other two *quasi*-rings the expected correlation
32 in not followed (Fig S2 in SI). In this case, the calculated HB distances in **A'** and
33 **B'** *quasi*-rings of **KP-1b** and *quasi*-ring **A'** of **KM-1** are much longer that expected
34 from the correlation of the series. Comparing these two compounds with the rest
35 of the series, it can be observed that when a second 6-MR is added to **KP-1b** and
36 **KM-1** an extra 2HRI between the H atom of the HCO group of the *quasi*-ring and
37 one of the H atoms of the added benzene-ring that decreasing the hydrogen bond
38 length. This repulsion is the same in the rest of the series, and therefore, the linear
39 correlation between aromaticity of the *quasi*-ring, the *ipso*-ring and the RAHB
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 distances is kept, except for the **KP-1b** and **KM-1** that are not affected by any
4
5 2HRI.
6
7

8 The different behavior of **KP-1a** and **KM-1** (ring **B'**) can also be assessed
9 analyzing the relative energy between conformers of the same size (Table 4).
10 These two conformers present a very different stability through (10.86 kcal mol⁻¹
11 for **KP-1a** and 9.09 kcal mol⁻¹ for **KP-1b** both with respect to **KM-1**) when
12 compared to the rest of the series. These relative energies keep constant (approx.
13 5 and 12 kcal mol⁻¹ respectively) through the series **KP-Lna**, **KP-Lnb** and **KM-Ln**
14 when there are more than one acene-like ring, as the number of 2HRI remains the
15 same. The relative stability of the three series (**KM-Ln** being the most stable,
16 followed by **KP-Lna** and **KP-Lnb**) is ruled by the 2HRI between *quasi*-ring and the
17 added benzene ring in **KP-Lnb** and between both *quasi*-rings and the added
18 benzene ring in **KP-Lna**.
19
20
21
22
23
24
25
26
27
28

29 In the series of triphenylene-like compounds studied, the substituent effect remains
30 the same. For this reason, in these systems we could assess the importance of π -
31 delocalization within the *ipso* and *quasi* ring (aromaticity). This delocalization
32 assists the HB, shortening it, so reinforcing the concept of resonance-assisted
33 hydrogen bond.
34
35
36
37
38

39 Conclusions

40 Double-headed compounds having two intramolecular resonance-assisted
41 hydrogen bonds (RAHB) have been designed with the aim of studying the relation
42 between the RAHB length and the aromaticity of the different *ipso*-rings. For these
43 purpose, we have built [*n*]acenes (linear) and [*n*]phenancenes (kinked) like
44 compounds with different number of benzene rings. Two different effects are
45 controlling the HB distances, namely, the connectivity between substituents of the
46 *ipso*-ring and the topology of the benzene chains (being linear or kinked). The first
47 effect has been decreased by adding up to five benzene rings between *quasi*-rings
48 in [*n*]acenes and [*n*]phenancenes, so being the topology of the chain the
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 responsible of ruling the different HB strength. The HB lengths observed can be
4 explained in most cases using an extended Clar's aromatic π -sextet model that
5 incorporates the *quasi*-rings as if they were rings in benzenoids. In general, shorter
6 HBs are found for the more aromatic *quasi*-rings. The existence of H...H bond
7 repulsions in some kinked systems explain the exceptions found to this extended
8 Clar's aromatic π -sextet model.
9
10
11
12
13
14

15 Finally, in order to assess the relation between aromaticity (based on PDI values)
16 and HB length, triphenylene-like compounds have been studied. In these systems,
17 where the connectivity between substituent is the same, we found good linear
18 correlation between aromaticity of the *ipso*- and *quasi*-ring and the RAHB
19 distances, thus reinforcing the Gilli's concept of resonance assisting the hydrogen
20 bond formations in the *quasi*-ring.
21
22
23
24
25
26

27 **Computational methods**

28
29
30 All molecules considered in this study were calculated using the long-range
31 corrected hybrid CAM-B3LYP level of theory⁴³ including the D3 version of
32 Grimme's dispersion with Becke-Johnson damping⁴⁴ to provide a correct
33 description of the distance-dependence of the hydrogen bonding interactions⁴⁵ as
34 well as the inter-ring electron delocalization effects in the condensed benzenoid
35 systems^{38,46,47,38} The standard 6-311++G(d,p) basis set was used for all atoms.
36 Vibrational analyses of optimized structures showed that the structures are minima
37 in the potential energy surface. All calculations were carried out with the Gaussian
38 09 package.⁴⁸ To avoid the numerical-integration accuracy issues⁴⁹ and the basis-
39 set dependence problems,⁵⁰ in this study the delocalization indices (DIs)⁵¹ have
40 been calculated within the representation of natural atomic orbitals (NAO)⁵²
41 available within the NBO6⁵³ and JaNPA⁵⁴ program interfaces.
42
43
44
45
46
47
48
49
50
51

52
53 DIs are among the most popular bonding indicators. They are closely related to
54 the covalent bond-order indices⁵⁵ and provide a quantitative information on the
55
56
57
58
59
60

1
2
3 electron density shared between two atoms/fragments A and B. Within the NAO-
4 based partitioning scheme the delocalization index for one-determinant closed-
5 shell systems is congeneric with the quadratic bond-order originally proposed by
6 Wiberg⁵⁶
7
8
9

$$\delta(A, B) = 2 \sum_{\mu \in A} \sum_{\nu \in B} |P_{\mu\nu}|^2 \quad (1)$$

10
11
12 where $P_{\mu\nu}$ are the elements of the reduced density matrix in the NAO basis.⁵²
13
14
15

16
17
18 Based on the finding of Fulton⁵⁵ and Bader⁵⁷ who showed that benzene has larger
19 DIs in *para*-related atoms than in *meta*-related ones, the PDI⁵⁸ index uses the DIs
20 of *para*-related atoms as a measure of aromaticity for six-membered rings (6-MRs):
21
22
23
24

$$PDI = \frac{\delta(A_1, A_4) + \delta(A_2, A_5) + \delta(A_3, A_6)}{3} \quad (2)$$

25
26
27
28
29 For PDI, the larger the index, the greater the aromaticity of the ring. Obviously, this
30 index can only be applied to 6-MRs. We use it to discuss the aromaticity of both
31 the benzene rings (PDI) and the *quasi*-ring (PDI_Q), also known as *quasi*-
32 aromaticity.⁵⁹ It was demonstrated⁶⁰ for simple derivatives of benzene that PDI
33 gives qualitatively the same results as other electronic indices as well as indices
34 like the HOMA⁶¹ structural index or the NICS magnetic index.⁴⁹ For this reason, in
35 our studies we are using exclusively the PDI index.
36
37
38
39
40
41
42
43
44

45 Supporting Information

46
47 Linear correlation between PDI for the *ipso* and *quasi*-rings (Figure S1) and linear
48 correlation between aromaticity of the *ipso*-ring and RAHB bond length (Figure S1) for the
49 triphenylene-like compound. Optimized Cartesian coordinates for all compounds
50 considered at the CAM-B3LYP/6-311+G(d,p)+GD3B level of theory are given in a
51 separated file.
52
53
54
55
56
57
58
59
60

Acknowledgements

We are grateful for financial support from the Spanish MINECO (CTQ2017-85341-P) and the Catalan DIUE (2017-SGR-39, XRQTC, and ICREA Academia 2014 Award to M.S.) and the FEDER fund (UNGI10-4E-801). D.W.S acknowledges the financial support of the Polish National Science Center within the Sonata Project 2015/17/D/ST4/00558.

References

- (1) Grabowski, S. J. *Hydrogen Bonding - New Insights*; Springer: The Netherlands, 2006; Vol. 3.
- (2) Jeffrey, G. A. *An Introduction to Hydrogen Bonding*; Oxford University Press: USA, 1997.
- (3) *Molecular Interactions: From Van Der Waals to Strongly Bound Complexes*; Scheiner, S., Ed.; Wiley & Sons: UK, 1997.
- (4) Steiner, T. The Hydrogen Bond in the Solid State. *Angew. Chem. Int. Ed.* **2002**, *41*, 48-76.
- (5) Desiraju, G. R.; Steiner, T. *The Weak Hydrogen Bond in Structural Chemistry and Biology*; Oxford University Press: New York, 1999.
- (6) Bertolasi, V.; Gilli, P.; Ferretti, V.; Gilli, G. Evidence for Resonance-Assisted Hydrogen Bonding. 2. Intercorrelation between Crystal Structure and Spectroscopic Parameters in Eight Intramolecularly Hydrogen Bonded 1,3-Diaryl-1,3-Propanedione Enols. *J. Am. Chem. Soc.* **1991**, *113*, 4917-4925.
- (7) Bertolasi, V.; Nanni, L.; Gilli, P.; Ferretti, V.; Gilli, G.; Issa, Y. M.; Sherif, O. E. Intramolecular N-H...O=C Hydrogen-Bonding Assisted by Resonance

1
2
3 - Intercorrelation between Structural and Spectroscopic Data for 6 Beta-Diketo-
4 Arylhydrazones Derived from Benzoylacetone or Acetylacetone. *New J. Chem.*
5 **1994**, *18*, 251-261.
6
7

8
9
10 (8) Gilli, G.; Bellucci, F.; Ferretti, V.; Bertolasi, V. Evidence for
11 Resonance-Assisted Hydrogen Bonding from Crystal-Structure Correlations on
12 the Enol Form of the β -Diketone Fragment. *J. Am. Chem. Soc.* **1989**, *111*, 1023.
13
14

15
16 (9) Gilli, P.; Bertolasi, V.; Pretto, L.; Ferretti, V.; Gilli, G. Covalent Versus
17 Electrostatic Nature of the Strong Hydrogen Bond: Discrimination among Single,
18 Double, and Asymmetric Single-Well Hydrogen Bonds by Variable-Temperature
19 X-Ray Crystallographic Methods in Beta-Diketone Enol RAHB Systems. *J. Am.*
20 *Chem. Soc.* **2004**, *126*, 3845-3855.
21
22
23
24

25
26 (10) Gora, R. W.; Maj, M.; Grabowski, S. J. Resonance-Assisted
27 Hydrogen Bonds Revisited. Resonance Stabilization vs. Charge Delocalization.
28 *Phys. Chem. Chem. Phys.* **2013**, *15*, 2514-2522.
29
30
31

32
33 (11) Grosch, A. A.; van der Lubbe, S. C. C.; Fonseca Guerra, C. Nature
34 of Intramolecular Resonance Assisted Hydrogen Bonding in Malonaldehyde and
35 Its Saturated Analogue. *J. Phys. Chem. A* **2018**, *122*, 1813-1820.
36
37
38

39
40 (12) Guerra, C. F.; Bickelhaupt, F. M.; Snijders, J. G.; Baerends, E. J. The
41 Nature of the Hydrogen Bond in DNA Base Pairs: The Role of Charge Transfer
42 and Resonance Assistance. *Chem. Eur. J.* **1999**, *5*, 3581-3594.
43
44
45

46
47 (13) Guevara-Vela, J. M.; Romero-Montalvo, E.; Costales, A.; Pendas, A.
48 M.; Rocha-Rinza, T. The Nature of Resonance-Assisted Hydrogen Bonds: A
49 Quantum Chemical Topology Perspective. *Phys. Chem. Chem. Phys.* **2016**, *18*,
50 26383-26390.
51
52
53
54
55
56
57
58
59
60

1
2
3 (14) Guevara-Vela, J. M.; Romero-Montalvo, E.; del Río-Lima, A.; Martín
4 Pendás, Á.; Hernández-Rodríguez, M.; Rocha Rinza, T. Hydrogen-Bond
5 Weakening through Π Systems: Resonance-Impaired Hydrogen Bonds (RIHB).
6 *Chem. Eur. J* **2017**, *23*, 16605-16611.
7
8
9

10
11 (15) Guillaumes, L.; Simon, S.; Fonseca Guerra, C. The Role of
12 Aromaticity, Hybridization, Electrostatics, and Covalency in Resonance-Assisted
13 Hydrogen Bonds of Adenine–Thymine (AT) Base Pairs and Their Mimics.
14 *ChemistryOpen* **2015**, *4*, 318-327.
15
16
17
18

19
20 (16) Jiang, X.; Zhang, H.; Wu, W.; Mo, Y. A Critical Check for the Role of
21 Resonance in Intramolecular Hydrogen Bonding. *Chem. Eur. J* **2017**, *23*, 16885-
22 16891.
23
24
25

26
27 (17) Kurczab, R.; Mitoraj, M. P.; Michalak, A.; Ziegler, T. Theoretical
28 Analysis of the Resonance Assisted Hydrogen Bond Based on the Combined
29 Extended Transition State Method and Natural Orbitals for Chemical Valence
30 Scheme. *J. Phys. Chem. A* **2010**, *114*, 8581-8590.
31
32
33
34

35 (18) Lin, X.; Zhang, H.; Jiang, X.; Wu, W.; Mo, Y. The Origin of the Non-
36 Additivity in Resonance-Assisted Hydrogen Bond Systems. *J. Phys. Chem. A*
37 **2017**, *121*, 8535-8541.
38
39
40

41 (19) Mo, Y. Probing the Nature of Hydrogen Bonds in DNA Base Pairs. *J.*
42 *Mol. Model.* **2006**, *12*, 665-672.
43
44
45

46 (20) Mo, Y.; Bao, P.; Gao, J. Energy Decomposition Analysis Based on a
47 Block-Localized Wavefunction and Multistate Density Functional Theory. *Phys.*
48 *Chem. Chem. Phys.* **2011**, *13*, 6760-6775.
49
50
51
52

53 (21) Mohajeri, A. Theoretical Evidences for Resonance-Assisted
54 Hydrogen Bonding. *Journal of Molecular Structure-Theochem* **2004**, *678*, 201-205.
55
56
57
58

1
2
3 (22) Paul, B. K. Lack of Resonance Assistance in a Classical
4 Intramolecular Hydrogen Bond: An Exploration from Quantum Theory of Atoms-in-
5 Molecules Perspective. *J. Phys. Org. Chem.* **2019**, *32*, e3999.
6
7

8
9
10 (23) Paul, B. K.; Guchhait, N. Geometrical Criteria Versus Quantum
11 Chemical Criteria for Assessment of Intramolecular Hydrogen Bond (IMHB)
12 Interaction: A Computational Comparison into the Effect of Chlorine Substitution
13 on IMHB of Salicylic Acid in Its Lowest Energy Ground State Conformer. *Chem.*
14 *Phys.* **2013**, *412*, 58-67.
15
16
17

18
19
20 (24) Romero-Montalvo, E.; Guevara-Vela, J. M.; Costales, A.; Pendás, Á.
21 M.; Rocha-Rinza, T. Cooperative and Anticooperative Effects in Resonance
22 Assisted Hydrogen Bonds in Merged Structures of Malondialdehyde. *Phys. Chem.*
23 *Chem. Phys.* **2017**, *19*, 97-107.
24
25
26

27
28 (25) Sanz, P.; Mó, O.; Yáñez, M.; Elguero, J. Non-Resonance-Assisted
29 Hydrogen Bonding in Hydroxymethylene and Aminomethylene Cyclobutanones
30 and Cyclobutenones and Their Nitrogen Counterparts. *ChemPhysChem* **2007**, *8*,
31 1950-1958.
32
33
34

35
36
37 (26) Sanz, P.; Yáñez, M.; Mó, O. Resonance-Assisted Intramolecular
38 Chalcogen–Chalcogen Interactions? *Chem. Eur. J* **2003**, *9*, 4548-4555.
39
40

41
42 (27) Sobczyk, L.; Grabowski, S. J.; Krygowski, T. M. Interrelation between
43 H-Bond and Pi-Electron Delocalization. *Chem. Rev.* **2005**, *105*, 3513-3560.
44
45

46
47 (28) Houjou, H.; Motoyama, T.; Banno, S.; Yoshikawa, I.; Araki, K.
48 Experimental and Theoretical Studies on Constitutional Isomers of 2,6-
49 Dihydroxynaphthalene Carbaldehydes. Effects of Resonance-Assisted Hydrogen
50 Bonding on the Electronic Absorption Spectra. *J. Org. Chem.* **2009**, *74*, 520-529.
51
52
53
54
55
56
57
58
59
60

1
2
3 (29) Houjou, H.; Shingai, H.; Yagi, K.; Yoshikawa, I.; Araki, K. Mutual
4 Interference between Intramolecular Proton Transfer Sites through the Adjoining
5 Π -Conjugated System in Schiff Bases of Double-Headed, Fused Salicylaldehydes.
6 *J. Org. Chem.* **2013**, *78*, 9021-9031.
7
8
9

10
11 (30) Mahmudov, K. T.; Pombeiro, A. J. L. Resonance-Assisted Hydrogen
12 Bonding as a Driving Force in Synthesis and a Synthone in the Design of Materials.
13 *Chem. Eur. J* **2016**, *22*, 16356-16398.
14
15
16
17

18 (31) Nguyen, Y. H.; Lampkin, B. J.; Venkatesh, A.; Ellern, A.; Rossini, A.
19 J.; VanVeller, B. Open-Resonance-Assisted Hydrogen Bonds and Competing
20 Quasiaromaticity. *J. Org. Chem.* **2018**.
21
22
23

24 (32) Dewar, M. J. S. Structure of Stipitatic Acid. *Nature* **1945**, *155*, 50.
25
26
27

28 (33) Palusiak, M.; Simon, S.; Solà, M. Interplay between Intramolecular
29 Resonance-Assisted Hydrogen Bonding and Aromaticity in *o*-Hydroxyaryl
30 Aldehydes. *J. Org. Chem.* **2006**, *71*, 5241-5248.
31
32
33

34 (34) Palusiak, M.; Simon, S.; Solà, M. Interplay between Intramolecular
35 Resonance-Assisted Hydrogen Bonding and Local Aromaticity. II. 1,3-
36 Dihydroxyaryl-2-Aldehydes. *J. Org. Chem.* **2009**, *74*, 2059-2066.
37
38
39
40

41 (35) Pareras, G.; Palusiak, M.; Duran, M.; Solà, M.; Simon, S. Tuning the
42 Strength of the Resonance-Assisted Hydrogen Bond in *o*-Hydroxybenzaldehyde
43 by Substitution in the Aromatic Ring. *J. Phys. Chem. A* **2018**, *122*, 2279-2287.
44
45
46
47

48 (36) Portella, G.; Poater, J.; Bofill, J. M.; Alemany, P.; Solà, M. Local
49 Aromaticity of [n]Acenes, [n]Phenacenes, and [n]Helicenes (n = 1–9). *J. Org.*
50 *Chem.* **2005**, *70*, 2509-2521.
51
52
53

54 (37) Clar, E. *The Aromatic Sextet*, Wiley: New York, 1972.
55
56
57
58
59
60

1
2
3 (38) Szczepanik, D. W.; Solà, M.; Krygowski, T. M.; Szatyłowicz, H.;
4 Andrzejak, M.; Pawełek, B.; Dominikowska, J.; Kukułka, M.; Dyduch, K.
5 Aromaticity of Acenes: The Model of Migrating Π -Circuits. *Phys. Chem. Chem.*
6 *Phys.* **2018**, *20*, 13430-13436.
7
8
9

10
11 (39) Poater, J.; Visser, R.; Solà, M.; Bickelhaupt, F. M. Polycyclic
12 Benzenoids: Why Kinked Is More Stable Than Straight. *J. Org. Chem.* **2007**, *72*,
13 1134-1142.
14
15
16

17
18 (40) Bendikov, M.; Duong, H. M.; Starkey, K.; Houk, K. N.; Carter, E. A.;
19 Wudl, F. Oligoacenes: Theoretical Prediction of Open-Shell Singlet Diradical
20 Ground States. *J. Am. Chem. Soc.* **2004**, *126*, 7416-7417.
21
22
23

24
25 (41) Yu, D.; Stuyver, T.; Rong, C.; Alonso, M.; Lu, T.; De Proft, F.;
26 Geerlings, P.; Liu, S. Global and Local Aromaticity of Acenes from the Information-
27 Theoretic Approach in Density Functional Reactivity Theory. *Phys. Chem. Chem.*
28 *Phys.* **2019**, *21*, 18195-18210.
29
30
31

32
33 (42) M. Krygowski, T.; Palusiak, M.; Goral, A.; E. Zachara-Horeglad, J.
34 Relationship between Substituent Effect and Aromaticity - Part III: Naphthalene as
35 a Transmitting Moiety for Substituent Effect. *J. Phys. Org. Chem.* **2007**, *20*, 297-
36 306.
37
38
39

40
41 (43) Yanai, T.; Tew, D. P.; Handy, N. C. A New Hybrid Exchange-
42 Correlation Functional Using the Coulomb-Attenuating Method (CAM-B3LYP).
43 *Chem. Phys. Lett.* **2004**, *393*, 51-57.
44
45
46

47
48 (44) Grimme, S.; Ehrlich, S.; Goerigk, L. Effect of the Damping Function
49 in Dispersion Corrected Density Functional Theory. *J. Comput. Chem.* **2011**, *32*,
50 1456-1465.
51
52
53

1
2
3 (45) Thanthiriwatte, K. S.; Hohenstein, E. G.; Burns, L. A.; Sherrill, C. D.
4 Assessment of the Performance of DFT and DFT-D Methods for Describing
5 Distance Dependence of Hydrogen-Bonded Interactions. *J. Chem. Theory*
6 *Comput.* **2011**, *7*, 88-96.
7
8
9

10
11 (46) Casademont-Reig, I.; Woller, T.; Contreras-García, J.; Alonso, M.;
12 Torrent-Sucarrat, M.; Matito, E. New Electron Delocalization Tools to Describe the
13 Aromaticity in Porphyrinoids. *Phys. Chem. Chem. Phys.* **2018**, *20*, 2787-2796.
14
15
16
17

18 (47) Szczepanik, D. W.; Andrzejak, M.; Dominikowska, J.; Pawelek, B.;
19 Krygowski, T. M.; Szatyłowicz, H.; Solà, M. The Electron Density of Delocalized
20 Bonds (Eddb) Applied for Quantifying Aromaticity. *Phys. Chem. Chem. Phys.*
21 **2017**, *19*, 28970-28981.
22
23
24
25

26 (48) Frisch, M. J. T., G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.;
27 Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.;
28 Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.;
29 Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.;
30 Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven,
31 T.; Montgomery, Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.;
32 Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.;
33 Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.;
34 Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo,
35 C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi,
36 R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.;
37 Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas,
38 Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian 09, Revision
39 A.02, Gaussian Inc., Wallingford CT, 2009.
40
41
42
43
44
45
46
47
48
49
50
51

52 (49) Feixas, F.; Matito, E.; Poater, J.; Solà, M. Quantifying Aromaticity
53 with Electron Delocalisation Measures. *Chem. Soc. Rev.* **2015**, *44*, 6434-6451.
54
55
56
57
58
59
60

1
2
3 (50) Jabłoński, M.; Palusiak, M. Basis Set and Method Dependence in
4 Quantum Theory of Atoms in Molecules Calculations for Covalent Bonds. *J. Phys.*
5 *Chem. A* **2010**, *114*, 12498-12505.
6
7

8
9
10 (51) Fradera, X.; Austen, M. A.; Bader, R. F. W. The Lewis Model and
11 Beyond. *J. Phys. Chem. A* **1999**, *103*, 304-314.
12
13

14
15 (52) Reed, A. E.; Weinstock, R. B.; Weinhold, F. Natural Population
16 Analysis. *J. Chem. Phys.* **1985**, *83*, 735-746.
17
18

19
20 (53) E. D. Glendening, J., K. Badenhoop, A. E. Reed, J. E. Carpenter, J.
21 A. Bohmann, C. M. Morales, C. R. Landis, and F. Weinhold. Nbo 6.0; Theoretical
22 Chemistry Institute, University of Wisconsin: Madison, 2013.
23
24

25
26 (54) Nikolaienko, T. Y.; Bulavin, L. A.; Hovorun, D. M. Janpa: An Open
27 Source Cross-Platform Implementation of the Natural Population Analysis on the
28 Java Platform. *Comp. Theor. Chem.* **2014**, *1050*, 15-22.
29
30

31
32
33 (55) Fulton, R. L.; Mixon, S. T. Comparison of Covalent Bond Indexes and
34 Sharing Indexes. *J. Phys. Chem.* **1993**, *97*, 7530-7534.
35
36

37
38 (56) Wiberg, K. B. Application of the Pople-Santry-Segal Cndo Method to
39 the Cyclopropylcarbanyl and Cyclobutyl Cation and to Bicyclobutane. *Tetrahedron*
40 **1968**, *24*, 1083-1096.
41
42

43
44
45 (57) Bader, R. F. W.; Johnson, S.; Tang, T. H.; Popelier, P. L. A. The
46 Electron Pair. *J. Phys. Chem.* **1996**, *100*, 15398-15415.
47
48

49
50 (58) Poater, J.; Fradera, X.; Duran, M.; Solà, M. The Delocalization Index
51 as an Electronic Aromaticity Criterion: Application to a Series of Planar Polycyclic
52 Aromatic Hydrocarbons. *Chem. Eur. J.* **2003**, *9*, 400-406.
53
54
55
56
57
58
59
60

1
2
3 (59) Krygowski, T. M.; Bankiewicz, B.; Czarnocki, Z.; Palusiak, M. Quasi-
4 Aromaticity—What Does It Mean? *Tetrahedron* **2015**, *71*, 4895-4908.
5
6

7
8 (60) Krygowski, T. M.; Ejsmont, K.; Stepień, B. T.; Cyrański, M. K.;
9 Poater, J.; Solà, M. Relation between the Substituent Effect and Aromaticity. *J.*
10 *Org. Chem.* **2004**, *69*, 6634-6640.
11
12

13
14 (61) Krygowski, T. M.; Szatyłowicz, H.; Stasyuk, O. A.; Dominikowska, J.;
15 Palusiak, M. Aromaticity from the Viewpoint of Molecular Geometry: Application to
16 Planar Systems. *Chem. Rev.* **2014**, *114*, 6383-6422.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60