

Title	Relationships between maternal obesity and maternal and neonatal iron status
Authors	Flynn, Angela C.;Begum, Shahina;White, Sara;Dalrymple, Kathryn;Gill, Carolyn;Alwan, Nisreen;Kiely, Mairead E.;Latunde-Dada, Gladys;Bell, Ruth;Briley, Annette;Nelson, Scott;Oteng-Ntim, Eugene;Sandall, Jane;Sanders, Thomas;Whitworth, Melissa;Murray, Deirdre M.;Kenny, Louise C.;Poston, Lucilla;SCOPE UPBEAT Consortiums
Publication date	2018
Original Citation	Flynn, A., Begum, S., White, S., Dalrymple, K., Gill, C., Alwan, N., Kiely, M., Latunde-Dada, G., Bell, R., Briley, A., Nelson, S., Oteng-Ntim, E., Sandall, J., Sanders, T., Whitworth, M., Murray, D., Kenny, L., Poston, L., and On behalf of the SCOPE and UPBEAT Consortiums [2018] 'Relationships between Maternal Obesity and Maternal and Neonatal Iron Status', <i>Nutrients</i> , 10(8), 1000 (10pp). doi: 10.3390/nu10081000
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.mdpi.com/2072-6643/10/8/1000 - 10.3390/nu10081000
Rights	© 2018, the Authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/). - http://creativecommons.org/licenses/by/4.0/
Download date	2023-06-05 13:50:51
Item downloaded from	http://hdl.handle.net/10468/6942



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Table S1. Association between maternal obesity and maternal inflammatory and iron status. Differences and 95% confidence intervals.

Indicator	BMI category	Adjusted difference	95% CI
Ferritin (ug/l)	Normal weight	Reference	
	Obese	0.03	(-0.15, 0.20)
Ferritin <15ug/l*	Normal weight	Reference	
	Obese	1.13	(0.58, 2.19)
Soluble transferrin receptor (nmol/l)	Normal weight	Reference	
	Obese	3.43	(2.50, 4.35)
sTfR:log ₁₀ sF	Normal weight	Reference	
	Obese	0.17	(0.08, 0.26)
Hepcidin (ng/ml)	Normal weight	Reference	
	Obese	0.04	(-1.79, 1.88)
CRP>5mg/l*	Normal weight	Reference	
	Obese	0.20	(0.12, 0.33)
IL-6>1pg/ml*	Normal weight	Reference	
	Obese	0.02	(0.00, 0.07)

Multiple linear regression, adjusted for ethnicity, parity, educational level and smoking. *Multiple logistic regression adjusted for ethnicity, parity, educational level and smoking. sF, ferritin; sTfR, soluble transferrin receptor; CRP, c-reactive protein; IL-6, interleukin 6.

Table S2. Association between maternal obesity and neonatal iron status. Differences and 95% confidence intervals.

Indicator	BMI category	Adjusted difference	95% CI
Ferritin (ug/l)	Normal weight	Reference	
	Obese	-0.12	(-0.26, 0.01)
Ferritin <76ug/l*	Normal weight	Reference	
	Obese	0.73	(0.39, 1.37)

Multiple linear or logistic regression, adjusted for ethnicity, parity, smoking, gestational age at delivery, and caesarean delivery. *Multiple logistic regression, adjusted for ethnicity, parity, smoking, gestational age at delivery, and caesarean delivery.