

Title	Trial of labour after caesarean section and the risk of neonatal and infant death: a nationwide cohort study
Authors	O'Neill, Sinéad M.;Agerbo, Esben;Khashan, Ali S.;Kearney, Patricia M.;Henriksen, Tine B.;Greene, Richard A.;Kenny, Louise C.
Publication date	2017-02-27
Original Citation	O'Neill, S. M., Agerbo, E., Khashan, A. S., Kearney, P. M., Henriksen, T. B., Greene, R. A. and Kenny, L. C. (2017) 'Trial of labour after caesarean section and the risk of neonatal and infant death: a nationwide cohort study', BMC Pregnancy and Childbirth, 17(74), pp. 1-9. doi:10.1186/s12884-017-1255-2
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1186/s12884-017-1255-2
Rights	© 2017, the Authors. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License ( <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> ), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver ( <a href="http://creativecommons.org/publicdomain/zero/1.0/">http://creativecommons.org/publicdomain/zero/1.0/</a> ) applies to the data made available in this article, unless otherwise stated. - <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>
Download date	2024-05-24 11:14:23
Item downloaded from	<a href="https://hdl.handle.net/10468/3827">https://hdl.handle.net/10468/3827</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

**Additional File 2 Neonatal and infant death according to mode of delivery – subgroup analysis restricted to women delivering between 38 and 40 weeks gestation**

<b>Mode of delivery 1<sup>st</sup> and 2<sup>nd</sup> birth</b>	<b>Neonatal death (<math>\leq 28</math> days) n=74</b>
	<b>* Model 1 AOR (95% CI)</b>
CS – ERCS	<i>Ref</i>
CS – TOLAC	1.78 (1.03, 3.08)
<b>Mode of delivery 1<sup>st</sup> and 2<sup>nd</sup> birth</b>	<b>Early neonatal death (<math>\leq 7</math> days) n=67</b>
	<b>* Model 1 AOR (95% CI)</b>
CS – ERCS	<i>Ref</i>
CS – TOLAC	1.99 (1.11, 3.58)
<b>Mode of delivery 1<sup>st</sup> and 2<sup>nd</sup> birth</b>	<b>Late neonatal death (<math>&gt; 7 \leq 28</math> days) n=7</b>
	<b>* Model 1 AOR (95% CI)</b>
CS – ERCS	<i>Ref</i>
CS – TOLAC	0.71 (0.13, 3.76)
<b>Mode of delivery 1<sup>st</sup> and 2<sup>nd</sup> birth</b>	<b>Infant death (<math>\leq 365</math> days) n=140</b>
	<b><sup>a</sup> Model 1 AOR (95% CI)</b>
CS – ERCS	<i>Ref</i>
CS – TOLAC	1.11 (0.76, 1.63)

**Table:** Data are adjusted odds ratios with 95% confidence intervals. **AOR**=adjusted odds ratio; **CI**=Confidence interval, **ERCS**: Elective repeat caesarean section; **TOLAC**=Trial of labour after caesarean section;

**\* Model 1:** adjusted for key covariates in the second birth including maternal age, maternal country of origin, educational attainment, mother and father's gross income, marital status, infant birthplace and infant birth weight, history of pregnancy loss and birth year (cohort n=48,711)