

Title	Metabolically healthy obesity across the life course: epidemiology, determinants and implications
Authors	Phillips, Catherine M.
Publication date	2016-10-10
Original Citation	Phillips, C. M. (2016) 'Metabolically healthy obesity across the life course: epidemiology, determinants, and implications', <i>Annals of the New York Academy of Sciences</i> , 1391(1), pp. 88-90 doi:10.1111/nyas.13230
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1111/nyas.13230
Rights	© 2016, New York Academy of Sciences. Published by John Wiley & Sons, Inc. This is the peer reviewed version of the following article: Phillips, C. M. (2016) 'Metabolically healthy obesity across the life course: epidemiology, determinants, and implications', <i>Annals of the New York Academy of Sciences</i> , which has been published in final form at http://onlinelibrary.wiley.com/doi/10.1111/nyas.13230/abstract . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Download date	2023-09-27 17:28:54
Item downloaded from	https://hdl.handle.net/10468/3320



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Metabolically healthy obesity across the life course: epidemiology, determinants and implications

Short title: Metabolically healthy obesity: a life course perspective

Catherine M. Phillips ^{1,2}

¹ HRB Centre for Diet and Health Research, Dept. of Epidemiology and Public Health, University College Cork, Ireland and ² HRB Centre for Diet and Health Research, School of Public Health, Physiotherapy and Population Science, University College Dublin, Ireland

Corresponding author: Dr. Catherine M. Phillips, HRB Centre for Diet and Health Research, Dept. of Epidemiology and Public Health, University College Cork, Ireland and School of Public Health, Physiotherapy and Population Science, University College Dublin, Ireland Email: c.phillips@ucc.ie
catherine.phillips@ucd.ie Telephone: +353 (1) 716 3483

Key words: Metabolically healthy obesity; Definitions; Determinants; Stability; Life course; Personalised medicine;

Word count: 6779

Abstract

In recent years different sub-phenotypes of obesity have been described including metabolically healthy obesity (MHO), whereby a proportion of obese individuals despite excess body fat remain free of metabolic abnormalities and increased cardiometabolic risk. In the absence of a universally accepted set of criteria to classify MHO the reported prevalence estimates vary widely. Our understanding of the determinants and stability of MHO over time and associated cardiometabolic and mortality risk is improving, but many questions remain. For example, whether MHO is truly benign is debatable and whether risk stratification of obese individuals based on their metabolic health status may offer new opportunities for more personalised approaches in diagnosis, intervention and treatment of diabetes remains speculative. Furthermore, as most research to date has focussed on MHO in adults little is known about childhood MHO. In this review we focus on the epidemiology, determinants, stability and health implications of MHO across the life course.

Introduction

Obesity has become a worldwide epidemic and represents a major public health challenge. Over the last four decades, there has been a global shift from a time when underweight was twice as prevalent as obesity to one in which the numbers of obese individuals now surpass those who are underweight, both globally and in all regions, with the exceptions of areas of Asia and sub-Saharan Africa ¹. Examination of body mass index (BMI) trends from 1698 population-based data sources involving more than 19.2 million adults (9.9 million men and 9.3 million women) revealed that in 2014 the prevalence of obesity exceeded underweight in both men and women in 68% and 83% of the countries for whom estimates were available ¹. Recent estimates indicate that about 266 million men and 375 million women are obese in the world ¹. Future projections predict that over one billion people, or approximately 20% of the world's entire adult population, will be obese by 2030 ². Not only is the prevalence of adult obesity increasing so too is childhood obesity. Approximately one quarter of children worldwide are overweight or obese ^{3, 4}. Although there are indications that the prevalence of overweight and obesity in children is plateauing in some populations ⁴, current levels remain high. This is particularly concerning as it has been shown that childhood obesity tracks to adulthood and is associated with increased risk of cardiometabolic disease and premature mortality ⁵. The current obesity epidemic is paralleled by escalating prevalence of type 2 diabetes mellitus (T2DM) and metabolic syndrome (MetS), a constellation of metabolic perturbations including obesity, insulin resistance, hypertension and dyslipidaemia. However over recent years it has become increasingly recognised that obesity is not a homogeneous condition and that a subset of obese individuals do not display disturbed metabolic profiles or increased risk of cardiometabolic disease. As such these individuals, who are characterised by preserved insulin sensitivity, normal blood pressure and lipid profiles despite their adiposity, may be described as being metabolically healthy obese (MHO). Similarly there exist normal weight individuals who display abnormal metabolic profiles and carry increased cardiometabolic risk. Thus a spectrum of metabolic health phenotypes according to BMI or body composition exist, ranging from metabolically healthy normal weight to metabolically unhealthy obese (MUO), along which MHO represents an intermediate stage. Research

on obesity phenotypes has mainly focussed on MHO among adults to date with a paucity of information available on childhood MHO. Therefore, in this review we focus on the epidemiology, determinants, stability and health implications of MHO over the life course from childhood into adulthood.

Metabolically healthy obesity in childhood and adolescence

The currently increasing prevalence of a range of adult chronic non-communicable diseases, including obesity, cannot be explained solely by genetic or adult lifestyle factors. Evidence from epidemiological studies suggest that the origins of many chronic diseases, including obesity, MetS, cardiovascular disease (CVD) and T2DM, actually begin in early life and that childhood insulin resistance and adiposity predispose to diabetes, MetS and CVD in adulthood⁶⁻¹⁰. Increasing evidence also suggests that early life exposure to a range of environmental factors, including nutrition for example, plays a critical role in defining offspring health, both in childhood and in later life. According to the Developmental Origins of Health and Disease hypothesis, transient environmental exposures during critical periods of development (such as the pre-conceptional, fetal and early infant phases of life) can alter normal physiology and have a persistent impact on metabolism and gene expression thereby influencing offspring phenotype and disease risk in later life^{11, 12}. For example, growth velocity in utero and in early life, which are a read-out of early nutritional status, may affect cardiovascular and cancer risk. Stefan *et al.*, recently reviewed the literature on height (a genetically determined phenotype which is influenced by maternal and early life exposures) and cardiometabolic disease and cancer risk later in life¹³. Accumulating data suggest divergent associations, such that height is associated with lower cardiometabolic risk and higher cancer risk. The authors speculate that overnutrition, particularly of milk and dairy products, during child development may play a role. Thus avoiding such overnutrition during critical developmental periods may attenuate accelerated growth and development of obesity in children leading to lower cancer risk in later life. Therefore, investigating metabolic health and adiposity in childhood may have implications in terms of

preventative strategies for adverse metabolic health phenotypes in early life which may have long term impacts.

As is the case for adults no standard metabolic health definition exists for use in children. **Table 1** details some currently used criteria to define MHO among children and adolescents ¹⁴⁻²⁰. While metabolic health may be simply defined as the absence of insulin resistance ¹⁷, in the majority of studies MHO characterisation centres on the absence of MetS (or some of its cardiometabolic risk factors) ¹⁴⁻²⁰ among those with excess body weight. It is important to note that how obesity is classified is also an issue, with different anthropometric measures and cut-offs being used. The Bogalusa Heart Study defined overweight/obesity as a BMI in the top quartile ¹⁶, the cross-sectional study by Prince *et al.*, defined obesity as a BMI $\geq 85^{\text{th}}$ percentile ¹⁷, the studies by Vukovic *et al.*, ^{18, 19} and Camhi *et al.*, ^{14, 21} defined obesity as a BMI $\geq 95^{\text{th}}$ percentile, whereas Zamrazilova *et al.*, ²⁰ used a BMI $\geq 97^{\text{th}}$ percentile to classify obesity. The Korean Children-Adolescent Study was based on abdominal obesity assessed by waist circumference ¹⁵. Such variation in classification of both obesity and metabolic health undoubtedly hinders comparisons between studies and contributes to the observed disparity in MHO prevalence in these studies (ranging from 4.2% ¹⁶, 21.5% ¹⁷, 21.7% ¹⁸, 25% ¹⁹, 53% ¹⁵ to 68% ¹⁴).

The determinants of and molecular mechanisms underlying MHO among children are under examined and as a result, poorly understood. Collectively the data indicate that MHO children are more likely to be younger (and in earlier stages of puberty) and female, of high birth weight with less visceral fat, preserved insulin sensitivity, high adiponectin concentrations, altered ghrelin levels, more favourable lipid profiles, reduced concentrations of transaminases and uric acid, and without hepatic steatosis ^{17-19, 22-28}. Although early weight gain was originally identified as a determinant of childhood MHO ²³ a larger study of Danish men for whom childhood BMI was available failed to find robust evidence to support a role for either rapid BMI growth or early-onset obesity in the development of MHO ²⁹. Aside from laboratory and clinical based predictors limited data regards the role of environmental, lifestyle, behavioural or genetic factors in determining MHO status among children exist. Prince *et*

al., examined lifestyle predictors of MHO among 8-17 year olds attending a weight management clinic ¹⁷. They identified dietary fat intake and moderate-to-physical activity as independent predictors. In contrast Camhi *et al.*, reported that physical activity, but not sitting or screen time differs between MHO and MUO in adults, but not in adolescents ¹⁴. In further work these authors also demonstrated that MHO adolescents have better compliance with dietary guidelines compared to their MUO counterparts ²¹. A study of 1213 Chinese children aged 6-18 years reported that walking to school and frequency of soft drink consumption, together with other demographic factors, contribute to the prediction of MHO status ³⁰. Interestingly this study also examined 22 genetic variants previously identified from genome wide association studies of obesity and diabetes. They found that both genetic predisposition and lifestyle factors and their interaction are independent predictors of MHO. More recently a study of teenage boys (13-17.9 years) investigated a range of potential determinants of MHO including duration of exposure to obesity, dietary intake and lifestyle factors. With the exception of carbohydrate intake, no other associations with MHO were identified with dietary or lifestyle factors. However a clear relationship between early onset of obesity and longer duration of its exposure with MUO was demonstrated ²⁰. Collectively these data suggest potential intervention windows and targets to improve cardiometabolic profiles in paediatric obesity with a view to achieving and maintaining better long term cardiometabolic health.

Metabolically healthy obesity in adulthood

The prevalence of MHO among adults varies greatly between studies. Although study specific differences such as age, ethnicity, sample size or environmental factors and genetics may be contributing factors, the lack of a universal definition of metabolic health and differences in obesity classification (BMI versus body fat percentage) account for a large proportion of the reported disparity ³¹. **Table 2** details some currently used criteria to define MHO among adults ³²⁻³⁸. At the very least metabolic health may be defined as the absence of insulin resistance ³⁵, but as is the case in children for the most part current characterisation of MHO in adults is based on the absence of MetS

(or some of its components) among those with excess body weight, generally defined by BMI³⁴⁻³⁸. Some definitions additionally include favourable inflammatory status determined by C reactive protein levels^{32, 33}.

Although limited, comparative studies examining MHO prevalence across a range of currently used criteria have reported considerable variation in MHO prevalence and poor agreement between MHO definitions^{39, 40}. In an Irish cross-sectional population representative sample of adults aged 50-69 years, among the obese (by BMI) participants MHO prevalence ranged from 6.8%-36.6%³⁹. MHO prevalence was generally higher among females and conflicting associations were noted with age, depending on which definitions were used. In a Korean study of 186 obese (by BMI) male subjects Yoo *et al.*,⁴¹ reported MHO prevalence ranging from 24.2% to 70.4%. This study did not observe any age related differences, which may be accounted for by the narrow and younger age range of the male only study participants (mean age 37 years). In a Swiss population-based sample of adults aged 35-75 years MHO prevalence ranged from 3.3-32.1% in men and 11.4-43.3% in women, with higher estimates reported when obesity was defined by body fat percentage (BF%)⁴⁰. The latter finding highlights the importance of how obesity is classified. Direct measurement of body fat using dual-energy X-ray absorptiometry (DEXA) is the gold standard, however DEXA is generally not available or practical for most researchers. BMI is the most widely used method to classify excess adiposity, with values $\geq 30 \text{ kg/m}^2$ indicating obesity. However BMI does not discriminate between fat and lean body mass, thus individuals of short stature or muscular build may be misclassified. Indeed, accumulating evidence indicates that BMI may actually under estimate obesity prevalence⁴². Similarly, obesity classification influences MHO prevalence, with estimates ranging from a third to half of obese individuals by BMI and DEXA respectively⁴³. Furthermore, recent data suggest that combined assessment of BMI and body fat percentage or other anthropometric measures to classify obesity may help identify individuals at greater cardiometabolic risk than BMI alone^{44, 45}. Collectively these data underscore the importance of accurate obesity diagnosis in the context of more precise classification and risk stratification.

MHO prevalence has been recently examined in a number of large scale studies. Analysis of 10 cohort studies involving 163,517 individuals from 7 European countries revealed significant diversity in MHO prevalence across Europe (7-28% in women and 2-19% in men) ³⁸. Generally, MHO, which was defined as obesity (by BMI) without any MetS component and no previous CVD diagnosis, was more prevalent among women and decreased with age in both genders ³⁸. A worldwide meta-analysis of 31 studies reported overall prevalence of MHO of 7.27%, with highest prevalence among American populations, although wide ranges were reported between individual studies (MHO prevalence of 1.3-22.9% in Americans, 2.1-23.9% in Europeans and 2.8-25.8% in Asians) ⁴⁶. Despite study design and population differences, the observed variation in MHO prevalence reported both in the comparative studies and meta-analyses highlight the need for larger scale population representative studies, improved obesity classification and a global consensus on a standard MHO definition.

Characterisation and determinants of metabolically healthy obesity

Although the determinants of MHO and the molecular mechanisms underlying the MHO phenotype are not fully elucidated, accumulating evidence is improving our understanding of the biological factors which distinguish MHO on the one hand from obesity per se, and on the other hand, from metabolically unhealthy obesity ^{47, 48}. Here we summarise the recent literature regards determinants of metabolic health status, with a focus on potential biological, environmental and genetic factors involved in the pathogenesis of MHO.

Adiposity and body composition

It has been recognised that different body fat distribution patterns, most notably increased visceral adipose tissue (VAT) and ectopic fat deposition (intramuscular, hepatic and epicardial) are related to different metabolic phenotypes and obesity-related cardiometabolic risk ⁴⁹⁻⁵¹. Greater VAT is associated with impaired glucose tolerance, insulin resistance, increased secretion of very low density lipoprotein (VLDL) and increased intra-hepatic triglyceride content (IHTG) ⁵²⁻⁵⁶. Hepatic fat has been

identified as a potential predictor of the MUO phenotype, type 2 diabetes and subclinical atherosclerosis⁵⁷⁻⁶¹. It has been hypothesised that how the body channels surplus energy, arising from a combination of excessive caloric intake, reduced physical activity and increased time spent in sedentary behaviour, may determine an individual's predisposition to MHO or MUO. The pathway to MUO may be characterised by dysfunctional adipose tissue (larger fat cells), increased immune cell infiltration, raised pro-inflammatory status and reduced capacity of subcutaneous adipose tissue to expand leading to increased ectopic fat deposition resulting in lipotoxicity, insulin resistance in peripheral tissues and a range of metabolic derangements (**Figure 1**)⁶². Whereas in MHO individuals the excess calories are channelled into insulin-sensitive subcutaneous adipose tissue which is capable of expansion, thus visceral and ectopic adiposity are reduced, macrophage infiltration and raised pro-inflammatory state are attenuated, insulin sensitivity is preserved and the individual is protected from development of the MetS^{47, 63-66}.

It has been questioned whether increased VAT is just an innocent bystander acting as a marker of ectopic fat deposition or is indeed the culprit⁶⁷. In a small study of 39 obesity-matched adolescents and adults Linder *et al.*, compared the impact of body fat distribution and ectopic fat, in particular liver fat, on insulin resistance with a view to establishing whether previously identified relationships in adults between liver fat and insulin resistance hold true among adolescents. Despite having lower VAT the overweight and obese adolescents were more insulin resistant than the gender and BMI-matched adults. Of note hepatic fat content, but not total body fat or VAT, was identified as an independent predictor of insulin resistance among both adolescents and adults⁶⁸. An elegant study by Fabbrini *et al.*, examined the independent association of VAT and IHTG to metabolic function. They demonstrated increased VLDL-TG secretion and impaired insulin action in adipose tissue, skeletal muscle and liver of obese subjects with high IHTG, but not among those with high VAT matched for IHTG. Furthermore high IHTG was associated with altered expression and protein levels of CD36 (a protein involved in fatty acid metabolism) in adipose tissue and skeletal muscle, suggesting a role in ectopic fat accumulation. The authors concluded that IHTG, not VAT, is a better

indicator of metabolic functionality associated with obesity⁶⁹. In later work this group examined the impact of weight gain among obese individuals defined by IHTG and insulin sensitivity as MHO and MUO⁷⁰. Both groups were challenged with a high-fat diet to achieve ~6% weight gain. Despite similar fat mass increases there were distinct differences in the response to weight gain between the groups. Insulin sensitivity in adipose tissue, liver and skeletal muscle deteriorated and blood pressure, plasma TG, VLDL apoB100 concentrations and secretion rates increased among the MUO, but not the MHO individuals, suggesting that MHO subjects are protected against the adverse effects of weight gain. Such protection may be derived from the increased biological pathways and genes associated with adipose tissue lipogenesis observed among the MHO, but not the MUO, individuals⁷⁰. From this evidence it seems that body composition and fat distribution, in particular VAT and hepatic fat, are both important players in determining cardiometabolic health status.

Diet and lifestyle factors

Environmental factors such as diet, physical activity, alcohol consumption and smoking play a role in the development of obesity. Simply put excessive caloric intake coupled with low levels of physical activity and/or increased sedentary time give rise to a positive energy balance leading to increased body fat accumulation. In the context of MHO diet, including dietary composition, dietary patterns and dietary quality, has been fairly widely studied. However the evidence supporting the role of diet in MHO has been surprisingly inconsistent to date. Interestingly similar total energy intake and dietary macronutrient intakes have been reported in MHO and MUO individuals^{39, 71-73}, leading researchers to examine dietary patterns, indices of dietary quality and compliance with dietary recommendations. A recent cross-sectional study involving 2415 middle-aged Australian adults reported that for every one standard deviation increase in the healthy dietary pattern, the likelihood of having a more metabolically healthy profile increased by 16% (OR 1.16, 95% CI 1.04-1.29)⁷⁴. Using NHANES data (2007-2008 and 2009-2010) Camhi *et al.*, examined dietary quality assessed by the Healthy Eating Index 2005 (HEI-2005) scores among obese adolescents (n=133) and adults (n=1102)²¹. MHO was defined as obesity with two or less abnormal risk factors for fasting glucose,

hypertension, triglycerides and HDL-C¹⁴. HEI-2005 scores were higher among the MHO adolescents and women (aged 19-44 years) relative to their MUO counterparts, whereas no differences were noted among MHO and MUO men (aged 19-44 or 45-85 years). Examination of scores from specific food groups revealed that MHO adolescents had higher milk scores and scores from added sugars, solid fats and alcoholic beverages. Among the 19-44 year-old women, higher scores for whole fruits, whole grain, meat and beans were reported. These findings highlight the potential of dietary quality indices as intervention targets and the importance of such intervention starting earlier in life, as differences were only observed among adolescents and younger women. Park *et al.*, investigated Mediterranean Diet Scores (MDS) among 1739 adult participants of the National Health and Nutrition Examination Survey III (1988-1994) who were followed up for deaths until 2011⁷⁵. Metabolic health was defined using the Wildman definition³². Consumption of red meat and dairy products were lower among the MHO individuals, who also had a higher ratio of monounsaturated to saturated fatty acids, which contributed to their higher MDS. Furthermore, adherence to a Mediterranean style diet was associated with lower all-cause mortality among the MHO individuals (multivariable-adjusted hazard ratio of 0.44, 95% confidence interval (0.26-0.75) comparing the highest tertile to the first tertile of MDS), but not among the MUO subjects, perhaps suggesting that alternative strategies are required for MUO.

Given the range of MHO criteria available, we investigated in a cross-sectional cohort of 2047 middle-aged men and women to what extent differences between metabolically healthy and unhealthy obese and non-obese subjects, defined using a variety of metabolic health definitions may be explained by dietary composition, dietary quality and food pyramid compliance³⁹. In keeping with previous findings total calorie intake, dietary macronutrient composition and also dietary quality were generally similar between the MHO and MUO individuals across MHO definitions. However better compliance with food pyramid recommendations was positively associated with MHO (defined by insulin resistance and Wildman). Furthermore, some differences in the number of daily servings of fruit and vegetables, dairy, meats, fats and high fat/sugar food and drinks were noted between the MHO and MUO subjects, depending on which MHO criteria used. Of note there was generally no effect of physical activity, smoking or alcohol intake observed between MHO and MUO individuals

across the range of MH definitions examined. Collectively these findings highlight the potential of dietary guidelines as intervention targets to improve cardiometabolic health status among obese individuals.

Diet and lifestyle interventions

Limited and inconsistent data regards the impact of dietary and exercise interventions in MHO exist⁷⁶⁻⁷⁹. Rondanelli *et al.*, reported significant improvements to a range of metabolic measures including HOMA, CRP, HDL-C, leptin, adiponectin, ghrelin, glucagon like peptide-1 and fatty acid profiles among 103 MHO individuals following a 2 month prudent dietary intervention⁸⁰. Unfortunately this study did not include MUO individuals. Kantartzis *et al.*, examined 262 MHO and MUO individuals, defined by HOMA and BMI, following a 9 month lifestyle intervention programme. Visceral fat was reduced in both groups post-intervention, however total body and liver fat was reduced among the MUO subjects only. These individuals also reported improvements in insulin sensitivity, although they remained insulin resistant⁷⁷. In contrast Ruiz *et al.*, did not observe any differences in the magnitude of change in anthropometric measures between MHO (n=25) and MUO (n=53) women following a 12-week energy-restricted dietary intervention⁸¹. Janiszewski *et al.*, reported reductions in body weight, total and visceral fat mass and enhanced insulin sensitivity in both MHO (n=63) and MUO (n=43) subjects following a 3-6 month exercise or diet-induced weight loss intervention, with greater improvements in insulin sensitivity among the MUO subjects⁷⁶. A 5-10% reduction in body weight is considered to be clinically significant and as such can improve metabolic health among obese individuals. Lui *et al.*, investigated the impact of a 5% lifestyle-based weight loss on the metabolic profiles of 392 MHO and MUO individuals⁸². Among those who achieved target weight loss, improvements to most risk factors were observed regardless of metabolic health status, suggesting that a clinically significant weight loss is beneficial to all obese individuals.

Cardiorespiratory fitness

Higher levels of cardiorespiratory fitness (CRF) are independently associated with healthier metabolic profiles and reduced risk of incident CVD and CVD mortality⁸³⁻⁸⁶. However the roles of physical and cardiorespiratory fitness (CRF) have not been extensively investigated in the context of understanding the determinants of MHO status. Dalleck *et al.*, examined metabolic syndrome components in 332 adults before and after a supervised 14 week community-based exercise program designed to improve cardiometabolic risk factors⁸⁷. This short-term intervention, which improved cardiorespiratory fitness (assessed by conventional submaximal exercise test protocols for walking or cycle ergometry) and eliminated MetS features, positively transitioned MUO individuals to MHO status. The greatest results were observed among those engaging in higher volumes of exercise, suggesting community based exercise as an effective model for primary prevention of cardiometabolic disease. Ortega *et al.*, examined fitness (assessed by a maximal exercise test on a treadmill), body fat composition and metabolic health status among 43, 265 adults participating in The Aerobics Center Longitudinal Study⁸⁸. Their findings suggest that the MHO phenotype, defined using BMI or BF%, is associated with better CRF in both men and women, and that once CRF level is accounted for the MHO phenotype may be benign in terms of mortality and morbidity risk. These authors recently reviewed the current evidence regards CRF and MHO from cross-sectional and longitudinal studies⁸⁹. They conclude that better CRF should be considered to be a characteristic of the MHO phenotype, signal that caution should be taken regards whether CRF plays a role in the prognosis of the MHO individuals, and recommend further investigation of the role of CRF in MHO.

Genetics

Limited data regards genetic predisposition to MHO exists. As mentioned already the Beijing Children and Adolescents Metabolic Syndrome study (BCAMS) explored the contribution of both genetic and environmental factors to the pathogenesis of MHO among 6-18 year olds³⁰. Although a limited number of genetic variants (22 single nucleotide polymorphisms (SNPs)) were examined, both the *KCNQ1* rs227892 and rs227897 SNPs were identified as independent predictors of MHO. Each additional C allele of either SNP was associated with reduced risk of being metabolically healthy

based on both cardiometabolic risk markers (23% lower risk) and insulin resistance (24% lower risk), with stronger associations identified when a composite genetic predisposition score was examined.

Interestingly this study provided the first evidence of gene-nutrient (soft drink consumption) and gene-environment (with walking to school) interactions predisposing to MHO. Clearly much work remains to be done in this area to uncover the influence of genetics, nutrigenetics and epigenetics on MHO pathogenesis. Berezina *et al.*, examined potential relationships between genetic variants of the adipocytokine genes (leptin, leptin receptor and adiponectin) and metabolically healthy (without CVD) abdominal obesity among adults ⁹⁰. Although genetic associations and gene-nutrient interactions have been described previously between genetic variants of these genes and MetS ^{91, 92}, this study identified for the first time a more than two fold greater likelihood of MUO among the T allele carriers of the adiponectin T45T polymorphism relative to the G allele carriers. Genetic predisposition to weight and metabolic health related traits has been investigated longitudinally in almost 4000 adult and 1380 adolescent participants of the Norwegian HUNT2 (1995-1997) and HUNT3 (2006-2008) surveys ⁹³. Examination of 27 SNPs previously associated with obesity, eating disorders or metabolic risk revealed novel genetic associations between a number of genes involved in regulation of food intake and energy expenditure, eating behaviour, food reward and satiety with longitudinal changes in BMI/WC and development of adverse metabolic phenotypes. Such data highlight the importance of not only improving our current understanding of the genetics but also the neurobiology of body weight regulation in the context of developing future strategies to combat obesity, and especially MUO.

Stability of metabolically healthy obesity across the life course

MHO was initially regarded as a static condition and, although some individuals can maintain their metabolic health status over time, it is becoming increasingly evident that MHO status is transient in nature. It has been suggested that the MHO phenotype starts in childhood and persists into adulthood. The Bogalusa Heart Study is unique in the context of examining MHO stability over time, in that the 1098 individuals participated in the study both as children (aged 5-17 years) and also as young adults

(24-43 years), with an average follow-up of 24 years (range 14.1-28.6 years)¹⁶. Importantly this study provides the opportunity to examine what happens to MHO status as we age from childhood, through adolescence into adulthood. The results are intriguing. The MHO children display similar favourable cardiometabolic profiles when adults relative to their childhood MUO counterparts. On the other hand the MUO children had the worst cardiometabolic profiles as adults. Although adult MHO status was maintained in only 13% of the MHO children, the MHO children were 2.7-9.3 times more likely to be MHO adults compared to children from the other metabolic health categories. Even though the MHO children displayed intermediate levels of insulin, glucose and blood pressure as adults, suggesting intermediate risk of T2DM and hypertension, examination of carotid intima media thickness (CIMT), a marker of atherosclerosis, did not reveal increased CIMT in adulthood¹⁶. As the cardiometabolic profiles of MHO adults have been shown to be more favourable than that of metabolically unhealthy normal weight individuals and more comparable to those of their normal weight counterparts, a better understanding of what factors contribute to achieving and maintaining good metabolic health from childhood into adulthood is critical.

As already mentioned while some individuals can retain MHO status over time a substantial proportion of individuals with MHO cannot and may become metabolically unhealthy. Indeed, it is possible for any individual to transition between metabolically healthy and unhealthy states regardless of their BMI. Such transitions may also contribute to the observed disparity in MHO prevalence, inverse association with age and conflicting findings regards cardiometabolic and mortality outcomes. Longitudinal investigations suggest that MUO is a progressive phenotype along which MHO represents a dynamic intermediate stage. Data from some of the earlier studies indicate that MHO status is transient for about a third of individuals. Follow-up (5.5-10.3 years) of the North West Adelaide Health Study cohort of 4,056 adults revealed that 33% of the MHO subjects became MUO over time, whereas for the remaining individuals persistent MHO status was associated with favourable cardiometabolic outcomes⁹⁴. In keeping with these findings data from the Pizarra study indicate that 37% of MHO subjects were no longer metabolically healthy after a 6-year follow-up⁹⁵. More recent data suggest that the numbers of MHO individuals becoming unhealthy over time are

actually greater. Longitudinal follow-up of the Tehran Lipid and Glucose Study revealed that 43.3% of the metabolically healthy abdominally obese transitioned to MUO over a 10 year period ⁹⁶. Data from the English Longitudinal Study of Ageing indicate that 44.5% of the MHO individuals became MUO over the 8-year follow-up ⁹⁷. Consistent with these findings data from the San Antonio Heart Study suggest that almost half (47.6%) of MHO subjects at baseline transitioned to MUO over the follow-up period (median 7.8 years) ⁹⁸.

Characterisation of the factors which distinguish those who progress to or maintain MHO from those who transition from MHO to MUO may uncover potential intervention targets. In the San Antonio Heart Study those who transitioned were older, had lower HDL cholesterol levels and increased adiposity compared to the individuals with persistent MHO ⁹⁸. Interestingly none of the adiposity measures (BMI, waist circumference and weight gain) were significant predictors of this change ⁹⁸. Moreover, lipid profiles emerged as the strongest determinants of metabolic health status likely to develop with weight gain. In addition to baseline lipid concentrations (triglycerides and HDL-cholesterol), findings from the Tehran Lipid and Glucose Study indicate that insulin resistance is a significant predictor of the change from MHO to MUO ⁹⁶. Additional predictors have been identified in the English Longitudinal Study of Ageing. Compared to those with persistent MHO, those who converted to MUO were more likely to have high blood pressure, display increased abdominal adiposity and have elevated levels of C-reactive protein, glycated haemoglobin and triglycerides ⁹⁷. Collectively these findings highlight the importance of healthy lipid and inflammatory profiles in achieving and maintaining optimal cardiometabolic health. Further characterisation of persistent metabolic health status and longitudinal investigation of the sustainability and predictors of the MHO phenotype over the life course is warranted.

MHO and long term health outcomes

The individual and joint contributions of metabolic health and BMI on long term cardiometabolic health outcomes and mortality are yet to be fully elucidated and further investigation is required. A large systematic review and meta-analysis of 2.88 million individuals confirmed significantly higher all-cause mortality with obesity when all grades are combined ⁹⁹. However, examination of individual obesity grades revealed that grade 1 obesity (BMI 30 to < 35 kg/m²) was not associated with higher mortality. These conflicting findings may be partly explained by the existence of different obesity associated metabolic health phenotypes. Examination of trends in metabolic health in the Northern Sweden MONICA study from 1986 to 2009 demonstrated that more people are becoming overweight and obese, and that a larger proportion of those individuals are metabolically healthy ¹⁰⁰, which has been suggested may reduce the impact of obesity as a CVD risk factor ¹⁰¹. Supporting this idea, a 20 year follow-up of the Atherosclerosis Risk in Communities Study recently reported intermediate risk for stroke, coronary heart disease and survival probability in individuals with suboptimal health (≤ 2 cardiometabolic risk factors), between that of the healthy and unhealthy subgroups, with no effect of BMI ¹⁰², suggesting that metabolic health may be more important than BMI in the context of adverse cardiometabolic outcomes.

Findings from prospective studies tracking the development of CVD, T2DM and mortality in MHO have been inconsistent ^{101, 103-108}. Thus whether MHO represents true health among obese individuals is controversial and remains the subject of ongoing debate ¹⁰⁹⁻¹¹². Examination of all-cause and CVD mortality after 17.7 years follow-up of the Whitehall II cohort of 5,269 adults aged 39-62 (prevalence of MHO 9-41% depending on definition used) revealed that both the MHO and MUO subjects had increased mortality risk (hazard ratio (HR) ranged from 1.81; 95% CI 1.16–2.84 to 2.30; 95% CI 1.13–4.70 for MHO and from 1.57; 95% CI 1.08–2.28 to 2.05; 95% CI 1.44–2.92 for the MAO) relative to the metabolically healthy normal weight subjects ¹⁰⁶. Furthermore, increased risk of both incident CVD and T2DM was reported among MHO individuals relative to their healthy normal weight counterparts ¹¹³. However, the MHO individuals were at a lower risk of T2DM but not CVD, compared to the MUO subjects. Thus MHO may not be as benign as initially thought and results are largely dependent on what outcome is examined and what reference group is used. In the Uppsala

Longitudinal Study of Adult Men (30 years follow-up of 1,758 subjects) increased mortality risk was identified in obese subjects with and without the MetS (2.4 and 1.7 fold higher, respectively) relative to the normal weight participants without the MetS ¹⁰³. The Third National Health and Nutrition Examination Survey (NHANES III) (8.7 years follow-up of 6,011 subjects), reported similar increased mortality risk (approximately 2.8-fold) between obese participants with ≤ 1 MetS feature and obese subjects with ≥ 2 MetS features relative to their metabolically normal non-obese counterparts ¹⁰⁷. Furthermore using waist circumference rather than BMI to classify obesity, follow-up (average time of 13.4 years) of the EPIC-MORGEN cohort of 22,654 individuals aged 20-59 years revealed higher mortality risk among metabolically healthy abdominally obese (MHAO) subjects relative to their metabolically healthy non-obese (MHNAO) counterparts (HR 1.43; 95% CI 1.00-2.04) ¹⁰⁸. Similar HRs were obtained for the metabolically unhealthy not abdominally obese subjects (MUNAO) (HR 1.31; 95% CI: 1.08-1.59), whereas higher HRs were identified for the metabolically unhealthy abdominally obese (MUAO) subjects (HR 1.99; 95% CI: 1.62-2.43 NS) ¹⁰⁸. Moreover, the Study of Women's Health across the Nation (SWAN) of 475 middle-aged women reported greater subclinical CVD burden among the metabolically healthy overweight/obese subjects relative to the MHNO women ¹¹⁴. Collectively these findings suggest that obese subjects, whether metabolically healthy or not, and regardless of how MHO is defined carry greater risk of CVD and mortality and thus MHO may not be as apparently healthy as originally considered.

Conversely several studies have not reported higher risk of CVD and all-cause mortality among their MHO participants. Seven year follow-up of 22,303 men and women (mean age 54.1 years) from the Health Survey for England and Scottish Health Survey failed to demonstrate increased risk of CVD (HR 1.26; 95% CI 0.74-2.13) or all-cause mortality (HR 0.91; 95% CI 0.64-1.29) among MHO subjects (defined by NCEP ATP III, the Wildman definition and BMI) relative to their metabolically healthy non-obese counterparts ¹⁰¹. Of note increased risk of all-cause mortality was observed among the MUO individuals (HR 1.72 95%CI 1.23-2.41) compared to the MHO subjects. Similar results were obtained when WC was used to define obesity. Calori *et al.*, in a follow-up of 2,011 middle-aged adults over 15 years, reported increased CVD, cancer and all-cause mortality risk (HR 1.40; 95% CI

1.08-1.81) among the obese insulin resistant individuals, but not in the obese insulin sensitive (MHO) subjects relative to their non-obese insulin sensitive counterparts ¹⁰⁴. In addition to higher prevalence of CVD, greater severity of angiographic CAD has also been reported in a study of 856 Korean subjects among the metabolically unhealthy (defined by NCEP ATP III) obese or normal weight subjects compared to the MHO or MHNO groups ¹¹⁵. Examination of mortality risk in NHANES III (12-18 years follow-up of 4,373 men and women) demonstrated that MHO individuals (defined according to HOMA, NCEP ATP III, Karelis definition) were not at increased risk of all-cause mortality compared to the MHNO individuals ¹⁰⁵. More recently Guo *et al.*, investigated the relative impact of body weight and metabolic health on health outcomes using data from 2 large cohorts (Coronary Artery Risk Development in Young Adults Study and the Atherosclerosis Risk in Communities Study, with 18.7 and 20 years follow-up, respectively) ¹⁰². They reported lower risk for T2DM, CVD, stroke and mortality among the MHO individuals relative to the MUO subjects, but increased diabetes risk compared to the MHNO subjects. Clearly the data on long term impact of MHO on cardiometabolic health and mortality risk is conflicting, which may be at least partly due to differences in study design, obesity classification, MHO definitions and reference groups. Whether obesity or metabolic health is a more important predictor of future health and/or disease remains unclear and further investigation of obesity associated metabolic health phenotypes is warranted.

MHO: role in risk stratification and personalised treatment?

Obesity is a multifaceted public health problem; the sheer complexity of the interacting biological, environmental and social determinants has been nicely illustrated by the UK Foresight obesity systems map ¹¹⁶. However, it is becoming apparent that the situation is further complicated at a personal level by the existence of subtypes of obesity based on an individuals' metabolic health status. Despite an ever increasing evidence base which has highlighted potential intervention points including food production and consumption, physiology, individual physical activity, the physical activity environment, and both individual and social psychology, obesity prevalence continues to rise.

This begs the question of whether more personalised strategies to combat obesity, based on an individual's metabolic health background, may offer new opportunities in obesity diagnosis, intervention and treatment.

The existence, clinical utility and limitations of the MHO phenotype have been widely questioned and debated ^{109-112, 117-119}. While the lack of a universally accepted MHO definition and usefulness of BMI to accurately classify obesity are clearly pertinent issues, the concept that any form of obesity could be described as healthy is controversial. In a recent commentary Rey-Lopez argued that “more efforts must be allocated to reducing the distal and actual causal agents that lead to weight gain, instead of the current disproportionate scientific interest in the biological processes that explain the heterogeneity of obesity” ¹¹². However perhaps the heterogeneity of obesity, in terms of an individuals' phenotype and inter-individual differences in responsiveness to dietary or lifestyle interventions, should not be ignored. Recent evidence indicates that despite similar overall dietary intake between metabolic health subtypes that favourable lifestyle factors including higher dietary quality, healthy diet pattern, greater compliance with food pyramid recommendations, being less sedentary and more (moderately) physically active may all be positively associated with MHO ^{21, 39, 74, 120}. Interestingly examination of stable and unstable MHO suggests that a healthy lifestyle index may determine transition to MUO ¹²¹. Furthermore a recent proteomics study identified dysregulated inflammatory and lipid processes as molecular hallmarks of MHO ¹²², confirming earlier findings that MHO individuals display more favorable lipoprotein ¹²³ and inflammatory profiles ⁶⁵. Collectively such investigations may identify new behavioural and biological targets which may aid the development of more effective evidence based risk stratification, intervention and treatment strategies to reduce both obesity and its metabolic complications.

Supporting this concept, the American Association of Clinical Endocrinologists (AACE) in 2014 suggested a complication-centric approach to the management of weight loss, whereby more aggressive therapeutic approaches for those patients with obesity-related complications were advocated ¹²⁴. More recently the AACE and the American College of Endocrinology (ACE),

motivated according to the Chair of the AACE Obesity Scientific Committee by “the lack of comprehensive and evidence-based guidelines to real-world clinical care of patients with obesity”, have developed new clinical practice guidelines (CGP) which acknowledge the need for a more individualised treatment approach to obesity ¹²⁵. These evidence-based CPGs address a range of aspects of obesity care including screening, diagnosis, clinical evaluation, treatment options, selection and goals. A notable shift here is the additional target of improving metabolic health rather than just weight loss per say. This development is timely, as it is evident that up to now approaches focussed on preventing and/or attenuating obesity and body weight have not achieved much success in halting the rising tide of obesity. While data on the impact of the new guidelines on the obesity epidemic will take some time to filter through it seems likely that high risk groups, such as the MUO who carry the greatest risk of both adverse cardiometabolic and mental health outcomes ^{101, 102, 104, 105, 115, 126}, could really benefit from such risk stratification. However, if metabolic health is a more important driver of future health than obesity it could be argued that improving metabolic health and attenuating development of cardiometabolic disease in intermediate risk subgroups (with or without obesity), such as MHO and metabolically unhealthy non-obese individuals, may also be worthwhile.

Conclusions

It is clear that a great body of research on obesity-associated metabolic health phenotypes has been performed to date. However much remains to be done. Despite the knowledge that different obesity subtypes exist the research community has been slow to refine obesity and metabolic health definitions. While advances in the development of new obesity treatment guidelines are encouraging whether these will have the desired impact on reducing obesity and its complications remains to be seen. Better understanding of both the lifestyle determinants of MHO and the molecular mechanisms that mediate the MHO phenotype is warranted. To advance the state-of-the-art future research will need to focus on these issues from a life course perspective, as well as conducting larger evidence-

based lifestyle intervention studies and longitudinal follow-up, with a view to opening up new avenues of personalised obesity medicine.

Acknowledgements

This work was supported by a research grant from the Irish Health Research Board (reference HRC/2014/13).

References

1. Non-Communicable Disease Risk Factor Collaboration, M. Di Cesare, J. Bentham, *et al.* 2016. Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet*. **387**: 1377-1396.
2. Kelly, T., W. Yang, C.S. Chen, *et al.* 2008. Global burden of obesity in 2005 and projections to 2030. *International journal of obesity (2005)*. **32**: 1431-1437.
3. Keane, E., P.M. Kearney, I.J. Perry, *et al.* 2014. Trends and prevalence of overweight and obesity in primary school aged children in the Republic of Ireland from 2002-2012: a systematic review. *BMC Public Health*. **14**: 974.
4. Olds, T., C. Maher, S. Zumin, *et al.* 2011. Evidence that the prevalence of childhood overweight is plateauing: data from nine countries. *Int J Pediatr Obes*. **6**: 342-360.
5. Reilly, J.J. & J. Kelly. 2011. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes (Lond)*. **35**: 891-898.
6. Dwyer, T., C. Sun, C.G. Magnussen, *et al.* 2013. Cohort Profile: the international childhood cardiovascular cohort (i3C) consortium. *Int J Epidemiol*. **42**: 86-96.
7. Franks, P.W., R.L. Hanson, W.C. Knowler, *et al.* 2010. Childhood obesity, other cardiovascular risk factors, and premature death. *The New England journal of medicine*. **362**: 485-493.
8. Charakida, M., A. Jones, E. Falaschetti, *et al.* 2012. Childhood obesity and vascular phenotypes: a population study. *J Am Coll Cardiol*. **60**: 2643-2650.
9. Liang, Y., D. Hou, X. Zhao, *et al.* 2015. Childhood obesity affects adult metabolic syndrome and diabetes. *Endocrine*. **50**: 87-92.
10. Raitakari, O.T., M. Juonala, M. Kahonen, *et al.* 2003. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *JAMA*. **290**: 2277-2283.
11. Barker, D.J. 1995. Fetal origins of coronary heart disease. *BMJ*. **311**: 171-174.
12. Godfrey, K.M. & D.J. Barker. 2001. Fetal programming and adult health. *Public Health Nutr*. **4**: 611-624.
13. Stefan, N., H.U. Haring, F.B. Hu, *et al.* 2016. Divergent associations of height with cardiometabolic disease and cancer: epidemiology, pathophysiology, and global implications. *The lancet. Diabetes & endocrinology*. **4**: 457-467.

14. Camhi, S.M., M.E. Waring, S.B. Sisson, *et al.* 2013. Physical activity and screen time in metabolically healthy obese phenotypes in adolescents and adults. *J Obes.* **2013**: 984613.
15. Chun, S., S. Lee, H.J. Son, *et al.* 2015. Clinical Characteristics and Metabolic Health Status of Obese Korean Children and Adolescents. *Korean J Fam Med.* **36**: 233-238.
16. Li, S., W. Chen, S.R. Srinivasan, *et al.* 2012. Relation of childhood obesity/cardiometabolic phenotypes to adult cardiometabolic profile: the Bogalusa Heart Study. *American journal of epidemiology.* **176 Suppl 7**: S142-149.
17. Prince, R.L., J.L. Kuk, K.A. Ambler, *et al.* 2014. Predictors of metabolically healthy obesity in children. *Diabetes care.* **37**: 1462-1468.
18. Vukovic, R., T. Milenkovic, K. Mitrovic, *et al.* 2015. Preserved insulin sensitivity predicts metabolically healthy obese phenotype in children and adolescents. *Eur J Pediatr.* **174**: 1649-1655.
19. Vukovic, R., K. Mitrovic, T. Milenkovic, *et al.* 2013. Insulin-sensitive obese children display a favorable metabolic profile. *Eur J Pediatr.* **172**: 201-206.
20. Zamrazilova, H., R. Weiss, V. Hainer, *et al.* 2016. Cardiometabolic health in obese adolescents is related to length of obesity exposure: A pilot study. *The Journal of clinical endocrinology and metabolism.* jc20161706.
21. Camhi, S.M., E. Whitney Evans, L.L. Hayman, *et al.* 2015. Healthy eating index and metabolically healthy obesity in U.S. adolescents and adults. *Prev Med.* **77**: 23-27.
22. Bluher, S. & P. Schwarz. 2014. Metabolically healthy obesity from childhood to adulthood - Does weight status alone matter? *Metabolism: clinical and experimental.* **63**: 1084-1092.
23. Bouhours-Nouet, N., S. Dufresne, F.B. de Casson, *et al.* 2008. High birth weight and early postnatal weight gain protect obese children and adolescents from truncal adiposity and insulin resistance: metabolically healthy but obese subjects? *Diabetes care.* **31**: 1031-1036.
24. Mangge, H., S. Zelzer, P. Puerstner, *et al.* 2013. Uric acid best predicts metabolically unhealthy obesity with increased cardiovascular risk in youth and adults. *Obesity (Silver Spring, Md.* **21**: E71-77.
25. Pacifico, L., E. Poggiogalle, F. Costantino, *et al.* 2009. Acylated and nonacylated ghrelin levels and their associations with insulin resistance in obese and normal weight children with metabolic syndrome. *European journal of endocrinology / European Federation of Endocrine Societies.* **161**: 861-870.
26. Reinehr, T., B. Wolters, C. Knop, *et al.* 2015. Strong effect of pubertal status on metabolic health in obese children: a longitudinal study. *The Journal of clinical endocrinology and metabolism.* **100**: 301-308.
27. Senechal, M., B. Wicklow, K. Wittmeier, *et al.* 2013. Cardiorespiratory fitness and adiposity in metabolically healthy overweight and obese youth. *Pediatrics.* **132**: e85-92.
28. Weghuber, D., S. Zelzer, I. Stelzer, *et al.* 2013. High risk vs. "metabolically healthy" phenotype in juvenile obesity - neck subcutaneous adipose tissue and serum uric acid are clinically relevant. *Exp Clin Endocrinol Diabetes.* **121**: 384-390.
29. Howe, L.D., E. Zimmermann, R. Weiss, *et al.* 2014. Do rapid BMI growth in childhood and early-onset obesity offer cardiometabolic protection to obese adults in mid-life? Analysis of a longitudinal cohort study of Danish men. *BMJ Open.* **4**: e004827.
30. Li, L., J. Yin, H. Cheng, *et al.* 2016. Identification of Genetic and Environmental Factors Predicting Metabolically Healthy Obesity in Children: Data From the BCAMS Study. *The Journal of clinical endocrinology and metabolism.* **101**: 1816-1825.
31. Phillips, C.M. 2013. Metabolically healthy obesity: definitions, determinants and clinical implications. *Rev Endocr Metab Disord.* **14**: 219-227.
32. Wildman, R.P., P. Muntner, K. Reynolds, *et al.* 2008. The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med.* **168**: 1617-1624.

33. Karelis, A.D. & R. Rabasa-Lhoret. 2008. Inclusion of C-reactive protein in the identification of metabolically healthy but obese (MHO) individuals. *Diabetes Metab.* **34**: 183-184.
34. Karelis, A., M. Brochu & R. Rabasa-Lhoret. 2004. Can we identify metabolically healthy but obese individuals (MHO)? *Diabetes & Metabolism.* **30**: 569-572.
35. Meigs, J.B., P.W.F. Wilson, C.S. Fox, *et al.* 2006. Body mass index, metabolic syndrome, and risk of type 2 diabetes or cardiovascular disease. *J Clin Endocr Metab.* **91**: 2906-2912.
36. Aguilar-Salinas, C.A., E.G. Garcia, L. Robles, *et al.* 2008. High adiponectin concentrations are associated with the metabolically healthy obese phenotype. *J Clin Endocr Metab.* **93**: 4075-4079.
37. 2001. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *Jama.* **285**: 2486-2497.
38. van Vliet-Ostaptchouk, J.V., M.L. Nuotio, S.N. Slagter, *et al.* 2014. The prevalence of metabolic syndrome and metabolically healthy obesity in Europe: a collaborative analysis of ten large cohort studies. *BMC Endocr Disord.* **14**: 9.
39. Phillips, C.M., C. Dillon, J.M. Harrington, *et al.* 2013. Defining metabolically healthy obesity: role of dietary and lifestyle factors. *PLoS one.* **8**: e76188.
40. Velho, S., F. Paccaud, G. Waeber, *et al.* 2010. Metabolically healthy obesity: different prevalences using different criteria. *European journal of clinical nutrition.* **64**: 1043-1051.
41. Yoo, H.K., E.Y. Choi, E.W. Park, *et al.* 2013. Comparison of Metabolic Characteristics of Metabolically Healthy but Obese (MHO) Middle-Aged Men According to Different Criteria. *Korean J Fam Med.* **34**: 19-26.
42. Gomez-Ambrosi, J., C. Silva, J.C. Galofre, *et al.* 2012. Body mass index classification misses subjects with increased cardiometabolic risk factors related to elevated adiposity. *International journal of obesity (2005).* **36**: 286-294.
43. Shea, J.L., E.W. Randell & G. Sun. 2011. The prevalence of metabolically healthy obese subjects defined by BMI and dual-energy X-ray absorptiometry. *Obesity (Silver Spring, Md.* **19**: 624-630.
44. Millar, S.R., I.J. Perry & C.M. Phillips. 2015. Assessing cardiometabolic risk in middle-aged adults using body mass index and waist-height ratio: are two indices better than one? A cross-sectional study. *Diabetol Metab Syndr.* **7**: 73.
45. Phillips, C.M., A.C. Tierney, P. Perez-Martinez, *et al.* 2013. Obesity and body fat classification in the metabolic syndrome: impact on cardiometabolic risk metabotype. *Obesity (Silver Spring, Md.* **21**: E154-161.
46. Wang, B., R. Zhuang, X. Luo, *et al.* 2015. Prevalence of Metabolically Healthy Obese and Metabolically Obese but Normal Weight in Adults Worldwide: A Meta-Analysis. *Hormone and metabolic research = Hormon- und Stoffwechselforschung = Hormones et metabolisme.* **47**: 839-845.
47. Stefan, N., H.U. Haring, F.B. Hu, *et al.* 2013. Metabolically healthy obesity: epidemiology, mechanisms, and clinical implications. *The lancet. Diabetes & endocrinology.* **1**: 152-162.
48. Stefan, N., K. Kantartzis, J. Machann, *et al.* 2008. Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med.* **168**: 1609-1616.
49. Arsenault, B.J., D. Lachance, I. Lemieux, *et al.* 2007. Visceral adipose tissue accumulation, cardiorespiratory fitness, and features of the metabolic syndrome. *Arch Intern Med.* **167**: 1518-1525.
50. Despres, J.P. & I. Lemieux. 2006. Abdominal obesity and metabolic syndrome. *Nature.* **444**: 881-887.
51. Shulman, G.I. 2014. Ectopic fat in insulin resistance, dyslipidemia, and cardiometabolic disease. *The New England journal of medicine.* **371**: 2237-2238.
52. Adiels, M., M.R. Taskinen, C. Packard, *et al.* 2006. Overproduction of large VLDL particles is driven by increased liver fat content in man. *Diabetologia.* **49**: 755-765.
53. Despres, J.P. 1998. The insulin resistance-dyslipidemic syndrome of visceral obesity: effect on patients' risk. *Obes Res.* **6 Suppl 1**: 8S-17S.


54. Hwang, J.H., D.T. Stein, N. Barzilay, *et al.* 2007. Increased intrahepatic triglyceride is associated with peripheral insulin resistance: in vivo MR imaging and spectroscopy studies. *Am J Physiol Endocrinol Metab.* **293**: E1663-1669.
55. Jakobsen, M.U., T. Berentzen, T.I. Sorensen, *et al.* 2007. Abdominal obesity and fatty liver. *Epidemiologic reviews.* **29**: 77-87.
56. Korenblat, K.M., E. Fabbrini, B.S. Mohammed, *et al.* 2008. Liver, muscle, and adipose tissue insulin action is directly related to intrahepatic triglyceride content in obese subjects. *Gastroenterology.* **134**: 1369-1375.
57. Chang, Y., H.S. Jung, J. Cho, *et al.* 2016. Metabolically Healthy Obesity and the Development of Nonalcoholic Fatty Liver Disease. *The American journal of gastroenterology.* **111**: 1133-1140.
58. Heianza, Y., Y. Arase, H. Tsuji, *et al.* 2014. Metabolically healthy obesity, presence or absence of fatty liver, and risk of type 2 diabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 20 (TOPICS 20). *The Journal of clinical endocrinology and metabolism.* **99**: 2952-2960.
59. Jung, C.H., Y.M. Kang, J.E. Jang, *et al.* 2016. Fatty liver index is a risk determinant of incident type 2 diabetes in a metabolically healthy population with obesity. *Obesity (Silver Spring, Md.* **24**: 1373-1379.
60. Sung, K.C., S.C. Cha, J.W. Sung, *et al.* 2014. Metabolically healthy obese subjects are at risk of fatty liver but not of pre-clinical atherosclerosis. *Nutr Metab Cardiovasc Dis.* **24**: 256-262.
61. Zhang, H., Z. Ma, L. Pan, *et al.* 2016. Hepatic fat content is a determinant of metabolic phenotypes and increased carotid intima-media thickness in obese adults. *Scientific reports.* **6**: 21894.
62. McMorrow, A.M., R.M. Connaughton, F.E. Lithander, *et al.* 2015. Adipose tissue dysregulation and metabolic consequences in childhood and adolescent obesity: potential impact of dietary fat quality. *The Proceedings of the Nutrition Society.* **74**: 67-82.
63. Kloting, N., M. Fasshauer, A. Dietrich, *et al.* 2010. Insulin-sensitive obesity. *Am J Physiol Endocrinol Metab.* **299**: E506-515.
64. O'Connell, J., L. Lynch, A. Hogan, *et al.* 2011. Preadipocyte factor-1 is associated with metabolic profile in severe obesity. *The Journal of clinical endocrinology and metabolism.* **96**: E680-684.
65. Phillips, C.M. & I.J. Perry. 2013. Does inflammation determine metabolic health status in obese and nonobese adults? *The Journal of clinical endocrinology and metabolism.* **98**: E1610-1619.
66. Stienstra, R. & N. Stefan. 2013. Tipping the inflammatory balance: inflammasome activation distinguishes metabolically unhealthy from healthy obesity. *Diabetologia.* **56**: 2343-2346.
67. Despres, J.P. 2012. Body Fat Distribution and Risk of Cardiovascular Disease An Update. *Circulation.* **126**: 1301-1313.
68. Linder, K., F. Springer, J. Machann, *et al.* 2014. Relationships of body composition and liver fat content with insulin resistance in obesity-matched adolescents and adults. *Obesity (Silver Spring, Md.* **22**: 1325-1331.
69. Fabbrini, E., F. Magkos, B.S. Mohammed, *et al.* 2009. Intrahepatic fat, not visceral fat, is linked with metabolic complications of obesity. *Proceedings of the National Academy of Sciences of the United States of America.* **106**: 15430-15435.
70. Fabbrini, E., J. Yoshino, M. Yoshino, *et al.* 2015. Metabolically normal obese people are protected from adverse effects following weight gain. *The Journal of clinical investigation.* **125**: 787-795.
71. Hankinson, A.L., M.L. Daviglius, L. Van Horn, *et al.* 2012. Diet composition and activity level of at risk and metabolically healthy obese American adults. *Obesity (Silver Spring, Md.*
72. Kimokoti, R.W., S.E. Judd, J.M. Shikany, *et al.* 2015. Metabolically Healthy Obesity Is Not Associated with Food Intake in White or Black Men. *The Journal of nutrition.* **145**: 2551-2561.

73. Manu, P., C. Ionescu-Tirgoviste, J. Tsang, *et al.* 2012. Dysmetabolic Signals in "Metabolically Healthy" Obesity. *Obes Res Clin Pract.* **6**: e9-e20.
74. Bell, L.K., S. Edwards & J.A. Grieger. 2015. The Relationship between Dietary Patterns and Metabolic Health in a Representative Sample of Adult Australians. *Nutrients.* **7**: 6491-6505.
75. Park, Y.M., S.E. Steck, T.T. Fung, *et al.* 2016. Mediterranean diet and mortality risk in metabolically healthy obese and metabolically unhealthy obese phenotypes. *International journal of obesity (2005).*
76. Janiszewski, P.M. & R. Ross. 2010. Effects of weight loss among metabolically healthy obese men and women. *Diabetes care.* **33**: 1957-1959.
77. Kantartzis, K., J. Machann, F. Schick, *et al.* 2011. Effects of a lifestyle intervention in metabolically benign and malign obesity. *Diabetologia.* **54**: 864-868.
78. Karelis, A.D., V. Messier, M. Brochu, *et al.* 2008. Metabolically healthy but obese women: effect of an energy-restricted diet. *Diabetologia.* **51**: 1752-1754.
79. Shin, M.J., Y.J. Hyun, O.Y. Kim, *et al.* 2006. Weight loss effect on inflammation and LDL oxidation in metabolically healthy but obese (MHO) individuals: low inflammation and LDL oxidation in MHO women. *International journal of obesity (2005).* **30**: 1529-1534.
80. Rondanelli, M., C. Klersy, S. Perna, *et al.* 2015. Effects of two-months balanced diet in metabolically healthy obesity: lipid correlations with gender and BMI-related differences. *Lipids in health and disease.* **14**: 139.
81. Ruiz, J.R., F.B. Ortega & I. Labayen. 2013. A weight loss diet intervention has a similar beneficial effect on both metabolically abnormal obese and metabolically healthy but obese premenopausal women. *Annals of nutrition & metabolism.* **62**: 223-230.
82. Liu, R.H., S. Wharton, A.M. Sharma, *et al.* 2013. Influence of a clinical lifestyle-based weight loss program on the metabolic risk profile of metabolically normal and abnormal obese adults. *Obesity (Silver Spring, Md).*
83. Hogstrom, G., A. Nordstrom & P. Nordstrom. 2014. High aerobic fitness in late adolescence is associated with a reduced risk of myocardial infarction later in life: a nationwide cohort study in men. *European heart journal.* **35**: 3133-3140.
84. Kodama, S., K. Saito, S. Tanaka, *et al.* 2009. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA.* **301**: 2024-2035.
85. Myers, J., P. McAuley, C.J. Lavie, *et al.* 2015. Physical activity and cardiorespiratory fitness as major markers of cardiovascular risk: their independent and interwoven importance to health status. *Prog Cardiovasc Dis.* **57**: 306-314.
86. Sato, M., S. Kodama, A. Sugawara, *et al.* 2009. Physical fitness during adolescence and adult mortality. *Epidemiology (Cambridge, Mass.* **20**: 463-464.
87. Dalleck, L.C., G.P. Van Guilder, T.B. Richardson, *et al.* 2014. A community-based exercise intervention transitions metabolically abnormal obese adults to a metabolically healthy obese phenotype. *Diabetes, metabolic syndrome and obesity : targets and therapy.* **7**: 369-380.
88. Ortega, F.B., D.C. Lee, P.T. Katzmarzyk, *et al.* 2013. The intriguing metabolically healthy but obese phenotype: cardiovascular prognosis and role of fitness. *European heart journal.* **34**: 389-397.
89. Ortega, F.B., C. Cadenas-Sanchez, X. Sui, *et al.* 2015. Role of Fitness in the Metabolically Healthy but Obese Phenotype: A Review and Update. *Prog Cardiovasc Dis.* **58**: 76-86.
90. Berezina, A., O. Belyaeva, O. Berkovich, *et al.* 2015. Prevalence, Risk Factors, and Genetic Traits in Metabolically Healthy and Unhealthy Obese Individuals. *Biomed Res Int.* **2015**: 548734.
91. Ferguson, J.F., C.M. Phillips, A.C. Tierney, *et al.* 2010. Gene-nutrient interactions in the metabolic syndrome: single nucleotide polymorphisms in ADIPOQ and ADIPOR1 interact with plasma saturated fatty acids to modulate insulin resistance. *The American journal of clinical nutrition.* **91**: 794-801.

92. Phillips, C.M., L. Goumidi, S. Bertrais, *et al.* 2010. Leptin receptor polymorphisms interact with polyunsaturated fatty acids to augment risk of insulin resistance and metabolic syndrome in adults. *The Journal of nutrition*. **140**: 238-244.
93. Kvaloy, K., J. Holmen, K. Hveem, *et al.* 2015. Genetic Effects on Longitudinal Changes from Healthy to Adverse Weight and Metabolic Status - The HUNT Study. *PloS one*. **10**: e0139632.
94. Appleton, S.L., C.J. Seaborn, R. Visvanathan, *et al.* 2013. Diabetes and Cardiovascular Disease Outcomes in the Metabolically Healthy Obese Phenotype: A cohort study. *Diabetes care*.
95. Soriguer, F., C. Gutierrez-Repiso, E. Rubio-Martin, *et al.* 2013. Metabolically healthy but obese, a matter of time? Findings from the prospective Pizarra study. *The Journal of clinical endocrinology and metabolism*.
96. Eshtiaghi, R., S. Keihani, F. Hosseinpanah, *et al.* 2015. Natural course of metabolically healthy abdominal obese adults after 10 years of follow-up: the Tehran Lipid and Glucose Study. *International journal of obesity (2005)*. **39**: 514-519.
97. Hamer, M., J.A. Bell, S. Sabia, *et al.* 2015. Stability of metabolically healthy obesity over 8 years: the English Longitudinal Study of Ageing. *European journal of endocrinology / European Federation of Endocrine Societies*. **173**: 703-708.
98. Achilike, I., H.P. Hazuda, S.P. Fowler, *et al.* 2015. Predicting the development of the metabolically healthy obese phenotype. *International journal of obesity (2005)*. **39**: 228-234.
99. Flegal, K.M., B.K. Kit, H. Orpana, *et al.* 2013. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *Jama*. **309**: 71-82.
100. Benckert, M., M. Lilja, S. Soderberg, *et al.* 2015. Improved metabolic health among the obese in six population surveys 1986 to 2009: the Northern Sweden MONICA study. *BMC Obes*. **2**: 7.
101. Hamer, M. & E. Stamatakis. 2012. Metabolically healthy obesity and risk of all-cause and cardiovascular disease mortality. *The Journal of clinical endocrinology and metabolism*. **97**: 2482-2488.
102. Guo, F. & W.T. Garvey. 2015. Cardiometabolic disease risk in metabolically healthy and unhealthy obesity: Stability of metabolic health status in adults. *Obesity (Silver Spring, Md)*.
103. Arnlov, J., J. Sundstrom, E. Ingelsson, *et al.* 2011. Impact of BMI and the metabolic syndrome on the risk of diabetes in middle-aged men. *Diabetes care*. **34**: 61-65.
104. Calori, G., G. Lattuada, L. Piemonti, *et al.* 2011. Prevalence, metabolic features, and prognosis of metabolically healthy obese Italian individuals: the Cremona Study. *Diabetes care*. **34**: 210-215.
105. Durward, C.M., T.J. Hartman & S.M. Nickols-Richardson. 2012. All-cause mortality risk of metabolically healthy obese individuals in NHANES III. *J Obes*. **2012**: 460321.
106. Hinnouho, G.M., S. Czernichow, A. Dugravot, *et al.* 2013. Metabolically healthy obesity and risk of mortality: Does the definition of metabolic health matter? *Diabetes care*.
107. Kuk, J.L. & C.I. Ardern. 2009. Are metabolically normal but obese individuals at lower risk for all-cause mortality? *Diabetes care*. **32**: 2297-2299.
108. van der, A.D., A.C. Nooyens, F.J. van Duijnhoven, *et al.* 2013. All-cause mortality risk of metabolically healthy abdominal obese individuals: The EPIC-MORGEN study. *Obesity (Silver Spring, Md)*.
109. Bradshaw, P.T. & J. Stevens. 2015. Invited commentary: limitations and usefulness of the metabolically healthy obesity phenotype. *American journal of epidemiology*. **182**: 742-744.
110. Phillips, C.M. 2016. Metabolically Healthy Obesity: Personalised and Public Health Implications. *Trends Endocrinol Metab*. **27**: 189-191.
111. Puri, R. 2014. Is it finally time to dispel the concept of metabolically-healthy obesity? *J Am Coll Cardiol*. **63**: 2687-2688.
112. Rey-Lopez, J.P., L.F. de Rezende, T.H. de Sa, *et al.* 2015. Is the metabolically healthy obesity phenotype an irrelevant artifact for public health? *American journal of epidemiology*. **182**: 737-741.

113. Hinnouho, G.M., S. Czernichow, A. Dugravot, *et al.* 2015. Metabolically healthy obesity and the risk of cardiovascular disease and type 2 diabetes: the Whitehall II cohort study. *European heart journal.* **36**: 551-559.
114. Khan, U.I., D. Wang, R.C. Thurston, *et al.* 2011. Burden of subclinical cardiovascular disease in "metabolically benign" and "at-risk" overweight and obese women: the Study of Women's Health Across the Nation (SWAN). *Atherosclerosis.* **217**: 179-186.
115. Kwon, B.J., D.W. Kim, S.H. Her, *et al.* 2013. Metabolically obese status with normal weight is associated with both the prevalence and severity of angiographic coronary artery disease. *Metabolism: clinical and experimental.* **62**: 952-960.
116. McPherson K, K.P., Butland B, Jebb S, Thomas S *et al.* . 2007. "Tackling Obesities: Future Choices-Project Report. 2nd Edition. ". In. G.O.f.S.A.h.w.b.g.u.a.f.d.o.p.A. 02/01/2012, Ed.
117. Bosello, O., M.P. Donataccio & M. Cuzzolaro. 2016. Obesity or obesities? Controversies on the association between body mass index and premature mortality. *Eat Weight Disord.* **21**: 165-174.
118. Munoz-Garach, A., I. Cornejo-Pareja & F.J. Tinahones. 2016. Does Metabolically Healthy Obesity Exist? *Nutrients.* **8**.
119. Payab, M., S. Hasani-Ranjbar & B. Larijani. 2014. Whether all obese subjects both in metabolic groups and non-metabolic groups should be treated or not. *J Diabetes Metab Disord.* **13**: 21.
120. de Rooij, B.H., J.D. van der Berg, C.J. van der Kallen, *et al.* 2016. Physical Activity and Sedentary Behavior in Metabolically Healthy versus Unhealthy Obese and Non-Obese Individuals - The Maastricht Study. *PloS one.* **11**: e0154358.
121. Schroder, H., R. Ramos, J.M. Baena-Diez, *et al.* 2014. Determinants of the transition from a cardiometabolic normal to abnormal overweight/obese phenotype in a Spanish population. *European journal of nutrition.* **53**: 1345-1353.
122. Doumatey, A.P., J. Zhou, M. Zhou, *et al.* 2016. Proinflammatory and lipid biomarkers mediate metabolically healthy obesity: A proteomics study. *Obesity (Silver Spring, Md.* **24**: 1257-1265.
123. Phillips, C.M. & I.J. Perry. 2015. Lipoprotein particle subclass profiles among metabolically healthy and unhealthy obese and non-obese adults: Does size matter? *Atherosclerosis.* **242**: 399-406.
124. Garvey, W.T., A.J. Garber, J.I. Mechanick, *et al.* 2014. American association of clinical endocrinologists and american college of endocrinology position statement on the 2014 advanced framework for a new diagnosis of obesity as a chronic disease. *Endocr Pract.* **20**: 977-989.
125. Garvey, W.T., J.I. Mechanick, E.M. Brett, *et al.* 2016. American Association of Clinical Endocrinologists and American College of Endocrinology Clinical Practice Guidelines for Comprehensive Medical Care of Patients with Obesity - Executive Summary. *Endocr Pract.*
126. Phillips, C.M. & I.J. Perry. 2015. Depressive symptoms, anxiety and well-being among metabolic health obese subtypes. *Psychoneuroendocrinology.* **62**: 47-53.

Figure 1



In this model it is hypothesized that the body’s coping mechanism to a positive energy balance, arising from a combination of excessive caloric intake, increased levels of sedentary behaviour and reduced physical activity, may determine an individual’s predisposition to MHO or MUO. The pathway to MUO may be characterised by dysfunctional adipose tissue, increased immune cell infiltration and reduced capacity of subcutaneous adipose tissue to expand leading to increased ectopic fat deposition resulting in lipotoxicity, insulin resistance in peripheral tissues and a range of metabolic derangements. Adapted from McMorrow *et al.*,⁶².