

Title	Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two o-hydroxyaldehyde groups - The importance of topology
Authors	Pareras, Gerard;Szczepanik, Dariusz Wojciech;Duran, Miquel;Solà, Miquel;Simon, Silvia
Publication date	2019-11-06
Original Citation	Pareras, G., Szczepanik, D. W., Duran, M., Solà, M. and Simon, S. (2019) 'Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two o-hydroxyaldehyde groups - The importance of topology', Journal of Organic Chemistry, 84(23), pp. 15538-15548. doi: 10.1021/acs.joc.9b02526
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1021/acs.joc.9b02526
Rights	© 2019, American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Journal of Organic Chemistry, copyright © American Chemical Society, after peer review and technical editing by the publisher. To access the final edited and published work see <a href="https://pubs.acs.org/doi/abs/10.1021/acs.joc.9b02526">https://pubs.acs.org/doi/abs/10.1021/acs.joc.9b02526</a>
Download date	2023-10-01 15:12:26
Item downloaded from	<a href="https://hdl.handle.net/10468/9436">https://hdl.handle.net/10468/9436</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

# SUPPORTING INFORMATION

## Tuning the strength of the resonance-assisted hydrogen bond in acenes and phenacenes with two *o*-hydroxyaldehyde groups. The importance of topology

Gerard Pareras<sup>a,‡</sup>, Dariusz W. Szczepanik<sup>b,‡</sup>, Miquel Duran<sup>b</sup>, Miquel Solà<sup>b,\*</sup> and Sílvia Simon<sup>b,\*</sup>

<sup>a</sup> Department of Chemistry, University College Cork, College Rd, Cork, Ireland


<sup>b</sup> Institut de Química Computacional i Catàlisi and Departament de Química, Universitat de Girona, C/ Maria Aurèlia Capmany 69, 17003 Girona, Catalonia, Spain

‡ These two authors contributed equally to this work.


Corresponding authors: [silvia.simon@udg.edu](mailto:silvia.simon@udg.edu), [miquel.sola@udg.edu](mailto:miquel.sola@udg.edu)

**Fig. S1** – Linear correlation between PDI for the *ipso* and *quasi*-rings for triphenylene-like compound. Grey for **KM-Ln A'** *quasi*-ring, yellow for **KM-Ln B'** *quasi*-ring, red for **KP-Lnb** and blue for **KP-Lna**, being *n* the number of added benzene rings (see Fig. 6 in text) **S2**

**Fig. S2** – Linear correlation between aromaticity of the *ipso*-ring and RAHB bond length for the triphenylene-like compound. Grey for **KM-Ln A'** *quasi*-ring, yellow for **KM-Ln B'** *quasi*-ring, red for **KP-Lnb** and blue for **KP-Lna**, being *n* the number of added benzene rings (see Fig. 6 in text). ..... **S2**



**Fig. S1** – Linear correlation between PDI for the *ipso* and *quasi*-rings for triphenylene-like compound. Grey for **KM-Ln A'** *quasi*-ring, yellow for **KM-Ln B'** *quasi*-ring, red for **KP-Lnb** and blue for **KP-Lna**, being n the number of added benzene rings (see Fig. 6 in text).



**Fig. S2** – Linear correlation between aromaticity of the *ipso*-ring and RAHB bond length for the triphenylene-like compound. Grey for **KM-Ln A'** *quasi*-ring, yellow for **KM-Ln B'** *quasi*-ring, red for **KP-Lnb** and blue for **KP-Lna**, being n the number of added benzene rings (see Fig. 6 in text).