


|                             |   |
|-----------------------------|---|
| Title                       | Breast milk-derived human milk oligosaccharides promote Bifidobacterium interactions within a single ecosystem  |
| Authors                     | Lawson, Melissa A. E.;O'Neill, Ian J.;Kujawska, Magdalena;Gowrinadh Javvadi, Sree;Wijeyesekera, Anisha;Flegg, Zak;Chalklen, Lisa;Hall, Lindsay J.   |
| Publication date            | 2019-11-18  |
| Original Citation           | Lawson, M. A. E., O'Neill, I. J., Kujawska, M., Gowrinadh Javvadi, S., Wijeyesekera, A., Flegg, Z., Chalklen, L. and Hall, L. J. (2019) 'Breast milk-derived human milk oligosaccharides promote Bifidobacterium interactions within a single ecosystem', The ISME Journal. (14pp.) doi: 10.1038/s41396-019-0553-2  |
| Type of publication         | Article (peer-reviewed)   |
| Link to publisher's version | 10.1038/s41396-019-0553-2   |
| Rights                      | © The Author(s) 2019. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> . - <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> |
| Download date               | 2023-03-29 23:28:22   |
| Item downloaded from        | <a href="http://hdl.handle.net/10468/9357">http://hdl.handle.net/10468/9357</a>   |



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh



**Supplementary figure 2:** Bifidobacterial survival after exposure to intestinal environmental stressors. (A) Bacterial survival in MRS supplemented with 0.3% bile salts was measured after 48h growth and compared to parallel cultures grown in MRS anaerobically at 37°C. Data shown from five independent experiments; n.d. denotes measurement not detectable. Blank/media only values are subtracted from all points in each experiment, all negative values are represented as zero. Isolates from Baby V1 are represented as black bars, V2 are grey bars, V3 are dotted bars, and the tested type and control strains are shown as white bars. Error bars represent SEM. (B) Bacterial growth in MRS was measured for each strain after 30h growth in an aerobic environment and compared to parallel samples grown anaerobically. Measurements shown are representative of three independent experimental repeats. (C) Ability to withstand acid shock for 4 h at a pH2. Following acid shock, strains were incubated for 48 h in an anaerobic cabinet at 37°C. Growth (optical density) was then measured. Data shown is mean ± standard deviation for 3 independent experiments