

Title	Efficient determination of non-steroidal anti-inflammatory drugs by micellar electrokinetic chromatography in wastewater
Authors	Alatawi, Hanan;Hogan, Anna;Albalawi, Ibtihaj;Alsefri, Samia;Moore, Eric
Publication date	2023-01
Original Citation	Alatawi, H., Hogan, A., Albalawi, I., Alsefri, S. and Moore, E. (2023) 'Efficient determination of non-steroidal anti-inflammatory drugs by micellar electrokinetic chromatography in wastewater', Analytical Methods, (8pp). doi: 10.1039/D2AY01807A
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1039/d2ay01807a
Rights	© The Royal Society of Chemistry 2023. - https://creativecommons.org/licenses/by-nc/3.0/
Download date	2023-10-01 23:01:07
Item downloaded from	https://hdl.handle.net/10468/14239



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

Supplementary Information for Analytical methods

Efficient determination of non-steroidal anti-inflammatory drugs by micellar electrokinetic chromatography in wastewater

Hanan Alatawi¹

Anna Hogan¹

Ibtihaj Albalawi¹

Samia Alsefri¹

Eric Moore^{1,2}

¹ School of Chemistry, University College Cork, Cork, Ireland

² Tyndall National Institute, Cork, Ireland

Correspondence to:

Eric Moore

School of Chemistry, University College Cork, Cork, Ireland.

Email: e.moore@ucc.ie

Supplementary figures




Figure. S1 Electropherogram of CZE separation: Peak: 1- PAC, 2- DIC, 3-KET, 4- IB, 5- ASA; using different BGE: (A) 20 mM acetate BGE at pH 5.5 , (B) 20 mM phosphate BGE at pH 9, (c) 20 mM borate BGE at pH 9.




Figure. S2 MEKC electropherograms show the effect of SDS concentration on the separation of NSAIDs (A) 50 mM, (B) 70 mM, (C) 90 mM, (D) 110 Mm; Peak: 1- PAC, 2- ASA, 3-KET, 4- IB, 5- DIC at concentration of 750 $\mu\text{g/L}$ each; Capillary: 50 μm id 31.5 cm (effective length) 40 cm (total length); BGE: 15 mM borate buffer, pH 9; separation voltage: 15 kV.




Figure. S3 The effect of injection size on the efficiency




Figure. S4 MEKC electropherograms show the effect of voltage on the separation of NSAIDs (A) 10, (B) 15, (c) 20, (D) 25 kV; Peak: 1- PAC, 2- ASA, 3-KET, 4- IB, 5- DIC at a concentration of 750 $\mu\text{g/L}$ each; Capillary: 50 μm id 31.5 cm (effective length) 40 cm (total length); BGE: 15 mM borate buffer, pH 9, 90 mM SDS, 10% methanol; separation voltage: 15 kV; hydrodynamic injection: 10 mbar/5s.

Table S1. Structures of the investigated compounds






Analytes	Chemical structure	MW	pKa
ASA		180.16	3.15
DIC		318.13	4.15
IB		206.28	4.60
KET		254.28	4.0
PAC		151.16	9.5

Table S2. The recoveries of investigated compounds in (100 µg/L) spiked wastewater samples

Analyte	Recovery (RSD%)	
	C18	Oasis HLB
	Wastewater sample	Wastewater sample
IB	83.8 (13.2%)	40 (19%)
KET	91.1 (11%)	28.1 (24%)
DIC	118.1 (15.5%)	22.7 (28.5%)
ASA	36.5 (14.7%)	65.2 (11.6%)
PAC	-	122.4 (9.3%)