

Title	Evolutionary stasis of a heritable morphological trait in a wild fish population despite apparent directional selection
Authors	O'Sullivan, Ronan J.;Aykanat, Tutku;Johnston, Susan E.;Kane, Adam;Poole, Russell;Rogan, Ger;Prodöhl, Paulo A.;Primmer, Craig R.;McGinnity, Philip;Reed, Thomas E.
Publication date	2019-06-11
Original Citation	O'Sullivan, R. J., Aykanat, T., Johnston, S. E., Kane, A., Poole, R., Rogan, G., Prodöhl, P. A., Primmer, C. R., McGinnity, P. and Reed, T. E. (2019) 'Evolutionary stasis of a heritable morphological trait in a wild fish population despite apparent directional selection', <i>Ecology and Evolution</i> , 9(12), pp. 7096-7111. doi: 10.1002/ece3.5274
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://onlinelibrary.wiley.com/doi/full/10.1002/ece3.5274 - 10.1002/ece3.5274
Rights	© 2019 The Authors. <i>Ecology and Evolution</i> published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited. - https://creativecommons.org/licenses/by/4.0/
Download date	2024-09-10 17:10:10
Item downloaded from	https://hdl.handle.net/10468/11527




Figure S4: Stylized four-year lifecycle of Burritshoole Atlantic salmon. Adult fish spawn in freshwater, G1. G1's offspring, G2, typically spend 3 years in freshwater (0+, 1+, 2+) before leaving the river (smoltification) during their third year of life, and undergoing an oceanic feeding stage (2+/3+). G2 typically return to freshwater to spawn during their fourth year of life (3+), giving rise to the next generation, G3.